

Topology of random real hypersurfaces

Jean-Yves Welschinger

September 8, 2014

Abstract:

These are notes of the mini-course I gave during the CIMPA summer school at Villa de Leyva, Colombia, in July 2014. The subject was my joint work with Damien Gayet on the topology of random real hypersurfaces, restricting myself to the case of projective spaces and focusing on our lower estimates. Namely, we estimate from (above and) below the mathematical expectation of all Betti numbers of degree d random real projective hypersurfaces. For any closed connected hypersurface Σ of \mathbb{R}^n , we actually estimate from below the mathematical expectation of the number of connected components of these degree d random real projective hypersurfaces which are diffeomorphic to Σ .

1 Random real polynomials

1.1 In one variable

Let $P \in \mathbb{R}_d[X]$ be a polynomial in one variable, of degree d and with real coefficients. Let $V_P = \{x \in \mathbb{R} \mid P(x) = 0\}$ be the set of its real roots. Everybody knows the following

Theorem 1.1 *Let $P \in \mathbb{R}_d[X]$, then $0 \leq \#V_P \leq d$ and moreover $\#V_P \equiv d \pmod{2}$ provided P is generic enough.*

This Theorem 1.1 already raises a question which is going to be the main theme of this course.

Question 1: What is the typical number of roots of P , choosing P at random?

The mathematical expectation of this number of real roots reads as the average

$$\mathbb{E}(\#V_P) = \int_{\mathbb{R}_d[X]} (\#V_P) d\mu(P),$$

where μ denotes some probability measure on $\mathbb{R}_d[X]$.

First answer:

A first answer to Question 1 has been given by M. Kac in the 40's

Theorem 1.2 (M. Kac, 1943, [11])

$$\mathbb{E}(\#V_P) \sim_{d \rightarrow +\infty} \frac{2}{\pi} \log(d).$$

In order to provide this answer, Kac did consider that the space $\mathbb{R}_d[X]$ of polynomials is Euclidean, a canonical orthonormal basis being given by the monomials $1, X, X^2, \dots, X^d$. Now, since this space is Euclidean, it carries a canonical probability measure, the Gaussian measure associated to its scalar product. The latter reads

$$d\mu(P) = \frac{1}{\sqrt{\pi}^{d+1}} \exp(-\|P\|^2) dP,$$

where $d+1$ corresponds to the dimension of $\mathbb{R}_d[X]$ and dP to its Lebesgue measure which is associated to the scalar product but has infinite volume. This Gaussian measure is thus the Lebesgue measure weighted with some exponential which reduces its total volume to one. It has the great properties to be a product measure which is invariant under the orthogonal group.

Second answer:

A second answer to Question 1 has been given by E. Kostlan.

Theorem 1.3 (Kostlan, Shub-Smale 1993, [13], [21]) *For every $d > 0$,*

$$\mathbb{E}(\#V_P) = \sqrt{d}.$$

In order to provide this answer, E. Kostlan also did equip the space $\mathbb{R}_d[X]$ of polynomials with some Gaussian measure, but associated to a different scalar product. For this new scalar product, an orthonormal basis is given by the monomials $\sqrt{\binom{d}{k}} X^k$, $0 \leq k \leq d$. This scalar product turns out to be more natural geometrically. I will give a geometric definition in §2.5, but let me already point out a nice property.

The space $\mathbb{R}_d[X]$ is well known to be isomorphic to the space $\mathbb{R}_d^{\text{hom}}[X, Y]$ of homogeneous polynomials of degree d in two variables and real coefficients. This isomorphism reads $X^k \in \mathbb{R}_d[X] \mapsto X^k Y^{d-k} \in \mathbb{R}_d^{\text{hom}}[X, Y]$. If we push forward the new scalar product under this isomorphism, then we get one which is invariant under the action of the orthogonal group of the plane, by composition on the right. That is, for every $Q \in \mathbb{R}_d^{\text{hom}}[X, Y]$ and every $h \in O_2(\mathbb{R})$, $\|Q\| = \|Q \circ h^{-1}\|$.

Let me explain the proof of Kostlan, which also recovers the result of Kac.

Proof: (see [4])

Let us fix the isomorphism $(a_0, \dots, a_d) \in \mathbb{R}^{d+1} \mapsto \sum_{i=0}^d a_i X^i \in \mathbb{R}_d[X]$ and focus on two objects of \mathbb{R}^{d+1} . First, the unit sphere S^d and for every $\underline{a} = (a_0, \dots, a_d) \in S^d$, let us denote by $\lambda_{\underline{a}}$ the linear form $(y_0, \dots, y_d) \in \mathbb{R}^{d+1} \mapsto \sum_{i=0}^d a_i y_i \in \mathbb{R}$.

Secondly, let us consider the curve $\tilde{\gamma} : t \in \mathbb{R} \mapsto (1, t, \dots, t^d) \in \mathbb{R}^{d+1}$ in the case of Kac, or $\tilde{\gamma} : t \in \mathbb{R} \mapsto (1, \sqrt{\binom{d}{1}}t, \dots, \sqrt{\binom{d}{d}}t^d) \in \mathbb{R}^{d+1}$ in the case of Kostlan.

If $P = \sum_{i=0}^d a_i X^i \in \mathbb{R}_d[X]$, then, as a function on the real line, $P = \lambda_{\underline{a}} \circ \tilde{\gamma}$ (in the case of Kac), so that $V_P \cong \ker \lambda_{\underline{a}} \cap \text{Im}(\tilde{\gamma})$.

The observation of Kostlan is then the following.

Theorem 1.4 ([4])

$$\mathbb{E}(\#V_P) = \frac{1}{\pi} \text{length}(\gamma), \text{ where } \gamma : t \in \mathbb{R} \mapsto \frac{\tilde{\gamma}(t)}{\|\tilde{\gamma}(t)\|} \in S^d.$$

Proof : (see [4])

This is Crofton's formula, the length of a curve is the average of the number of intersection points with the hyperplanes. Note that the formula is obvious when the curve is a closed geodesic on the sphere. This formula follows from the fact that it also holds true for a piece of such a geodesic, since every smooth curve can then be approximated by some piecewise geodesic curve. \square

The end of the proof of Theorem 1.4 is just a computation of the length of the curve γ , which gives $\text{length}(\gamma) \sim_{d \rightarrow +\infty} 2 \log(d)$ in the case of Kac (a bit tough) and $\text{length}(\gamma) = \pi\sqrt{d}$ in the case of Kostlan (easy). \square

1.2 In several variables

What about polynomials in several variables?

If $P \in \mathbb{R}_d[X_1, \dots, X_n]$ is a polynomial in n variables, degree d and real coefficients, then $V_P = \{x \in \mathbb{R}^n \mid P(x) = 0\}$ is no more a finite set, but rather an affine real algebraic hypersurface. It is not compact in general, but has a standard compactification. Namely, this space of polynomial is again canonically isomorphic to the space $\mathbb{R}_d^{\text{hom}}[X_0, \dots, X_n]$ of homogeneous polynomials of degree d , $n+1$ variables and real coefficients. This isomorphism reads $X_1^{\alpha_1} \dots X_n^{\alpha_n} \in \mathbb{R}_d[X_1, \dots, X_n] \mapsto X_0^{d-\alpha_1-\dots-\alpha_n} X_1^{\alpha_1} \dots X_n^{\alpha_n} \in \mathbb{R}_d^{\text{hom}}[X_0, \dots, X_n]$ and if $Q \in \mathbb{R}_d^{\text{hom}}[X_0, \dots, X_n] \setminus \{0\}$, then $V_Q = \{x \in \mathbb{R}P^n \mid Q(x) = 0\}$ is a compact hypersurface, smooth for generic polynomials and which then contains V_P as a dense subset. I will come back to projective spaces in §2.1.

Again, the topology of V_Q depends on the choice of Q , as in one variable. For example, in degree $d = 2$ and $n = 3$ variables, V_Q is a quadric surface which may be empty, homeomorphic to a sphere in the case of the ellipsoid or to a torus in the case of the hyperboloid. If we denote, for every $i \in \{0, \dots, n-1\}$, by $b_i(V_Q; \mathbb{Z}/2\mathbb{Z}) = \dim H_i(V_Q; \mathbb{Z}/2\mathbb{Z})$ the i -th Betti number with $\mathbb{Z}/2\mathbb{Z}$ coefficients of V_Q , then

Theorem 1.5 (Smith-Thom's inequality, 1965, [22])

$$0 \leq \sum_{i=0}^{n-1} b_i(V_Q; \mathbb{Z}/2\mathbb{Z}) \leq \sum_{i=0}^{2n-2} b_i(V_Q^{\mathbb{C}}; \mathbb{Z}/2\mathbb{Z}) = d^n + o(d^n).$$

Moreover, $\sum_{i=0}^{n-1} b_i(V_Q; \mathbb{Z}/2\mathbb{Z}) \equiv \sum_{i=0}^{2n-2} b_i(V_Q^{\mathbb{C}}; \mathbb{Z}/2\mathbb{Z}) \pmod{2}$, provided $Q \in \mathbb{R}_d^{\text{hom}}[X_0, \dots, X_n]$ is generic enough for V_Q to be smooth.

In Theorem 1.5, $V_Q^{\mathbb{C}} = \{x \in \mathbb{C}P^n \mid Q(x) = 0\}$ denotes the set of complex roots of Q in the complex projective space. This Theorem 1.5 extends Theorem 1.1, which corresponds to the case $n = 1$. The case $n = 2$ was also previously known as the (famous in real algebraic geometry) Harnack-Klein's inequality, see [9], [12].

Again, this raises the question

Question 2: What is the typical topology of V_Q , choosing Q at random in $\mathbb{R}_d^{\text{hom}}[X_0, \dots, X_n]$? e.g. which Betti numbers to expect?

Let me give a formulation of our joint results with Damien Gayet.

For every $i \in \{0, \dots, n-1\}$, let me set

$$\begin{aligned} b_i(V_Q; \mathbb{R}) &= \dim H_i(V_Q; \mathbb{R}), \\ m_i(V_Q) &= \inf_{f \text{ Morse on } V_Q} \#\text{Crit}_i(f). \end{aligned}$$

Here, $\text{Crit}_i(f)$ denotes the number of critical points of index i of the Morse function f . Recall that a real function of class C^2 is said to be Morse if and only if all of its critical points are non-degenerated. This means that the Hessian of this function at all of its critical points is a non-degenerated quadratic form. The index of such a quadratic form is then the maximal dimension of a linear subspace of the tangent space at the critical point where it restricts to a negative definite one, see [17]. The latter is called the index of the critical point.

It follows from Morse theory that $b_i(V_Q; \mathbb{R}) \leq b_i(V_Q; \mathbb{Z}/2\mathbb{Z}) \leq m_i(V_Q)$, see [17].

The mathematical expectations for these Betti or Morse numbers read as the averages

$$\begin{aligned} \mathbb{E}(b_i) &= \int_{\mathbb{R}_d^{\text{hom}}[X_0, \dots, X_n]} b_i(V_Q; \mathbb{R}) d\mu(Q), \\ \mathbb{E}(m_i) &= \int_{\mathbb{R}_d^{\text{hom}}[X_0, \dots, X_n]} m_i(V_Q) d\mu(Q). \end{aligned}$$

The probability measure μ we consider extends the one considered in Theorem 1.3. It is the Gaussian measure associated to the scalar product for which the monomials $\sqrt{\frac{(d+n)!}{n!\alpha_0! \dots \alpha_n!}} X_0^{\alpha_0} \dots X_n^{\alpha_n}$, $\alpha_0 + \dots + \alpha_n = d$, define an orthonormal basis. Again, the action of the orthogonal group of the $(n+1)$ -dimensional Euclidean space by composition on the right preserves this scalar product. That is, for every $Q \in \mathbb{R}_d^{\text{hom}}[X_0, \dots, X_n]$ and every $h \in O_{n+1}(\mathbb{R})$, $\|Q\| = \|Q \circ h^{-1}\|$. Let me finally observe that the coefficient $(d+n)!$ instead of $d!$ in the numerator of the monomials has only the effect to rescale the scalar product and does not affect the results. We will see in §2.5 how this scalar product shows up.

Theorem 1.6 (joint with Damien Gayet, [5], [6]) *There exist (universal) constants c_i^+, c_i^- such that*

$$c_i^- \leq \liminf_{d \rightarrow +\infty} \frac{\mathbb{E}(b_i)}{\sqrt{d^n} \text{Vol}_{FS} \mathbb{R}P^n} \leq \limsup_{d \rightarrow +\infty} \frac{\mathbb{E}(m_i)}{\sqrt{d^n} \text{Vol}_{FS} \mathbb{R}P^n} \leq c_i^+.$$

Unfortunately, by lack of time, I will only explain the proof of the lower estimates given by Theorem 1.6 in this course. The term $\text{Vol}_{\text{FS}}\mathbb{R}P^n$ denotes the total volume of the real projective space for the Fubini-Study metric, see §2.5. Though it is some constant, I distinguish it from c_i^\pm . In fact, in [5], [6], we not only prove Theorem 1.6 for projective spaces, but for any smooth real projective manifold. The term $\text{Vol}_{\text{FS}}\mathbb{R}P^n$ has then to be replaced by the total Kählerian volume of the real locus of the manifold, for the Kähler metric induced by the curvature form of a metric with positive curvature chosen on some ample real line bundle, the tensor powers of which we consider random sections. The constants c_i^+, c_i^- are, they, unchanged and only depend on i and n , see §1.3.

1.3 The universal constants c_i^+, c_i^-

Let me tell you more about these universal constants c_i^+, c_i^- .

The constant c_i^+ is related to random symmetric matrices. Namely $c_i^+ = \frac{1}{\sqrt{\pi}}e_{\mathbb{R}}(i, n - 1 - i)$, where

$$e_{\mathbb{R}}(i, n - 1 - i) = \int_{\text{Sym}(i, n-1-i; \mathbb{R})} |\det(A)| d\mu(A). \quad (1)$$

Here, $\text{Sym}(i, n - 1 - i; \mathbb{R})$ denotes the open cone of non-degenerated symmetric matrices of size $(n - 1) \times (n - 1)$, signature $(i, n - 1 - i)$ and real coefficients. It is included in the vector space $\text{Sym}(n - 1; \mathbb{R})$ of real symmetric matrices of size $(n - 1) \times (n - 1)$. The latter is Euclidean, equipped with the scalar product $(A, B) \in \text{Sym}(n - 1; \mathbb{R})^2 \mapsto \frac{1}{2}\text{tr}(AB) \in \mathbb{R}$, see [16]. So again this space inherits some Gaussian measure μ , which is the one we consider in the integral (1).

In particular,

$$\sum_{i=0}^{n-1} c_i^+ = \frac{1}{\sqrt{\pi}} \mathbb{E}(|\det(A)|) = \frac{1}{\sqrt{\pi}} \int_{\text{Sym}(n-1; \mathbb{R})} |\det(A)| d\mu(A).$$

Theorem 1.7 (joint with Damien Gayet, [5]) 1. $\sum_{i=0}^{n-1} c_i^+ \sim_{n \rightarrow +\infty} \frac{2\sqrt{2}}{\pi} \Gamma(\frac{n+1}{2})$, where Γ denotes Euler's function.

2. For every $\alpha \in [0, \frac{1}{2}[$, there exists $c_\alpha > 0$ such that for n large enough $\sum_{i=0}^{\lfloor \alpha n \rfloor} c_i^+ \leq \exp(-c_\alpha n^2)$. \square

Note that the first part of Theorem 1.7 was known for n even, see [16] and [5], while the second part quickly follows from some large deviation estimates established in [1].

For instance, for $i = 0$, $b_i(V_Q; \mathbb{R}) = b_0(V_Q; \mathbb{R})$ denotes the number of connected components of V_Q , Theorem 1.6 estimates the expected number of connected components of V_Q and $c_i^+ = c_0^+$ provides the upper estimate. This constant more than exponentially decreases as the dimension n grows to $+\infty$.

As for the constant c_i^- , we set

$\mathcal{H}_n = \{\text{closed connected hypersurfaces of } \mathbb{R}^n\}/\text{diffeomorphisms}$.

For every $[\Sigma] \in \mathcal{H}_n$, we set $b_i(\Sigma; \mathbb{R}) = \dim H_i(\Sigma; \mathbb{R})$ and associate some positive constant $c_{[\Sigma]}$, see §2.7. This constant $c_{[\Sigma]}$ is defined via some quantitative transversality, but turns out at the end to bound from below the expected number of connected components of V_Q that are diffeomorphic to Σ , which is what we actually estimate from below, see §2.7. Then, c_i^- is the infinite serie

$$c_i^- = \sum_{[\Sigma] \in \mathcal{H}_n} c_{[\Sigma]} b_i(\Sigma; \mathbb{R}).$$

This serie converges since it is bounded from above by c_i^+ .

Proposition 1.8 (joint with Damien Gayet, [6]) *For every $i \in \{0, \dots, n-1\}$, $c_i^- \geq c_{[S^i \times S^{n-1-i}]} \geq \exp(-\exp(5n+69))$. \square*

Indeed, the product of the i -dimensional unit sphere with the $(n-1-i)$ -dimensional unit sphere turns out to embed as a closed connected hypersurface of \mathbb{R}^n . The i -th Betti number of this hypersurface is one and we will see in §2.7 that the constant $c_{[\Sigma]}$ is actually explicit, so that it can be estimated for this product of spheres, see [6].

2 The lower estimates

2.1 Projective spaces

Recall that the n -dimensional projective space is by definition the space of one-dimensional linear subspaces of the affine $(n+1)$ -dimensional space. That is,

$$\begin{aligned} \mathbb{C}P^n &= \{\text{space of lines in } \mathbb{C}^{n+1}\} \\ &= \mathbb{C}^{n+1} \setminus \{0\} / x \sim \lambda x, \forall \lambda \in \mathbb{C}^*, \end{aligned}$$

and likewise,

$$\begin{aligned} \mathbb{R}P^n &= \{\text{space of lines in } \mathbb{R}^{n+1}\} \\ &= \mathbb{R}^{n+1} \setminus \{0\} / x \sim \lambda x, \forall \lambda \in \mathbb{R}^*. \end{aligned}$$

The points in $\mathbb{C}P^n$ are represented by their homogeneous coordinates $[x_0 : \dots : x_n]$, where $x_0, \dots, x_n \in \mathbb{C}$ do not all vanish, being understood that for every $\lambda \in \mathbb{C}^*$, $[x_0 : \dots : x_n] = [\lambda x_0 : \dots : \lambda x_n]$.

These complex projective spaces are smooth compact complex manifolds without boundary. They are covered by $n+1$ standard affine charts. Namely, for every $i \in \{0, \dots, n\}$, set $U_i = \{[x_0 : \dots : x_n] \in \mathbb{C}P^n \mid x_i \neq 0\}$. This dense open subset U_i corresponds to the lines of \mathbb{C}^{n+1} that are not contained in the hyperplane $\{(x_0, \dots, x_n) \in \mathbb{C}^{n+1} \mid x_i = 0\}$. Every such line intersects the affine hyperplane $\{(x_0, \dots, x_n) \in \mathbb{C}^{n+1} \mid x_i = 1\}$ at exactly one point, defining the chart

$$\phi_i : [x_0 : \dots : x_n] \in U_i \mapsto \left(\frac{x_0}{x_i}, \dots, \frac{x_{i-1}}{x_i}, \frac{x_{i+1}}{x_i}, \dots, \frac{x_n}{x_i} \right) \in \mathbb{C}^n.$$

2.2 Line bundles

The projective space $\mathbb{C}P^n$ is the space of lines of \mathbb{C}^{n+1} , so that every point $x = [x_0 : \dots : x_n]$ of $\mathbb{C}P^n$ represents a complex line $\gamma_x \subset \mathbb{C}^{n+1}$, the line generated by (x_0, \dots, x_n) in \mathbb{C}^{n+1} . The collection of all lines γ_x , $x \in \mathbb{C}P^n$, defines what is called a holomorphic line bundle γ over $\mathbb{C}P^n$. It is in particular a complex manifold equipped with a holomorphic submersion onto the base $\mathbb{C}P^n$, see [2], [7]. Since all these lines are included in \mathbb{C}^{n+1} , the tautological line bundle γ is a subline bundle of the trivial vector bundle $\mathbb{C}P^n \times \mathbb{C}^{n+1} \rightarrow \mathbb{C}P^n$ of rank $n + 1$.

Now, every vector space comes with its dual space, the space of linear forms over it. This defines the dual bundle $\gamma^* = \{\text{linear forms on } \gamma\} \rightarrow \mathbb{C}P^n$. Likewise, for every $d > 0$, I denote by γ_d^* the space of homogeneous forms of degree d on γ , so that $\gamma_1^* = \gamma^*$. Again, all these define holomorphic line bundles over $\mathbb{C}P^n$. Note that another standard notation for these bundle is $\gamma = \mathcal{O}_{\mathbb{C}P^n}(-1)$, $\gamma_d^* = \mathcal{O}_{\mathbb{C}P^n}(d)$.

We denote by $H^0(\mathbb{C}P^n; \gamma_d^*)$ the space of global holomorphic sections of the bundle γ_d^* , that is the space of holomorphic maps $s : \mathbb{C}P^n \rightarrow \gamma_d^*$ such that $\pi \circ s = \text{id}_{\mathbb{C}P^n}$, where $\pi : \gamma_d^* \rightarrow \mathbb{C}P^n$ denotes the tautological projection. Hence, for every point $x \in \mathbb{C}P^n$, $s(x)$ denotes a homogeneous form of degree d on the complex line γ_x .

Now, complex homogeneous polynomials of degree d in $n + 1$ variables define homogeneous functions of degree d on \mathbb{C}^{n+1} . These thus restrict to homogeneous functions of degree d on every line γ_x , whatever $x \in \mathbb{C}P^n$ is. As a consequence, these complex homogeneous polynomials of degree d define global holomorphic sections of the bundle γ_d^* so that we get an injective morphism $\mathbb{C}_d^{\text{hom}}[X_0, \dots, X_n] \hookrightarrow H^0(\mathbb{C}P^n; \gamma_d^*)$. It is not that hard to prove that this injective morphism is also surjective, but requires though two theorems in complex analysis, namely Hartog's theorem and the decomposition of entire functions into power series, see [2].

Upshot: It is important here to understand that a homogeneous polynomial $Q \in \mathbb{C}_d^{\text{hom}}[X_0, \dots, X_n]$ does not define a holomorphic function $\mathbb{C}P^n \rightarrow \mathbb{C}$ (any such function would be constant due to maximum's principle). Its vanishing subset in $\mathbb{C}^{n+1} \setminus \{0\}$ is a cone, and thus defines on the quotient $\mathbb{C}P^n$ the hypersurface $V_Q^{\mathbb{C}} = \{x \in \mathbb{C}P^n \mid Q(x) = 0\}$, provided Q is of positive degree. But the other level sets of Q in $\mathbb{C}^{n+1} \setminus \{0\}$ are not left invariants under homotheties and thus do not pass to the quotient $\mathbb{C}P^n$.

What is true is that these polynomials $Q \in \mathbb{C}_d^{\text{hom}}[X_0, \dots, X_n]$ define global sections of γ_d^* , and $V_Q^{\mathbb{C}}$ coincides with the vanishing locus of these as sections of γ_d^* .

2.3 Fubini-Study metric

Let me now equip \mathbb{C}^{n+1} with its standard Hermitean product, defined for every $v = (v_0, \dots, v_n)$ and $w = (w_0, \dots, w_n)$ in \mathbb{C}^{n+1} by $h(v, w) = \sum_{i=0}^n v_i \bar{w}_i \in \mathbb{C}$.

It restricts on every line γ_x of \mathbb{C}^{n+1} to a Hermitean product h . This is called a Hermitean metric on the line bundle γ . It also induces then a Hermitean metric h_d on all the line bundles γ_d^* , $d > 0$. Indeed, if $x \in \mathbb{C}P^n$ and $s(x) \in \gamma_d^*|_x$, then $s(x) : \gamma_x \rightarrow \mathbb{C}$ is a

homogeneous form of degree d and we set

$$\|s(x)\| = \frac{|s(x)(v)|}{\|v\|^d},$$

where this definition does not depend on the choice of $v \in \gamma_x \setminus \{0\}$.

Fundamental example: Let us compute the pointwise Fubini-study norm of $Q = X_0^d \in \mathbb{R}_d^{\text{hom}}[X_0, \dots, X_n]$, viewed as a section of γ_d^* .

I restrict myself to $U_0 \cong \mathbb{C}^n$, since it vanishes outside of U_0 . Let $x = [1 : x_1 : \dots : x_n] \in U_0$. Then, $v = (1, x_1, \dots, x_n)$ generates γ_x and $\|v\|^2 = h(v, v) = 1 + \sum_{i=1}^n |x_i|^2$. Since $Q(v) = |Q(v)| = 1$, we get

$$h_d(Q, Q)|_x = \frac{1}{(1 + \|x\|^2)^d} = \exp(-d \log(1 + \|x\|^2)).$$

This means that the norm of Q at the origin $[1 : 0 : \dots : 0] \in U_0$ equals one, but at every other point it decays exponentially fastly to zero as the degree grows to $+\infty$. Since $\log h_d(Q, Q)|_x = -d\|x\|^2 + o(\|x\|^2)$ near $x = 0$, we deduce that the Fubini-Study norm of Q gets concentrated in a ball of radius $\frac{1}{\sqrt{d}}$ centered at the origin. Such a section defined by Q is called a peak section, and the scale $\frac{1}{\sqrt{d}}$ is a fundamental scale in Kähler geometry. Peak sections exist over any projective or Stein manifolds, following the theory of L. Hörmander, see [10], [14].

Finally, it is possible to define sections of γ_d^* which peak near any point $x \in \mathbb{C}P^n$. Indeed, the group $GL_{n+1}(\mathbb{C})$ acts by linear automorphisms of \mathbb{C}^{n+1} and the unitary group $U_{n+1}(\mathbb{C})$ even by isometries. These actions are transitive on lines of \mathbb{C}^{n+1} and thus they induce actions on $\mathbb{C}P^n$ which are transitive on points. Moreover, these actions lift to actions on γ and thus on any line bundle γ_d^* , $d > 0$.

For every $x \in \mathbb{C}P^n$, there exists $r \in U_{n+1}(\mathbb{C})$ such that $x = r([1 : 0 : \dots : 0])$. Then, $Q \circ r^{-1} \in \mathbb{C}_d^{\text{hom}}[X_0, \dots, X_n]$ defines a section of γ_d^* which peaks near x .

2.4 Implementation of affine hypersurfaces

Let $\Sigma \subset \mathbb{R}^n$ be a closed hypersurface, not necessarily connected. It is a theorem of H. Seifert, see [20] or also [18], that there exists a polynomial $P \in \mathbb{R}[X_1, \dots, X_n]$ of some degree k such that $V_P = P^{-1}(0)$ contains a union of connected components $\tilde{\Sigma}$ which is isotopic to Σ . This means that there exists a path $(\phi_t)_{t \in [0,1]}$ of diffeomorphisms of \mathbb{R}^n such that ϕ_0 is the identity and $\phi_1(\tilde{\Sigma}) = \Sigma$. Note that this theorem of Seifert is similar to Stone-Weierstrass theorem, except that one needs some approximation in C^1 -norm. Note also that I could have immediately taken any polynomial P for which zero is a regular value and then defined Σ to be any union of closed connected components of V_P . From now on, let me fix P and denote by Σ such a union of closed connected components of V_P .

There exists $R > 0$ such that Σ is included in the ball $B(0, R) \subset \mathbb{R}^n$ of radius R . Let me replace, for every $d > 0$, P by the polynomial $P_d = P(\sqrt{d}\cdot)$. It is still a polynomial of

degree k , whose coefficients are $O(\sqrt{d}^k)$. Indeed, if $P = \sum_{(\alpha_1, \dots, \alpha_n) \in \mathbb{N}^n} a_{\alpha_1, \dots, \alpha_n} X_1^{\alpha_1} \dots X_n^{\alpha_n}$, then $P_d = \sum_{(\alpha_1, \dots, \alpha_n) \in \mathbb{N}^n} a_{\alpha_1, \dots, \alpha_n} \sqrt{d}^{\alpha_1 + \dots + \alpha_n} X_1^{\alpha_1} \dots X_n^{\alpha_n}$.

Under the isomorphism $\mathbb{R}_k[X_1, \dots, X_n] \mapsto \mathbb{R}_k^{\text{hom}}[X_0, \dots, X_n]$, P_d is mapped to the polynomial $Q_d = \sum_{(\alpha_1, \dots, \alpha_n) \in \mathbb{N}^n} a_{\alpha_1, \dots, \alpha_n} \sqrt{d}^{\alpha_1 + \dots + \alpha_n} X_0^{k - \alpha_1 - \dots - \alpha_n} X_1^{\alpha_1} \dots X_n^{\alpha_n}$. After multiplication by X_0^{d-k} , it provides a section $\sigma_P = Q_d X_0^{d-k} \in \mathbb{R}_d^{\text{hom}}[X_0, \dots, X_n] = \mathbb{R}H^0(\mathbb{C}P^n; \gamma_d^*)$ which vanishes in the ball $B([1 : 0 : \dots : 0], R/\sqrt{d}) \subset \mathbb{R}U_0 \subset \mathbb{R}P^n$ centered at the origin $[1 : 0 : \dots : 0]$ and of radius R/\sqrt{d} . Moreover, $\sigma_P^{-1}(0) \cap B([1 : 0 : \dots : 0], R/\sqrt{d})$ contains a union of components $\tilde{\Sigma}$ such that the pair $(B([1 : 0 : \dots : 0], R/\sqrt{d}), \tilde{\Sigma})$ gets diffeomorphic to (\mathbb{R}^n, Σ) . In addition, the pointwise Fubini-Study norm $h_d(\sigma_P, \sigma_P) = \|Q_d\|^2 \|X_0^{d-k}\|^2$ decays exponentially outside the origin as d grows to $+\infty$. This is indeed the case for X_0^{d-k} as we saw in the previous paragraph, while Q_d has fixed degree and coefficients $O(\sqrt{d}^k)$. We deduce that the Fubini-Study norm of σ_P is concentrated in the ball $(B([1 : 0 : \dots : 0], R/\sqrt{d}))$.

Finally, after composition on the right by some suitable $r \in O_{n+1}(\mathbb{R}) \subset U_{n+1}(\mathbb{C})$, we get for every $x \in \mathbb{R}P^n$ a section $\sigma_P \in \mathbb{R}_d^{\text{hom}}[X_0, \dots, X_n]$ such that $\sigma_P^{-1}(0) \cap B(x, R/\sqrt{d})$ contains a union of components $\tilde{\Sigma}$ for which the pair $(B(x, R/\sqrt{d}), \tilde{\Sigma})$ gets diffeomorphic to (\mathbb{R}^n, Σ) and such that the Fubini-Study norm of σ_P exponentially decreases outside of this ball $B(x, R/\sqrt{d})$. Note that since the radius of this ball converges to zero, the Riemannian metric of $\mathbb{R}P^n$ for which we take the ball does not matter. We will however introduce the Fubini-Study metric of $\mathbb{C}P^n$ in the next paragraph.

2.5 The probability measure μ revisited

Recall that I did introduce the projective spaces in §2.1 and their tautological line bundles in §2.2. These are line subbundles of some trivial vector bundle. Let $x \in \mathbb{C}P^n$ and $\gamma_x \subset \mathbb{C}^{n+1}$ be the line it represents. Let $y \in \gamma_x \setminus \{0\}$ and $p : \mathbb{C}^{n+1} \setminus \{0\} \rightarrow \mathbb{C}P^n$ be the canonical projection. Then, the differential map $d_y p : T_y(\mathbb{C}^{n+1} \setminus \{0\}) = \mathbb{C}^{n+1} \rightarrow T_x \mathbb{C}P^n$ contains γ_x in its kernel and restricts to an isomorphism $\gamma_x^\perp \rightarrow T_x \mathbb{C}P^n$, where γ_x^\perp stands for the orthogonal of γ_x with respect to the standard Hermitian product of \mathbb{C}^{n+1} , see §2.3.

This hyperplane γ_x^\perp does not depend on the choice of $y \in \gamma_x \setminus \{0\}$, but the isomorphism $d_y p|_{\gamma_x^\perp}$ does. By the way, the quotient of the trivial vector bundle $\mathbb{C}P^n \times \mathbb{C}^{n+1}$ by the tautological bundle γ is not isomorphic to the tangent bundle $T\mathbb{C}P^n$.

Exercise: Prove that the latter tangent bundle $T\mathbb{C}P^n$ is rather isomorphic to the bundle of morphisms from γ to the former quotient bundle (while the quotient bundle is isomorphic to the space of morphisms from the trivial line bundle to itself).

Let us now choose y of norm one, so that it lies in the intersection of the unit sphere with γ_x . This intersection is a circle, the orbit of the action of the unitary group $U_1(\mathbb{C})$ by homothety. The circle fibration $S^{2n+1} \rightarrow \mathbb{C}P^n$ this action produces is called the Hopf fibration. Still, the isomorphism $d_y p|_{\gamma_x^\perp}$ depends on the choice of $y \in S^{2n+1} \cap \gamma_x$, but up to an isometry, so that if we push forward under $d_y p$ the Hermitian product of γ_x^\perp , induced by restriction of the ambient one of \mathbb{C}^{n+1} , we get a well defined Hermitian product on

$T_x\mathbb{C}P^n$, which does not depend on the choice of $y \in S^{2n+1} \cap \gamma_x$. The collection of all these Hermitian products on all tangent spaces of $\mathbb{C}P^n$ defines a Hermitian metric on $\mathbb{C}P^n$ called the Fubini-Study metric. The action of $U_{n+1}(\mathbb{C})$ on $\mathbb{C}P^n$ we already discussed provides isometries for this metric.

Remark: A projective line for this Fubini-Study metric has total area π (exercise). We actually rescale in [5], [6] this metric by a factor $\frac{1}{\sqrt{\pi}}$ to normalize this area to one. This is quite natural from another point of view, since this Hermitian Fubini-Study metric, which is actually a Kähler metric, also originates from the curvature form of the canonical connection associated to the Fubini-Study metric of γ introduced in §2.3. Since the cohomology class of this form is the first Chern class of the line bundle γ , the volume of a projective line gets one for this metric. This Fubini-Study metric restricts to a Riemannian metric on $\mathbb{R}P^n$ and the quantity $\text{Vol}_{\text{FS}}\mathbb{R}P^n$ in Theorem 1.6 is the total volume of $\mathbb{R}P^n$ for this Riemannian Fubini-Study metric.

Now, since the line bundles γ_d^* are equipped with Hermitian metrics and their base $\mathbb{C}P^n$ with some volume form dx , induced by the Fubini-Study metric, the spaces $H^0(\mathbb{C}P^n; \gamma_d^*) = \mathbb{C}_d^{\text{hom}}[X_0, \dots, X_n]$ of global holomorphic sections of these bundles inherit some L^2 -Hermitian products, namely

$$(Q_1, Q_2) \in H^0(\mathbb{C}P^n; \gamma_d^*)^2 \mapsto \int_{\mathbb{C}P^n} h_d(Q_1, Q_2) dx \in \mathbb{C}.$$

These L^2 -Hermitian products restrict on the spaces $\mathbb{R}H^0(\mathbb{C}P^n; \gamma_d^*) = \mathbb{R}_d^{\text{hom}}[X_0, \dots, X_n]$ of real holomorphic sections to the L^2 -scalar products

$$(Q_1, Q_2) \in \mathbb{R}H^0(\mathbb{C}P^n; \gamma_d^*)^2 \mapsto \int_{\mathbb{C}P^n} h_d(Q_1, Q_2) dx \in \mathbb{R}.$$

Finally, now that the space of real homogeneous polynomials $\mathbb{R}_d^{\text{hom}}[X_0, \dots, X_n]$ is again Euclidean, it inherits some Gaussian measure $d\mu(P) = \frac{1}{\sqrt{\pi}^{N_d}} \exp(-\|P\|^2) dP$, where N_d denotes the dimension of $\mathbb{R}_d^{\text{hom}}[X_0, \dots, X_n]$ and dP the Lebesgue measure associated to this L^2 -scalar product.

Exercise: The monomials $X_0^{\alpha_0} \dots X_n^{\alpha_n}$ are orthogonal to each other and in fact the probability measure μ is the one considered in Theorem 1.6, so that $\sqrt{\frac{(d+n)!}{n!\alpha_0! \dots \alpha_n!}} X_0^{\alpha_0} \dots X_n^{\alpha_n}$ is an orthonormal basis (provided the Fubini-Study metric on $\mathbb{C}P^n$ is normalized so that its total volume is one ; it is $\pi^n/n!$ for the metric just defined).

Example: What is the L^2 -norm of the section σ_P we did construct in the previous §2.4?

Recall that $\sigma_P = Q_d X_0^{d-k}$, so that

$$\begin{aligned}
\|\sigma_P\|^2 &= \int_{\mathbb{C}P^n} h_d(\sigma_P, \sigma_P) dx \\
&= \int_{U_0} h_d(\sigma_P, \sigma_P) dx \text{ since } dx(\mathbb{C}P^n \setminus U_0) = 0 \\
&= \int_{\mathbb{C}^n} \frac{|P(\sqrt{d}x)|^2}{(1 + \|x\|^2)^d} dx \\
&\sim_{d \rightarrow +\infty} \frac{1}{d^n} \int_{\mathbb{C}^n} |P(y)|^2 \exp(-\|y\|^2) dy
\end{aligned}$$

The last equivalence is obtained after the change of variable $y = \sqrt{d}x$ and $dy = dx|_{[1:0:\dots:0]}$ denotes the standard Lebesgue measure of \mathbb{C}^n .

Now P has been fixed once for all, so that $\int_{\mathbb{C}^n} |P(y)|^2 \exp(-\|y\|^2) dy$ is a constant. From now on I will normalize σ_P by setting

$$\sigma_P = \sqrt{d}^n \frac{Q_d X_0^{d-k}}{\sqrt{\int_{\mathbb{C}^n} |P(y)|^2 \exp(-\|y\|^2) dy}}. \quad (2)$$

This section has L^2 -norm one asymptotically, this L^2 -norm being still concentrated in a ball of radius R/\sqrt{d} , but near the origin $[1:0:\dots:0]$, its pointwise Fubini-Study norm is of the order \sqrt{d}^n .

Note that the same holds true for the section $\sqrt{\frac{(d+n)!}{n!d!}} X_0^d$ above, which corresponds to σ_P for $P = 1$ (modulo the normalization of the volume) and this sheds some light on the coefficients $\sqrt{\frac{(d+n)!}{n!\alpha_0! \dots \alpha_n!}}$ instead of $\sqrt{\binom{d}{\alpha_0 \dots \alpha_n}}$ in the orthonormal basis obtained in the above exercise and introduced before Theorem 1.6.

2.6 Probability of presence of Σ

Recall that I did fix a closed hypersurface $\Sigma \subset B(0, R) \subset \mathbb{R}^n$ which does not need to be connected. I then did construct, for every $x \in \mathbb{R}P^n$, a homogeneous polynomial $\tilde{\sigma}_P \in \mathbb{R}_d^{\text{hom}}[X_0, \dots, X_n]$ such that $\sigma_P^{-1}(0) \cap B(x, R/\sqrt{d})$ contains a union of components $\tilde{\Sigma}$ for which the pair $(B(x, R/\sqrt{d}), \tilde{\Sigma})$ is diffeomorphic to (\mathbb{R}^n, Σ) . I now claim much more.

Theorem 2.1 (joint with Damien Gayet, [6]) *There exist $\tilde{c}_\Sigma > 0$ such that for every $x \in \mathbb{R}P^n$,*

$$\liminf_{d \rightarrow +\infty} \mu \left\{ \sigma \in \mathbb{R}_d^{\text{hom}}[X_0, \dots, X_n] \left| \begin{array}{l} \sigma^{-1}(0) \cap B(x, R/\sqrt{d}) \supset \tilde{\Sigma} \\ (B(x, R/\sqrt{d}), \tilde{\Sigma}) \cong (\mathbb{R}^n, \Sigma) \end{array} \right. \right\} \geq \tilde{c}_\Sigma.$$

Hence, not only there exists a polynomial $\sigma_P \in \mathbb{R}_d^{\text{hom}}[X_0, \dots, X_n]$ with our desired properties, but moreover we had a positive probability to find such, probability uniformly

bounded from below by a positive constant.

Proof:

First step: Let me choose tubular neighborhoods K and U of Σ , K being compact, such that $\Sigma \subset K \subset U \subset B(0, R)$ and

1. $|P|_{U \setminus K} > \delta$, so that in particular P does not vanish in $U \setminus K$,
2. If $|P(y)| \leq \delta, y \in U$, then $|d_y P| > \epsilon$,

for some $\delta, \epsilon > 0$.

Now, let me denote by Σ_d, K_d, U_d the images of Σ, K and U under the homothety of rate $\frac{1}{\sqrt{d}}$, so that $\Sigma_d \subset K_d \subset U_d \subset B(0, R/\sqrt{d})$. I get likewise,

1. $|\sigma_P|_{U_d \setminus K_d} > \delta \sqrt{d}^n$ and
2. If $|\sigma_P(y)| \leq \delta \sqrt{d}^n, y \in U_d$, then $|d_y \sigma_P| > \epsilon \sqrt{d}^{n+1}$,

for some may be slightly different constants $\delta, \epsilon > 0$.

This first step is called quantitative transversality. I knew that 0 is a regular value of σ_P , but I am quantifying how much transversal to the zero section σ_P is. Such kind of quantitative transversality played a key role in the construction by S. K. Donaldson of symplectic divisors in any closed symplectic manifold, see [3].

Second step:

Proposition 2.2 (joint with Damien Gayet, [6]) *There exist $C_1, C_2 > 0$ such that*

$$\mathbb{E}\left(\sup_{B(0, R/\sqrt{d})} |\sigma|\right) \leq C_1 \sqrt{d}^n \text{ and } \mathbb{E}\left(\sup_{B(0, R/\sqrt{d})} |d\sigma|\right) \leq C_2 \sqrt{d}^{n+1}. \quad \square$$

Let me skip the proof of this proposition, but point out however that

$$\begin{aligned} \mathbb{E}(|\sigma([1 : 0 : \cdots : 0])|) &= \int_{\mathbb{R}_d^{\text{hom}}[X_0, \dots, X_n]} |\sigma([1 : 0 : \cdots : 0])| d\mu(\sigma) \\ &= \int_{\langle X_0^d \rangle} |\sigma([1 : 0 : \cdots : 0])| d\mu(\sigma) \\ &\sim_{d \rightarrow +\infty} \frac{\sqrt{d}^n}{\sqrt{n!}}, \end{aligned}$$

if the volume of $\mathbb{C}P^n$ has been normalized to one.

Third step: (from now on we follow an approach similar to the one used by Nazarov and Sodin in [19]). Recall the following.

Theorem 2.3 (Markov's inequality) Let (Ω, μ) be a probability space and $f : \Omega \rightarrow \mathbb{R}^+$ be a random variable. Let $e = \mathbb{E}(f) = \int_{\Omega} f d\mu$ be its expectation. Then, for every $C > 0$, $\mu\{\omega \in \Omega \mid f(\omega) \geq C\} \leq e/C$.

Proof:

$$\begin{aligned} e &= \int_{\Omega} f d\mu \geq \int_{\{\omega \in \Omega \mid f(\omega) \geq C\}} f d\mu \geq C \mu\{\omega \in \Omega \mid f(\omega) \geq C\} \\ &\implies \mu\{\omega \in \Omega \mid f(\omega) \geq C\} \leq e/C. \quad \square \end{aligned}$$

Application: Since $\mathbb{E}(\sup_{B(0, R/\sqrt{d})} |\sigma|) \leq C_1 \sqrt{d}^n$, from Markov's inequality we deduce that

$$\mu\{\sigma \in \mathbb{R}_d^{\text{hom}}[X_0, \dots, X_n] \mid \sup_{B(0, R/\sqrt{d})} |\sigma| \geq 4C_1 \sqrt{d}^n\} \leq \frac{1}{4}$$

and likewise

$$\mu\{\sigma \in \mathbb{R}_d^{\text{hom}}[X_0, \dots, X_n] \mid \sup_{B(0, R/\sqrt{d})} |d\sigma| \geq 4C_2 \sqrt{d}^{n+1}\} \leq \frac{1}{4},$$

so that

$$\mu \left\{ \sigma \in \mathbb{R}_d^{\text{hom}}[X_0, \dots, X_n] \mid \begin{array}{l} \sup_{B(0, R/\sqrt{d})} |\sigma| \leq 4C_1 \sqrt{d}^n \\ \sup_{B(0, R/\sqrt{d})} |d\sigma| \leq 4C_2 \sqrt{d}^{n+1} \end{array} \right\} \geq \frac{1}{2}.$$

Last step:

Recall that I have to find a subset $E \subset \mathbb{R}_d^{\text{hom}}[X_0, \dots, X_n]$ of measure uniformly bounded from below by some positive constant, such that any polynomial σ in E has the property that $\sigma^{-1}(0) \cap B(x, R/\sqrt{d}) \supset \tilde{\Sigma}$ and $(B(x, R/\sqrt{d}), \tilde{\Sigma}) \cong (\mathbb{R}^n, \Sigma)$. This subset is going to be the set

$$E_M = \left\{ a\sigma_P + \tau \mid \begin{array}{l} a \geq M \\ \tau \in \sigma_P^{\perp} \\ \sup_{B(0, R/\sqrt{d})} |\tau| \leq 4C_1 \sqrt{d}^n \\ \sup_{B(0, R/\sqrt{d})} |d\tau| \leq 4C_2 \sqrt{d}^{n+1} \end{array} \right\},$$

with $M = \sup\{\frac{4C_1}{\delta}, \frac{4C_2}{\epsilon}\}$. The measure of E_M satisfies

$$\begin{aligned} \mu(E_M) &= \left(\int_M^{+\infty} \exp(-t^2) \frac{dt}{\sqrt{\pi}} \right) \mu \left\{ \tau \in \sigma_P^{\perp} \mid \begin{array}{l} \sup_{B(0, R/\sqrt{d})} |\tau| \leq 4C_1 \sqrt{d}^n \\ \sup_{B(0, R/\sqrt{d})} |d\tau| \leq 4C_2 \sqrt{d}^{n+1} \end{array} \right\} \\ &\geq \frac{1}{2} \int_M^{+\infty} \exp(-t^2) \frac{dt}{\sqrt{\pi}} \\ &= \tilde{c}_{\Sigma} > 0. \end{aligned}$$

One checks indeed that

$$\mu \left\{ \tau \in \sigma_P^\perp \left| \begin{array}{l} \sup_{B(0,R/\sqrt{d})} |\tau| \leq 4C_1\sqrt{d}^n \\ \sup_{B(0,R/\sqrt{d})} |d\tau| \leq 4C_2\sqrt{d}^{n+1} \end{array} \right. \right\} \geq \frac{1}{2}$$

as

$$\mu \left\{ \sigma \in \mathbb{R}_d^{\text{hom}}[X_0, \dots, X_n] \left| \begin{array}{l} \sup_{B(0,R/\sqrt{d})} |\sigma| \leq 4C_1\sqrt{d}^n \\ \sup_{B(0,R/\sqrt{d})} |d\sigma| \leq 4C_2\sqrt{d}^{n+1} \end{array} \right. \right\} \geq \frac{1}{2}.$$

Now, let $\sigma \in E_M$, $\sigma = a\sigma_P + \tau$. For every $t \in [0, 1]$, set $\sigma_t = a\sigma_P + t\tau$, so that $\sigma_0 = \sigma_P$ and $\sigma_1 = \sigma$. Then, for every $t \in [0, 1]$, σ_t vanishes transversely in the open set U_d . Indeed, let $x \in U_d$ and $t \in [0, 1]$ such that $\sigma_t(x) = 0$. Then

$$\begin{aligned} |a\sigma_P(x)| &= |t\tau(x)| \leq 4C_1\sqrt{d}^n \\ \implies |\sigma_P(x)| &\leq \delta\sqrt{d}^n \\ \implies |d\sigma_P(x)| &> \epsilon\sqrt{d}^{n+1}, \end{aligned}$$

so that

$$\begin{aligned} |d\sigma_t| &= |ad\sigma_P + td\tau(x)| \\ &\geq |ad\sigma_P| - |d\tau(x)| \\ &> 4C_2\sqrt{d}^{n+1} - 4C_2\sqrt{d}^{n+1} \\ &> 0. \end{aligned}$$

We deduce that the smooth vanishing locus $\sigma_t^{-1}(0) \cap U_d = \Sigma_t$ remains trapped in the compact set K_d , thus cannot leave U_d and that it realizes an isotopy between $\Sigma = \sigma_P^{-1}(0) \cap U_d$ and $\sigma^{-1}(0) \cap U_d$. Hence the result. \square

2.7 Proof of the lower estimates

Let me address the following problem.

Question 3: Given a closed Riemannian manifold (M, g) and $\epsilon > 0$, how many disjoint balls of radius ϵ can be packed in M ?

In our case, (M, g) is going to be the real projective space $\mathbb{R}P^n$ equipped with its Fubini-Study metric and ϵ is going to be R/\sqrt{d} . Note that if $\mathbb{C}P^n$ is the quotient of the unit sphere S^{2n+1} under the action of the unit circle $U_1(\mathbb{C})$ by isometries, giving rise to the Hopf fibration, $\mathbb{R}P^n$ is just the quotient of the sphere S^{n+1} under the action of the group $\{\pm 1\}$ of unit elements of \mathbb{R} . This antipodal action is also isometric for the round metric of S^{n+1} and the Fubini-Study metric of $\mathbb{R}P^n$ is just the metric on the quotient $\mathbb{R}P^n = S^{n+1}/\{\pm 1\}$ induced by this round metric.

Proposition 2.4 *Let (M, g) be a closed Riemannian manifold of dimension n and $\epsilon > 0$. Let N_ϵ be the maximal number of disjoint balls of radius ϵ that can be packed in M . Then,*

$$\liminf_{\epsilon \rightarrow 0} (\epsilon^n N_\epsilon) \geq \frac{\text{Vol}_g(M)}{2^n \text{Vol}_{\text{eucl}}(B(0, 1))},$$

where $\text{Vol}_g(M)$ denotes the total Riemannian volume of M and $\text{Vol}_{\text{eucl}}(B(0, 1))$ the Euclidean volume of the unit ball in \mathbb{R}^n .

Note that it is of course not possible to fill more than the total volume of M by disjoint balls, so that

$$\limsup_{\epsilon \rightarrow 0} (\epsilon^n N_\epsilon) \leq \frac{\text{Vol}_g(M)}{\text{Vol}_{\text{eucl}}(B(0, 1))},$$

but from Proposition 2.4 we know that it is possible to fill a fraction of it. This packing problem is a classical one. For instance, in the case of the Euclidean space \mathbb{R}^n , the question may be, given a box, can we fill its whole volume with apples. Of course not and actually even if the radius of the apples was converging to zero. The question then becomes what is the best way to fill the box in order to loose the minimal amount of space, but we do not address this question. If instead of Euclidean balls, we just wanted to fill the manifolds with balls of a given volume, that is by disjoint images of embeddings of the Euclidean balls by diffeomorphisms which preserve the volume form, then it would be possible to fill the whole volume, say asymptotically due to Moser's trick. Finally, a famous theorem of M. Gromov establishes that it is not possible to fill the whole Fubini-Study volume of $\mathbb{C}P^2$ by packing two disjoint symplectic balls, see [8], [15], meaning two disjoint embeddings of some ball of \mathbb{C}^2 into $\mathbb{C}P^2$ which preserve the symplectic form.

Proof:

Let Λ_ϵ be a subset of points of M with the property that for every $x \neq y \in \Lambda_\epsilon$, $d(x, y) > 2\epsilon$ and that Λ_ϵ is maximal with respect to this property. Then, the balls centered at the points of Λ_ϵ and of radius ϵ are disjoint to each other, so that $\#\Lambda_\epsilon \leq N_\epsilon$. But the balls centered at the points of Λ_ϵ and of radius 2ϵ cover M since a point y in the complement of these balls in M could be added to Λ_ϵ to get a strictly larger set with our desired property, contradicting the maximality of Λ_ϵ . Thus

$$\text{Vol}_g(M) \leq \sum_{x \in \Lambda_\epsilon} \text{Vol}_g(B(x, 2\epsilon)) \sim_{\epsilon \rightarrow 0} \#\Lambda_\epsilon \epsilon^n 2^n \text{Vol}_{\text{eucl}}(B(0, 1)),$$

so that $\liminf_{\epsilon \rightarrow 0} (\epsilon^n \#\Lambda_\epsilon) \geq \frac{\text{Vol}_g(M)}{2^n \text{Vol}_{\text{eucl}}(B(0, 1))}$. \square

Let me now come back to the proof of the lower estimates in Theorem 1.6. Let $\epsilon = R/\sqrt{d}$ and Λ_ϵ be a subset of $(\mathbb{R}P^n, g_{\text{FS}})$ maximal with the property that for every $x \neq y \in \Lambda_\epsilon$, $d(x, y) > 2\epsilon$. For every closed connected hypersurface Σ of \mathbb{R}^n and $P \in \mathbb{R}_d^{\text{hom}}[X_0, \dots, X_n]$, let $N_\Sigma(V_P)$ be the number of connected components of $V_P = P^{-1}(0) \subset \mathbb{R}P^n$ which are diffeomorphic to Σ . For every $x \in \mathbb{R}P^n$, we set $N_{\Sigma, x}(V_P)$ to be one if $V_P \cap B(x, \epsilon) \supset \tilde{\Sigma}$

such that $(B(x, \epsilon), \tilde{\Sigma}) \cong (\mathbb{R}^n, \Sigma)$ and $N_{\Sigma, x}(V_P) = 0$ otherwise. We deduce in particular, $N_{\Sigma}(V_P) \geq \sum_{x \in \Lambda_{\epsilon}} N_{\Sigma, x}(V_P)$. Then,

$$\begin{aligned}
\mathbb{E}(b_i) &= \int_{\mathbb{R}_d^{\text{hom}}[X_0, \dots, X_n]} b_i(V_P) d\mu(P) \\
&\geq \int_{\mathbb{R}_d^{\text{hom}}[X_0, \dots, X_n]} \left(\sum_{[\Sigma] \in \mathcal{H}_n} b_i(\Sigma) N_{\Sigma}(V_P) \right) d\mu(P) \\
&\geq \sum_{[\Sigma] \in \mathcal{H}_n} b_i(\Sigma) \sum_{x \in \Lambda_{\epsilon}} \int_{\mathbb{R}_d^{\text{hom}}[X_0, \dots, X_n]} N_{\Sigma, x}(V_P) d\mu(P) \\
&\geq \sum_{[\Sigma] \in \mathcal{H}_n} b_i(\Sigma) \tilde{c}_{\Sigma} \# \Lambda_{\epsilon} \text{ from Theorem 2.1.}
\end{aligned}$$

Note that d being fixed, all the sums involved are finite. We then use Proposition 2.4 to deduce

$$\liminf_{\epsilon \rightarrow 0} \left(\frac{\# \Lambda_{\epsilon}}{\sqrt{d}^n} \right) \geq \frac{\text{Vol}_{\text{FS}}(\mathbb{R}P^n)}{2^n \text{Vol}_{\text{eucl}}(B(0, R))}.$$

We finally set

$$c_{\Sigma} = \frac{\tilde{c}_{\Sigma}}{2^n \text{Vol}_{\text{eucl}}(B(0, R))}$$

to get

$$\liminf_{d \rightarrow +\infty} \left(\frac{\mathbb{E}(b_i)}{\sqrt{d}^n \text{Vol}_{\text{FS}}(\mathbb{R}P^n)} \right) \geq \sum_{[\Sigma] \in \mathcal{H}_n} c_{\Sigma} b_i(\Sigma) = c_i^-. \quad \square$$

Note that we have actually proved that the expected number $\mathbb{E}(N_{\Sigma})$ of connected components of V_P diffeomorphic to Σ satisfies

$$\liminf_{d \rightarrow +\infty} \left(\frac{\mathbb{E}(N_{\Sigma})}{\sqrt{d}^n \text{Vol}_{\text{FS}}(\mathbb{R}P^n)} \right) \geq c_{\Sigma}.$$

We have even proved this lower estimate for a smaller quantity, the expected number of disjoint balls B of $\mathbb{R}P^n$ that contain a component $\tilde{\Sigma}$ of V_P for which the pair $(B, \tilde{\Sigma})$ gets diffeomorphic to (\mathbb{R}^n, Σ) .

References

- [1] G. Ben Arous and A. Guionnet. Large deviations for Wigner's law and Voiculescu's non-commutative entropy. *Probab. Theory Related Fields*, 108(4):517–542, 1997.
- [2] J.-P. Demailly. *Complex Analytic and Differential Geometry*. online book. <http://www-fourier.ujf-grenoble.fr/~demailly/manuscripts/agbook.pdf>.
- [3] S. K. Donaldson. Symplectic submanifolds and almost-complex geometry. *J. Differential Geom.*, 44(4):666–705, 1996.

- [4] A. Edelman and E. Kostlan. How many zeros of a random polynomial are real? *Bull. Amer. Math. Soc. (N.S.)*, 32(1):1–37, 1995.
- [5] D. Gayet and J.-Y. Welschinger. Betti numbers of random real hypersurfaces and determinants of random symmetric matrices. *To appear in J. Eur. Math. Soc., arXiv:1207.1579*, 2012.
- [6] D. Gayet and J.-Y. Welschinger. Lower estimates for the expected betti numbers of random real hypersurfaces. *J. London Math. Soc., published online, arXiv:1303.3035*, 2013.
- [7] P. Griffiths and J. Harris. *Principles of algebraic geometry*. Wiley Classics Library. John Wiley & Sons, Inc., New York, 1994. Reprint of the 1978 original.
- [8] M. Gromov. Pseudoholomorphic curves in symplectic manifolds. *Invent. Math.*, 82(2):307–347, 1985.
- [9] A. Harnack. Ueber die Vieltheiligkeit der ebenen algebraischen Curven. *Math. Ann.*, 10(2):189–198, 1876.
- [10] L. Hörmander. *An introduction to complex analysis in several variables*. D. Van Nostrand Co., Inc., Princeton, N.J.-Toronto, Ont.-London, 1966.
- [11] M. Kac. On the average number of real roots of a random algebraic equation. *Bull. Amer. Math. Soc.*, 49:314–320, 1943.
- [12] F. Klein. Ueber den Verlauf der Abel’schen Integrale bei den Curven vierten Grades. *Math. Ann.*, 10(3):365–397, 1876.
- [13] E. Kostlan. On the distribution of roots of random polynomials. In *From Topology to Computation: Proceedings of the Smalefest (Berkeley, CA, 1990)*, pages 419–431. Springer, New York, 1993.
- [14] X. Ma and G. Marinescu. *Holomorphic Morse inequalities and Bergman kernels*, volume 254 of *Progress in Mathematics*. Birkhäuser Verlag, Basel, 2007.
- [15] D. McDuff and L. Polterovich. Symplectic packings and algebraic geometry. *Invent. Math.*, 115(3):405–434, 1994. With an appendix by Yael Karshon.
- [16] M. L. Mehta. *Random matrices*, volume 142 of *Pure and Applied Mathematics (Amsterdam)*. Elsevier/Academic Press, Amsterdam, third edition, 2004.
- [17] J. Milnor. *Morse theory*. Based on lecture notes by M. Spivak and R. Wells. Annals of Mathematics Studies, No. 51. Princeton University Press, Princeton, N.J., 1963.
- [18] J. Nash. Real algebraic manifolds. *Ann. of Math. (2)*, 56:405–421, 1952.

- [19] F. Nazarov and M. Sodin. On the number of nodal domains of random spherical harmonics. *Amer. J. Math.*, 131(5):1337–1357, 2009.
- [20] H. Seifert. Algebraische Approximation von Mannigfaltigkeiten. *Math. Z.*, 41(1):1–17, 1936.
- [21] M. Shub and S. Smale. Complexity of Bezout’s theorem. II. Volumes and probabilities. In *Computational algebraic geometry (Nice, 1992)*, volume 109 of *Progr. Math.*, pages 267–285. Birkhäuser Boston, 1993.
- [22] R. Thom. Sur l’homologie des variétés algébriques réelles. In *Differential and Combinatorial Topology (A Symposium in Honor of Marston Morse)*, pages 255–265. Princeton Univ. Press, Princeton, N.J., 1965.

Université de Lyon
CNRS UMR 5208
Université Lyon 1
Institut Camille Jordan
43 blvd. du 11 novembre 1918
F-69622 Villeurbanne cedex
France
welschinger@math.univ-lyon1.fr