

BLOWUP FOR THE NONLINEAR HEAT EQUATION WITH SMALL INITIAL DATA IN SCALE-INVARIANT BESOV NORMS

LORENZO BRANDOLESE AND FERNANDO CORTEZ

ABSTRACT. We consider the Cauchy problem of the nonlinear heat equation $u_t - \Delta u = u^b$, $u(0, x) = u_0$, with $b \geq 2$ and $b \in \mathbb{N}$. We prove that initial data $u_0 \in \mathcal{S}(\mathbb{R}^n)$ (the Schwartz class) arbitrarily small in the scale invariant Besov-norm $\dot{B}_{n(b-1)b/2, q}^{-2/b}(\mathbb{R}^n)$, can produce solutions that blow up in finite time. The case $b = 3$ answers a question raised by Yves Meyer. Our result also proves that the smallness assumption put in an earlier work by C. Miao, B. Yuan and B. Zhang, for the global-in-time solvability, is essentially optimal.

1. INTRODUCTION

In this paper we study the Cauchy problem for the nonlinear heat equation

$$(1.1) \quad \begin{cases} \partial_t u = \Delta u + |u|^\alpha u, & x \in \mathbb{R}^n, \quad t \in [0, T] \\ u(0, x) = u_0(x), \end{cases}$$

where $\alpha > 0$, $0 < T \leq \infty$, and $u: \mathbb{R}^+ \times \mathbb{R}^n \rightarrow \mathbb{R}$ is a real function. This problem attracted a considerable interest and we refer to, *e.g.*, [1, 6–9, 12, 15, 16, 19, 25, 28] for a small sample of the huge existing literature.

Several well-posedness results are available for the Cauchy problem (1.1). For example, if $u_0 \in C_0(\mathbb{R}^n)$, then there is $T = T(u_0) > 0$ and a unique $u \in C([0, T], C_0(\mathbb{R}^n))$ which is a classical solution to (1.1) on $(0, T) \times \mathbb{R}^n$. For more singular data, say $u_0 \in L^p(\mathbb{R}^n)$, we know the following, see [4, 27, 28].

- When $p > \frac{n\alpha}{2}$ and $p \geq 1$, there exists a constant $T = T(u_0) > 0$ and a unique function $u(t) \in C([0, T], L^p(\mathbb{R}^n))$ that is a classical solution to (1.1) on $(0, T) \times \mathbb{R}^n$.
- When $p < \frac{n\alpha}{2}$, there is no general theory of existence. Besides, A. Haraux and F. Weissler [13] established the non-uniqueness, by showing that there is a positive solution in $C([0, T], L^p(\mathbb{R}^n)) \cap L_{loc}^\infty((0, T), L^\infty(\mathbb{R}^n))$, arising from zero initial data.
- When $p = \frac{n\alpha}{2}$, see Theorem 2.1 below.

We will be interested in the issues of the blowup in finite time *v.s.* the global existence of the solutions. The first works in this direction are due to H. Fujita. Fujita proved that for the positive solutions of (1.1), if the initial data u_0 is of class $C^2(\mathbb{R}^n)$ with derivatives up to the second order bounded on \mathbb{R}^n , then a necessary condition for u to be unique in

Date: February 13, 2019.

2010 Mathematics Subject Classification. 35K55 (primary), 30H25 (secondary).

Key words and phrases. Blowup, Nonlinear Heat Equation, Besov.

$C(\mathbb{R}^n \times [0, T])$ is that

$$\forall x \in \mathbb{R}^n, \quad |u_0(x)| \leq M e^{|x|^\beta},$$

for some constants $M > 0$ and $0 < \beta < 2$. See [10, 11].

About the problem of the existence of regular global solutions, there are two possible scenarios: if $n\alpha/2 < 1$, then no nontrivial positive solution of this problem can be global (a situation now referred as *Fujita's phenomenon*), while for $n\alpha/2 > 1$, there are global non-trivial positive solutions under small initial data assumptions. K. Hayakawa [14] and F. Weissler [27, 28] later proved that Fujita's phenomenon occurs in the case of the critical exponent $n\alpha/2 = 1$.

2. MOTIVATIONS AND OVERVIEW OF THE MAIN RESULT

To motivate our results, we introduce the concept of a scale-invariant space. For $\lambda > 0$, let us set

$$(2.1) \quad u_\lambda(t, x) = \lambda^{\frac{2}{\alpha}} u(\lambda^2 t, \lambda x) \quad \text{and} \quad u_{0,\lambda}(x) = \lambda^{\frac{2}{\alpha}} u_0(\lambda x).$$

For every solution $u(t, x)$ of (1.1), $u_\lambda(t, x)$ is also a solution of (1.1) with initial data $u_{0,\lambda}(x)$. In this setting, we say that a Banach space E is *scale-invariant*, if

$$(2.2) \quad \|u(t, \cdot)\|_E = \|u_\lambda(t, \cdot)\|_E.$$

Scale-invariant space are known to play an essential role in issues like well-posedness, global existence or blow-up of the solution.

The purpose of the present paper is to study the borderline cases of explosion and global existence for solutions of (1.1), in a scale-invariant Banach space. In the case of problem (1.1), the only $L^p(\mathbb{R}^n)$ -space invariant under the above scaling (2.1) is obtained for $p = n\alpha/2$. Notice that $p \geq 1$ if and only if α is larger or equal to the Fujita critical exponent. Therefore, we will be especially interested in solutions in $L^{n\alpha/2}(\mathbb{R}^n)$.

Our starting point is the following theorem, where we collect some of the results of Brezis, Cazenave and Weissler, in this scaling invariant setting.

Theorem 2.1 (See [27]. See also [4] for the uniqueness). *Let $u_0 \in L^{n\alpha/2}(\mathbb{R}^n)$, and assume that $n\alpha/2 > 1$. There exists a time $T = T(u_0) > 0$ and a function $u \in C([0, T], L^{n\alpha/2}(\mathbb{R}^n)) \cap L_{loc}^\infty((0, T], L^\infty(\mathbb{R}^n))$ such that u is a classical solution of (1.1) on $(0, T) \times \mathbb{R}^n$. Moreover,*

- (i) $\sup_{0 < t < T} t^{\sigma/2} \|u(\cdot, t)\|_p < +\infty$,
- (ii) $\lim_{t \rightarrow 0} t^{\sigma/2} \|u(\cdot, t)\|_p = 0$,

for any $\frac{n\alpha}{2} < p < \frac{n\alpha(\alpha+1)}{2}$ and $\sigma = \frac{2}{\alpha} - \frac{n}{p}$.

The uniqueness of classical solutions to (1.1) holds in the class $C([0, T], L^{n\alpha/2}(\mathbb{R}^n))$.

Moreover, there exists $\delta = \delta(\alpha, n)$ such that if $\|u_0\|_{n\alpha/2} < \delta$ then such solution is global, i.e., one can take T arbitrarily large.

The solution of Theorem 2.1 satisfies the integral equation

$$(2.3) \quad u(t) = e^{t\Delta} u_0(x) + \int_0^t e^{(t-s)\Delta} |u|^\alpha u(s) ds,$$

where $e^{t\Delta}f = G_t * f$ and

$$G_t(x) = (4\pi t)^{-n/2} e^{-|x|^2/(4t)}$$

is the standard Gaussian. The uniqueness of *weak solutions* to the integral equation (2.3) have been also addressed in [4]. The authors show that there is at most one *weak solution* to (3.1) in the class $C([0, T], L^{n\alpha/2}(\mathbb{R}^n)) \cap L_{loc}^\infty((0, T], L^\infty(\mathbb{R}^n))$, provided $\frac{n}{2}\alpha \geq \alpha + 1 > 1$. Under the additional restriction $\frac{n}{2}\alpha > \alpha + 1 > 1$ the uniqueness of weak solutions to (3.1) holds in the larger class $C([0, T], L^{n\alpha/2}(\mathbb{R}^n))$. We refer to E. Terraneo's paper [26] for further uniqueness/non uniqueness results of weak solutions.

The problem of obtaining *global-in-time* solutions by relaxing the stringent smallness assumption $\|u_0\|_{n\alpha/2} \ll 1$ was also addressed. New ideas in this direction were brought by M. Cannone and Y. Meyer's works on the Navier–Stokes equations [5, 18].

In the model case $\alpha = 2$ and $n = 3$, *i.e.* for the cubic heat equation in \mathbb{R}^3 ,

$$\partial_t u = \Delta u + u^3,$$

Y. Meyer observed in his lecture notes [18] that if $u_0 \in L^3(\mathbb{R}^3)$, with

$$\|u_0\|_{\dot{B}_{6,\infty}^{-1/2}} \ll 1,$$

(this condition is considerably weaker than requiring $\|u_0\|_3 \ll 1$) then the maximal time T^* of the solution is $T^* = +\infty$. In fact, the method described therein would go through provided $\|u_0\|_{\dot{B}_{p,\infty}^{-1+3/p}} \ll 1$ and $3 < p < 9$. In [18], he also raised the question whether or not, for $u_0 \in L^3(\mathbb{R}^3)$, the even weaker smallness condition

$$\|u_0\|_{\dot{B}_{\infty,\infty}^{-1}} \ll 1$$

would still imply $T^* = +\infty$. See next section for the definition of Besov spaces. Notice that these Besov spaces enjoy the same scaling invariance properties as $L^3(\mathbb{R}^3)$ and we have the continuous injections $L^3(\mathbb{R}^3) \subset \dot{B}_{p,\infty}^{-1+3/p}(\mathbb{R}^3) \subset \dot{B}_{\infty,\infty}^{-1}(\mathbb{R}^3)$ ($3 < p \leq +\infty$). Moreover, $\dot{B}_{\infty,\infty}^{-1}$ is known to be the largest function space invariant under translation and satisfying such scaling property. In this sense, a smallness condition on the $\dot{B}_{\infty,\infty}^{-1}$ -norm would be the least demanding restriction that one could put in a scale-invariant setting.

In the same spirit, but for the general case of problem (1.1), the best result for the global-in-time existence are due to Miao, Yuan, and Zhang [20]. They proved (among other things) that the solution of Theorem 2.1 is global, provided $u_0 \in L^{n\alpha/2}(\mathbb{R}^n)$, with $n\alpha/2 > 1$, under the smallness condition

$$\|u_0\|_{\dot{B}_{p,q}^{-2/\alpha+n/p}} \ll 1, \quad \text{for some } 1 < \frac{n\alpha}{2} < p < \frac{n\alpha(\alpha+1)}{2}, \quad 1 \leq q \leq \infty.$$

The restriction $\frac{n\alpha}{2} < p$, together with the condition $q \geq p$, ensure the embedding of $L^{n\alpha/2}(\mathbb{R}^n)$ into $\dot{B}_{p,q}^{-2/\alpha+n/p}(\mathbb{R}^n)$. On the other hand the authors of [20] left open the limit case $p = n\alpha(\alpha+1)/2$. In other words, they left open the question whether or not initial data $u_0 \in L^{n\alpha/2}(\mathbb{R}^n)$, small in the $\dot{B}_{n\alpha(\alpha+1)/2,q}^{-2/(\alpha+1)}$ -norm, give rise to global-in-time solutions.

Our main result below provides a negative answer to the above problem, thus settling the borderline problem of the global solvability of (2.4), at least in the case of integer nonlinearity exponents.

More specifically, for $b \in \mathbb{N}$, we consider the Cauchy problem for the non-linear heat equation

$$(2.4) \quad \begin{cases} \partial_t u = \Delta u + u^b \\ u(0, x) = u_0(x) \end{cases} \quad x \in \mathbb{R}^n, t \in (0, T).$$

where $0 < T \leq \infty$. The results recalled for the problem (1.1) —in particular Theorem 2.1— remain valid for (2.4), with $b = \alpha + 1$. These two Cauchy problems in fact agree for positive solutions, or for real solutions of any sign, when b is an odd integer.

Theorem 2.2. *For any $\delta > 0$ and $b < q \leq +\infty$, with $b \in \mathbb{N}$ and $n(b-1)/2 > 1$, there exists $u_0 \in \mathcal{S}(\mathbb{R}^n)$ (the Schwartz class) such that*

$$(2.5) \quad \|u_0\|_{\dot{B}_{nb(b-1)/2, q}^{-2/b}} \leq \delta,$$

and such that the maximal time T^* of the solution $u \in C([0, T^*), L^{n(b-1)/2}(\mathbb{R}^n))$ to (2.4) arising from u_0 is finite.

As mentioned in the introduction, for $n(b-1)/2 \leq 1$, because of Fujita's phenomenon, finite time blow up occurs for positive solutions, no matter which norms of the initial data are assumed to be small.

In the case $b = n = 3$, Theorem 2.2 negatively answers Y. Meyer's question [18, Conjecture 1].

There are several blowup results for (1.1) based on the maximum principle, energy functionals, concavity methods, or the spectral properties of the Laplacian, etc. See e.g. [2] for a review of these classical methods. But none of them seems to be effective to establish Theorem 2.2, as the smallness condition (2.5) represents a severe obstruction for their applicability.

The proof of Theorem 2.2 is constructive: suitable initial data are given by (4.5) below with $N = N(\delta)$ large enough. Our approach, inspired by Palais [22], rather uses the positivity of the Fourier transform inherited from its initial condition u_0 . Even though conceptually similar to [22], our paper is technically completely different (for example, we are able to remove the restriction $b \leq 1 + 2/n$ that appears therein). From the technical point of view, our paper is somehow closer to [17, 21], where the authors studied the blowup for different equations, namely diffusion problems with *nonlocal quadratic* nonlinearity. Our method bears also some relation with that of [23]. However, the blowup result in $\mathcal{F}(L^1)$ of [23] is not put in relation with the size of the data in scale-invariant norms. As such, our blowup result looks more precise, and its proof shorter.

Since the work of Cannone [5] we know that fast enough oscillations of the initial give rise to global-in-time smooth solutions for a large class of semilinear dissipative system, and that size conditions on Besov norms with negative regularity represent an effective way to measure such oscillations. The main interest of our result is to illustrate a limitation of this principle, by showing that there are scale invariant Besov norms that turn out to be too weak to be used for this purpose.

In our blowup result, the maximal time T^* can be taken arbitrarily small, as one easily checks applying Theorem 1 to rescaled data $u_{0,\lambda}$, that have the same Besov norm as in (2.5), and existence time $T_\lambda^* = \lambda^{-2}T^*$.

In the more difficult case of the Navier–Stokes equations, a similar problem was addressed by Bourgain and Pavlović [3] (see also [29]). These authors considered the Cauchy problem for Navier–Stokes with small data in $\dot{B}_{\infty,\infty}^{-1}$. While they left open the hard problem of the blowup, they succeeded in constructing a solution featuring a “norm-inflation” phenomenon in such Besov space, after an arbitrarily short time. But it was later realized by O. Sawada [24] that Bourgain’s–Pavlović solution, in fact, *does not blow up* in finite time.

3. PRELIMINARIES

Let us recall the definition of the Besov norms and the Littlewood-Paley decomposition: let $\psi \in \mathcal{S}(\mathbb{R}^n)$ such that $\text{supp } \widehat{\psi} \subset \{3/4 \leq |\xi| \leq 8/3\}$ and

$$1 = \sum_{j=-\infty}^{\infty} \widehat{\psi}_j(\xi) \quad (\xi \in \mathbb{R}^n, \xi \neq 0),$$

where $\psi_j(x) = 2^{nj}\psi(2^jx)$, $j \in \mathbb{Z}$. Here and throughout, \widehat{f} denotes the Fourier transform of f . The homogeneous Besov spaces $\dot{B}_q^{s,p}$ can be defined as follows, at least for $s < n/p$ and $1 \leq p, q \leq \infty$, which will be our case (in this paper we will only deal with the case $s < 0$):

$$\dot{B}_{p,q}^s = \{f \in \mathcal{S}'(\mathbb{R}^n) : f = \sum_{j \in \mathbb{Z}} (2^{js}\psi_j * f) \text{ in the } \mathcal{S}'(\mathbb{R}^n)\text{-sense, and } \|f\|_{\dot{B}_{p,q}^s} < \infty\},$$

where, for $1 \leq q < +\infty$,

$$\|f\|_{\dot{B}_{p,q}^s} = \left(\sum_{j=-\infty}^{\infty} \|2^{js}\psi_j * f\|_p^q \right)^{1/q}$$

and $\|f\|_{\dot{B}_{p,\infty}^s} = \sup_{q \in \mathbb{Z}} \|2^{js}\psi_j * f\|_p$.

As mentioned before, one can obtain in Theorem 2.1 the global existence of the solution, dropping the smallness assumption on $\|u_0\|_{n(b-1)/2}$, and putting instead a smallness assumption on the $\dot{B}_{p,q}^{-2/(b-1)+n/p}$ -norm of the data, which is weaker than the $L^{n(b-1)/2}$ -norm. Let us sketch a proof of this fact, following the arguments of [18, 20], putting in evidence the admissible range for p , which is $n(b-1)/2 < p < nb(b-1)/2$.

One rewrites Equation (2.4) in the equivalent Duhamel formulation

$$(3.1) \quad u(t, x) = e^{t\Delta}u_0(x) + \int_0^t e^{(t-\tau)\Delta}u^b(\tau, x) d\tau =: \Phi(u)(t, x).$$

If $u_0 \in L^{n(b-1)/2}(\mathbb{R}^n)$, then the solution $u \in C([0, T], L^{n(b-1)/2}(\mathbb{R}^n))$ of Theorem 2.1 (recall that $\alpha = b-1$) is obtained through the contraction mapping theorem, as the limit $u = \lim u_l$ of approximate solutions (where $u_1 = e^{t\Delta}u_0$ and $u_{l+1} = \Phi(u_l)$, for $l = 1, 2, \dots$)

in the X -norm, where

$$\begin{aligned} \|u\|_X &:= \sup_{0 < t < T} \|u(t)\|_{n(b-1)/2} + \sup_{0 < t < T} t^{1/(b-1)-n/(2p)} \|u(t)\|_p \\ &=: \|u\|_Y + \|u\|_Z. \end{aligned}$$

Indeed, first notice that $e^{t\Delta}u_0 \in X$ by standard heat kernel estimates. Next, the key estimates for the nonlinear term are the following:

$$\begin{aligned} (3.2) \quad \left\| \int_0^t e^{(t-s)\Delta} u^b(s) ds \right\|_p &\leq C \int_0^t (t-s)^{-\frac{n}{2}(\frac{b}{p}-\frac{1}{p})} \|u^b(s)\|_{p/b} ds \\ &\leq C \|u\|_Z^b \int_0^t (t-s)^{-\frac{n}{2}(\frac{b}{p}-\frac{1}{p})} s^{-b/(b-1)+nb/(2p)} ds \\ &\leq C \|u\|_Z^b t^{-1/(b-1)+n/(2p)}, \end{aligned}$$

and

$$\begin{aligned} (3.3) \quad \left\| \int_0^t e^{(t-s)\Delta} u^b(s) ds \right\|_{n(b-1)/2} &\leq C \int_0^t (t-s)^{-\frac{n}{2}(\frac{b}{p}-\frac{2}{n(b-1)})} \|u^b(s)\|_{p/b} ds \\ &\leq C \|u\|_Z^b. \end{aligned}$$

These estimates are valid when $1 < n(b-1)/2 < p < nb(b-1)/2$ (one also needs here $1 < b \leq p$, but the restriction $b \leq p$ can be dropped after the solution is constructed, by interpolation). These estimate ensure that

$$\|\Phi(u)\|_X \leq C \|u\|_Z^b.$$

The Lipschitz estimates

$$\|\Phi(u) - \Phi(v)\|_X \leq C(\|u\|_Z^{b-1} + \|v\|_Z^{b-1})\|u - v\|_Z,$$

is established in a similar way. But $\|u_0\|_{\dot{B}_{p,\infty}^{-2/(b-1)+n/p}} \simeq \|u_1\|_Z$ owing to the heat kernel characterization of Besov spaces (see [5]). Hence, starting with u_0 small enough in the $\dot{B}_{p,\infty}^{-2/(b-1)+n/p}$ -norm allow to construct a solution with maximal lifetime $T^* = +\infty$.

Without any smallness assumption, a well known variant [4, 27] of the above argument still allows to construct a solution $u = \lim u_l$ in the X -norm, at least when $T > 0$ is small enough. This relies on the observation that the approximate solutions (and hence the solution u itself) satisfy the additional condition $\lim_{t \rightarrow 0} t^{1/(b-1)-n/(2p)} \|u_l(t)\|_p = 0$, for all l .

4. PROOF OF MAIN THEOREM

We start with a simple general remark about the properties of solutions of Theorem 2.1, arising from initial data in the Schwartz class. In this case, or more in general when $u_0 \in L^1 \cap L^{n(b-1)/2}$, the corresponding solution obtained in Theorem 2.1 remains in $L^1(\mathbb{R}^n)$ during the whole lifetime of the solution, and $u \in C([0, T], L^1(\mathbb{R}^n))$. This could be seen applying Gronwall-type estimates, or otherwise with the following argument: our claim is

immediate if $b > n(b-1)/2$. Indeed, in this case we may take $p = b$ in Theorem 2.1, and we have $\sup_{0 < t < T} t^{1/(b-1)-n/(2b)} \|u(t)\|_b < \infty$. So,

$$\|u(t)\|_1 = \|\Phi(u)(t)\|_1 \leq \|u_0\|_1 + C(u_0) \int_0^t s^{-b/(b-1)+n/2} ds \leq C(u_0, T),$$

because our condition $n(b-1)/2 > 1$ ensures that the above integral is finite for all finite $T > 0$. Moreover, the continuity with respect to t is obvious. On the other hand, if $n(b-1)/2 \geq b$ then we first observe that $u^b \in C([0, T], L^{n(b-1)/(2b)})$, next that $e^{t\Delta} u_0 \in L^{n(b-1)/(2b)}$ by interpolation, and from the integral equation $u(t) = \Phi(u)(t)$ we deduce $u \in C([0, T], L^{n(b-1)/(2b)} \cap L^{n(b-1)/2})$. If $n(b-1)/(2b^2) \leq 1$, then using again the equation $u = \Phi(u)$ we get by interpolation that $u^b \in C([0, T], L^1(\mathbb{R}^n))$ and so $u \in C([0, T], L^1(\mathbb{R}^n))$. Otherwise we iterate this argument, until we find $m \in \mathbb{N}$ such that $n(b-1)/(2b^m) \leq 1$ and we conclude as before.

In the same way, going back to the sequence (u_l) of approximate solutions introduced in the previous section, one can prove that when $u_0 \in L^1 \cap L^{n(b-1)/2}$ not only the convergence $u_l \rightarrow u$ holds in the X -norm, but also $u_l \rightarrow u$ in the $C([0, T], L^1)$ -norm, as $l \rightarrow \infty$.

Later on we will choose a specific $u_0 \in \mathcal{S}(\mathbb{R}^n)$ such that $\widehat{u}_0 \geq 0$ and \widehat{u}_0 is even (in a such way that u is real-valued). All the approximate solutions u_l constructed from such datum u_0 satisfy $\widehat{u}_l(t, \cdot) \geq 0$. The convergence of (u_l) in the $C([0, T], L^1(\mathbb{R}^n))$ -norm implies that $\widehat{u}(t, \cdot) \geq 0$ during the whole lifetime of the solution.

We introduce the following notation: for $b \in \mathbb{N}$, and a non-negative measurable function f , we denote

$$(4.1) \quad f^{*b} = \underbrace{f * \dots * f}_{b \text{ times}}.$$

Let us now state a useful lemma.

Lemma 4.1. *Let $\delta > 0$, $b \in \mathbb{N}$ ($b \geq 2$) and $w \in \mathcal{S}(\mathbb{R}^n)$, such that $\widehat{w} \geq 0$. Let $c_\delta = 1 - e^{-\frac{\delta}{2}(b^2-1)}$. Also assume that the support of \widehat{w} is contained in the ball $B(0, 1)$. Let w_k , α_k and t_k be defined by the recursive relations ($k \geq 1$):*

$$\begin{cases} w_0 = w \\ w_k = w_{k-1}^b, \end{cases} \quad \begin{cases} \alpha_0 = 1 \\ \alpha_k = \alpha_{k-1}^b b^{-2k} c_\delta, \end{cases} \quad \begin{cases} t_0 = 0 \\ t_k = t_{k-1} + b^{-2k} \frac{\delta}{2} (b^2 - 1). \end{cases}$$

Then, if u is the solution of (3.1) with initial condition $u_0(x) \in L^{n(b-1)/2}(\mathbb{R}^n)$, and if $\widehat{u}_0(\xi) \geq A\widehat{w}(\xi)$ with $A > 0$, then, for any $k \in \mathbb{N}$,

$$(4.2) \quad \widehat{u}(t, \xi) \geq A^{b^k} \alpha_k e^{-b^k t} \mathbf{1}_{t \geq t_k} \widehat{w}_k(\xi),$$

where $\mathbf{1}_{t \geq t_k}$ is the indicator function of the interval $[t_k, +\infty)$.

Proof. Using Fourier transform, we have that (3.1) becomes

$$(4.3) \quad \widehat{u}(\xi, t) = e^{-t|\xi|^2} \widehat{u}_0(\xi) + \int_0^t e^{(s-t)|\xi|^2} [\widehat{u}(s, \xi)]^{*b} ds.$$

We start with the case $k = 0$. We have $\widehat{u}(t, \cdot) \geq 0$, because $\widehat{u}_0(\xi) \geq A\widehat{w}(\xi) \geq 0$, as observed at the beginning of this section. Then, using that $\text{supp } \widehat{w} \subset \{|\xi| \leq 1\}$, we get

$$(4.4) \quad \begin{aligned} \widehat{u}(\xi, t) &\geq e^{-t|\xi|^2} \widehat{u}_0(\xi) \geq e^{-t|\xi|^2} A\widehat{w}(\xi) \geq A e^{-t} \widehat{w}(\xi) \\ &= A \alpha_0 e^{-t} \widehat{w}_0(\xi), \quad \forall t \geq 0. \end{aligned}$$

This agrees with (4.2) for $k = 0$. Suppose now that inequality (4.2) holds for $k - 1$. Then we get, for all $t \geq t_k$:

$$\begin{aligned} \widehat{u}(\xi, t) &\geq \int_0^t e^{(s-t)|\xi|^2} [\widehat{u}(s, \xi)]^{*b} ds \\ &\geq \int_{t_{k-1}}^t e^{(s-t)|\xi|^2} (A^{b^{k-1}} \alpha_{k-1})^b e^{-b^k s} [\widehat{w}_{k-1}(\xi)]^{*b} ds. \\ &\geq A^{b^k} \alpha_{k-1}^b \widehat{w}_k(\xi) \int_{t_{k-1}}^t e^{(s-t)|\xi|^2} e^{-b^k s} ds \\ &\geq A^{b^k} \alpha_{k-1}^b \widehat{w}_k(\xi) e^{-b^k t} \int_{t_{k-1}}^t e^{(s-t)b^{2k}} ds, \end{aligned}$$

where in the last inequality we used that $\text{supp } w_k \subset \{|\xi| \leq b^k\}$.

But, for $t \geq t_k$, we have

$$\begin{aligned} \int_{t_{k-1}}^t e^{(s-t)b^{2k}} ds &= b^{-2k} (1 - e^{-b^{2k}(t-t_{k-1})}) \\ &\geq b^{-2k} c_\delta, \end{aligned}$$

because $t_k - t_{k-1} = b^{-2k} \frac{\delta}{2}(b^2 - 1)$, and so $1 - e^{-b^{2k}(t_k - t_{k-1})} = c_\delta$.

Hence we get,

$$\begin{aligned} \widehat{u}(\xi, t) &\geq A^{b^k} \alpha_{k-1}^b b^{-2k} c_\delta e^{-b^k t} \mathbf{1}_{t \geq t_k} \widehat{w}_k(\xi) \\ &\geq A^{b^k} \alpha_k e^{-b^k t} \mathbf{1}_{t \geq t_k} \widehat{w}_k(\xi), \end{aligned}$$

by the recursive relation defining α_k . Our claim now follows by induction. \square

For later use, let us observe that closed form for the sequences introduced in the previous lemma w_k , α_k and t_k , are

$$w_k = w^{b^k} \quad (k \geq 0),$$

next

$$\alpha_k = b^{-\frac{2b}{(b-1)^2} b^k + \frac{2}{b-1} k + \frac{2b}{(b-1)^2}} c_\delta^{\frac{b^k - 1}{b-1}} \quad (k \geq 0),$$

and

$$t_k = \frac{\delta}{2} (b^2 - 1) \sum_{j=1}^k b^{-2j} \quad (k \geq 1)$$

as it is easily checked.

Next lemma provides a first blowup result for equation (3.1).

Lemma 4.2. *Let $\delta > 0$ and $w \in \mathcal{S}(\mathbb{R}^n)$ ($w \neq 0$) be a Schwartz function such that $\widehat{w} \geq 0$. Also assume that the support of \widehat{w} is contained in the ball $B(0, 1)$. Let $u_0 \geq Aw$, with $A \geq b^{2b/(b-1)^2} c_\delta^{-1/(b-1)} e^{\delta/2} \|\widehat{w}\|_1^{-1}$. If u is the classical solution of (3.1) arising from u_0 and belonging to $C([0, T^*], L^{n(b-1)/2}(\mathbb{R}^n))$, then $T^* \leq \frac{\delta}{2}$.*

Proof. Assume, by contradiction, $T^* > \frac{\delta}{2}$. Applying Lemma 4.1, and using that $t_k \uparrow \frac{\delta}{2}$ as $k \rightarrow +\infty$, we get, for $t = \delta/2$ and all $k \in \mathbb{N}$,

$$\begin{aligned} \|\widehat{u}(\delta/2, \cdot)\|_1 &\geq A^{b^k} \alpha_k e^{-b^k \delta/2} \|\widehat{w}_k\|_1 \\ &= A^{b^k} \alpha_k e^{-b^k \delta/2} \|\widehat{w}\|_1^{b^k}, \end{aligned}$$

by Tonelli's theorem and the non-negativity of \widehat{w} . The size condition on A ensures that, taking $\sup_{k \in \mathbb{N}}$ in the right-hand side, one gets $\|\widehat{u}(\delta/2, \cdot)\|_1 = +\infty$. But by the positivity of $\widehat{u}(t, \cdot)$ and Fourier inversion formula,

$$\|u(\delta/2, \cdot)\|_{L^\infty} \geq (2\pi)^{-n} \|\widehat{u}(\delta/2, \cdot)\|_{L^1} = +\infty.$$

This contradicts the fact that the lifetime of Weissler solution satisfies $T^* > \delta/2$. \square

Proof of Theorem 2.2. Let $\delta > 0$ fixed and $w \in \mathcal{S}(\mathbb{R}^n)$ such that $\widehat{w} \neq 0$ and $\widehat{w} \geq 0$. We also assume that \widehat{w} is an even function and its support is contained in the ball $B(0, \frac{1}{2b})$. Let also $u_{0,N} \in \mathcal{S}(\mathbb{R}^n)$ be defined as

$$(4.5) \quad u_{0,N}(x) = \epsilon_N \sum_{k=0}^N 2^{2k/b} \eta_k \cos\left(\frac{3}{2} 2^k x_1\right) w(x) \quad (N \in \mathbb{N}),$$

where the sequences (η_k) and (ϵ_N) are chosen in the following way:

$$(4.6) \quad \eta_k = 1/(1+k)^{1/b}, \quad \epsilon_N = 1/\log(\log(3+N)).$$

In fact, the only thing that does matter in what follows are the following properties of (η_k) and (ϵ_N) : they must be nonnegative and such that $(\eta_k) \in \ell^q$ if and only if $q > b$, and $\epsilon_N \rightarrow 0$, with $\epsilon_N^b \sum_{k=0}^N \eta_k^b \rightarrow +\infty$ as $N \rightarrow +\infty$. In the case $b \geq 3$ is odd we will additionally need $\epsilon_N^b \sum_{k=0}^{N-1} \eta_k^{b-1} \eta_{k+1} \rightarrow +\infty$. The choice (4.6) does satisfy these requirements.

Observe that the Fourier transform of $\cos(\frac{3}{2} 2^k x_1) w(x)$ is $\frac{1}{2} [\widehat{w}(\xi + \frac{3}{2} 2^k e_1) + \widehat{w}(\xi - \frac{3}{2} 2^k e_1)]$, which is contained in the union of two balls centered at $\pm \frac{3}{2} 2^k e_1$ and radius $1/(2b)$ (and hence in a single dyadic annulus). Let us consider the homogeneous Littlewood–Paley decomposition, $\sum_{\xi \in \mathbb{Z}} \widehat{\psi}(2^{-j}\xi) = 1$, for $\xi \neq 0$, obtained using a radial function $\widehat{\psi} \in C_0^\infty(\mathbb{R}^n)$ which is supported in $\{\frac{3}{4} \leq |\xi| \leq \frac{8}{3}\}$ and constant equal to 1 in $\{\frac{5}{4} \leq |\xi| \leq \frac{7}{4}\}$. If we denote by $\Delta_j f = \psi_j * f$ the Littlewood–Paley dyadic blocks, then we see that

$$\Delta_j \cos\left(\frac{3}{2} 2^k x_1\right) w(x) = 0, \quad \text{if } k \neq j,$$

and

$$\Delta_j \cos\left(\frac{3}{2} 2^j x_1\right) w(x) = \cos\left(\frac{3}{2} 2^j x_1\right) w(x).$$

Thus, if $q \geq 1$ we get

$$\begin{aligned}
\|u_{0,N}\|_{\dot{B}_{n(b-1)b/2,q}^{-2/b}} &= \epsilon_N \left(\sum_{j \in \mathbb{Z}} 2^{-\frac{2}{b}qj} \|\Delta_j u_{0,N}\|_{n(b-1)b/2}^q \right)^{\frac{1}{q}} \\
&= \epsilon_N \left(\sum_{j=0}^N \eta_j^q \|w(x) \cos(\frac{3}{2}2^j x_1)\|_{n(b-1)b/2}^q \right)^{\frac{1}{q}} \\
&\leq \epsilon_N \left(\sum_{j=0}^N \eta_j^q \|w\|_{n(b-1)b/2}^q \right)^{\frac{1}{q}} \\
&= \epsilon_N \left(\sum_{j=0}^N \eta_j^q \right)^{1/q} \|w\|_{n(b-1)b/2}.
\end{aligned}$$

Thus, for any fixed $\delta > 0$ and $q > b$, we can find $N_0 \in \mathbb{N}$ such that

$$(4.7) \quad \|u_{0,N_0}\|_{\dot{B}_{n(b-1)b/2,q}^{-2/b}} < \delta.$$

Now, let T_N^* be the maximal time of the solution obtained in Theorem 2.1, arising from $u_{0,N}(x)$. We denote u_N this solution. We are going to prove that $T_N^* < +\infty$, and more precisely that $T_N^* < \delta$.

If $N \geq N_0$ and $T_N^* < \delta$ then there is nothing to prove. We thus pick $N \geq N_0$ and assume $T_N^* \geq \delta$. By the remark at the beginning of Section 4, we have $\widehat{u}_N(t, \xi) \geq 0$ for all $t \in [0, T_N^*]$. Thus, if $0 < t < T_N^*$, we get

$$\begin{aligned}
\widehat{u}_N(t, \xi) &= e^{-t|\xi|^2} \widehat{u}_{0,N}(\xi) + \int_0^t e^{-(t-s)|\xi|^2} [\widehat{u}_N(s, \xi)]^{*b} ds \\
&\geq e^{-t|\xi|^2} \widehat{u}_{0,N}(\xi)
\end{aligned}$$

We have

$$\widehat{u}_{0,N}(\xi) = \epsilon_N \sum_{k=0}^N 2^{2k/b} \eta_k \frac{1}{2} [\widehat{w}(\xi + \frac{3}{2}2^k e_1) + \widehat{w}(\xi - \frac{3}{2}2^k e_1)].$$

Hence, using that the support of \widehat{w} is contained in $\{|\xi| \leq \frac{1}{4}\}$,

$$\widehat{u}_N(t, \xi) \geq \epsilon_N \sum_{k=0}^N \left(\underbrace{e^{-t2^{2k+2}} 2^{2k/b} \eta_k \frac{1}{2} \widehat{w}(\xi + \frac{3}{2}2^k e_1)}_{=A_k(t,\xi)} + \underbrace{e^{-t2^{2k+2}} 2^{2k/b} \eta_k \frac{1}{2} \widehat{w}(\xi - \frac{3}{2}2^k e_1)}_{=B_k(t,\xi)} \right).$$

This implies that

$$(4.8) \quad (\widehat{u}_N)^{*b}(t, \xi) \geq \epsilon_N^b \sum_{k_1=0}^N \cdots \sum_{k_b=0}^N \left((A_{k_1} + B_{k_1}) * \cdots * (A_{k_b} + B_{k_b})(t, \xi) \right).$$

It is now convenient to distinguish two cases.

The case b even. We bound from below (4.8) retaining just a few terms of the above summation:

$$(\widehat{u}_N)^{*b}(t, \xi) \geq \epsilon_N^b \sum_{k=0}^N (A_k * B_k)^{*b/2}(t, \xi).$$

But,

$$\widehat{w}(\cdot + \frac{3}{2}2^k e_1) * \widehat{w}(\cdot - \frac{3}{2}2^k e_1) = \widehat{w} * \widehat{w},$$

hence,

$$(\widehat{u}_N)^{*b}(t, \xi) \geq \epsilon_N^b \sum_{k=0}^N e^{-bt2^{2k+2}} 2^{2k-b} \eta_k^b (\widehat{w})^{*b}(\xi).$$

Using that $\text{supp}(\widehat{w}^{*b}) \subset B(0, 1)$, we deduce that

$$\begin{aligned} \widehat{u}_N(t, \xi) &\geq \int_0^t e^{(s-t)|\xi|^2} \widehat{u}_N(s, \cdot)^{*b} ds \\ &\geq \epsilon_N^b \sum_{k=0}^N 2^{2k-b} \eta_k^b \left(\int_0^t e^{(s-t)-bs2^{2k+2}} ds \right) (\widehat{w})^{*b}(\xi) \\ &\geq \epsilon_N^b \sum_{k=0}^N \frac{2^{-b} \eta_k^b}{4b} \left(1 - e^{-t(b2^{2k+2}-1)} \right) e^{-t} (\widehat{w})^{*b}(\xi) \\ &\geq \epsilon_N^b \left(\sum_{k=0}^N \eta_k^b \right) \frac{2^{-b}}{4b} (1 - e^{-t(4b-1)}) e^{-t} (\widehat{w})^{*b}(\xi). \end{aligned}$$

Our choice of (η_k) and (ϵ_k) ensure that $\epsilon_N^b (\sum_{k=0}^N \eta_k^b) \rightarrow +\infty$ as $N \rightarrow +\infty$. Now let us take $t = \delta/2$ and $N \geq N_0$ large enough in a such way that

$$\epsilon_N^b \left(\sum_{k=0}^N \eta_k^b \right) \frac{2^{-b}}{4b} (1 - e^{-\delta(4b-1)/2}) e^{-\delta/2} \geq \frac{b^{2b/(b-1)^2} e^{\delta/2}}{c_\delta^{1/(b-1)} \|\widehat{w}\|_1^b}.$$

Hence Lemma 4.2 applies and implies that the lifetime of the solution of $u_t = \Delta u + u^b$ arising from the initial datum $u_N(\delta/2, \cdot)$ must blow up before the time $\delta/2$. By the uniqueness result of Theorem 2.1, this implies that $T_N^* < \delta$.

The case b odd. In this case we can write $b = 2m + 3$, with $m \in \mathbb{N}$. Going back to (4.8), we bound this expression from below in the following way:

$$(\widehat{u}_N)^{*b}(t, \xi) \geq \epsilon_N^b \sum_{k=0}^{N-1} \left((A_k * B_k)^{*m} * A_{k+1} * B_k * B_k \right)(t, \xi).$$

By the invariance of convolution products under translation, $\text{supp}(A_k * B_k)$ is contained in $\{|\xi| \leq 1/b\}$ and $\text{supp}(A_{k+1} * B_k * B_k)$ is contained in $\{|\xi| \leq 3/(2b)\}$. Hence,

$$\text{supp} \left((A_k * B_k)^{*m} * A_{k+1} * B_k * B_k \right)(t, \cdot) \subset \{|\xi| \leq 1/2\}.$$

Hence,

$$(\widehat{u}_N)^{*b}(t, \xi) \geq \epsilon_N^b \sum_{k=0}^{N-1} e^{-(m+3)t 2^{2k+3}} 2^{2k-b} \eta_k^{2m+2} \eta_{k+1} \widehat{w}^{*b}(\xi).$$

Arguing as before, we obtain,

$$\begin{aligned} \widehat{u}_N(t, \xi) &\geq \int_0^t e^{(s-t)|\xi|^2} (\widehat{u}_N)^{*b}(s, \cdot) ds \\ &\geq \epsilon_N^b \sum_{k=0}^{N-1} 2^{2k-b} \eta_k^{2m+2} \eta_{k+1} \left(\int_0^t e^{(s-t)-(m+3)s 2^{2k+3}} ds \right) \widehat{w}^{*b}(\xi) \\ &\geq \epsilon_N^b \left(\sum_{k=0}^{N-1} \eta_k^{b-1} \eta_{k+1} \right) \frac{2^{-b}}{8(m+3)} (1 - e^{-t(8(m+3)-1)}) e^{-t} \widehat{w}^{*b}(\xi). \end{aligned}$$

But $\epsilon_N^b \sum_{k=0}^{N-1} \eta_k^{b-1} \eta_{k+1} \rightarrow +\infty$ as $N \rightarrow +\infty$ and therefore we can conclude taking $t = \delta/2$ and applying Lemma 4.2, exactly as in the case b even. \square

5. CONCLUSIONS

The global-in-time solvability of the Cauchy problem for the nonlinear heat equation (2.4) in \mathbb{R}^n is usually obtained putting a smallness assumption on a suitable scale invariant norm of the initial data. However, in the present paper we proved that the scale-invariant norm of the Besov space $\dot{B}_{nb(b-1)/2, q}^{-2/b}$ is not suitable for this purpose: in fact, arbitrarily small initial data in this space (or in any larger scale invariant space) can give rise to solutions that blow up in finite time. While our method provides a few quantitative estimates on the solution, it gives little information on the nature of the blowup. The issue of the type of blowup has been thoroughly investigated, e.g., in [6, 16, 19].

Our result is sharp, in the sense that, for all $s > -2/b$, a smallness condition on the $\dot{B}_{p, q}^s$ -norm of u_0 (with $s - n/p = -2/b - 2/(b(b-1))$, to respect the scale invariance), which is slightly more stringent, does ensure that the solution is globally defined. On the other hand, the precise role of the third index q (that does not affect the scaling of the Besov norm) on this blowup issue is less clear: as the proof of Theorem 2.2 requires $q > b$, and breaks down when $q = b$, the following open problem naturally arises. (Here b does not need to be an integer): *Let $n(b-1)/2 > 1$ and u_0 a smooth and well decaying initial data as $|x| \rightarrow +\infty$; does the smallness assumption $\|u_0\|_{\dot{B}_{nb(b-1)/2, b}^{-2/b}} \ll 1$ imply that the solution of the Cauchy problem for $u_t = \Delta u + |u|^{b-1}u$ in \mathbb{R}^n is global in time ?*

ACKNOWLEDGEMENT

The authors are grateful to Paul Acevedo and to the Referee for their useful comments and suggestions.

Fernando Cortez was supported by Escuela Politecnica Nacional, Proyecto PII-DM-08-2016.

REFERENCES

- [1] J.M. Ball, *Remarks on blow-up and nonexistence theorems for nonlinear evolution equations.*, The Quarterly Journal of Mathematics **28** (1977), no. 4, 473–486.
- [2] C. Bandle and H. Brunner, *Blowup in diffusion equations: a survey*, J. Comput. Appl. Math. **97** (1998), no. 1-2, 3–22.
- [3] J. Bourgain and N. Pavlović, *Ill-posedness of the Navier–Stokes equations in a critical space in 3D*, Journal of Functional Analysis **255** (2008), no. 9, 2233–2247.
- [4] H. Brezis and T. Cazenave, *A nonlinear heat equation with singular initial data*, Journal D’Analyse Mathématique **68** (1996), no. 1, 277–304.
- [5] M. Cannone., *Harmonic analysis tools for solving the incompressible Navier-Stokes equations.*, Wavelets, paraproducts and Navier-Stokes With a preface by Yves Meyer., 1995.
- [6] C. Collot, F. Merle, and Pierre Raphaël, *Dynamics near the ground state for the energy critical nonlinear heat equation in large dimensions*, Comm. Math. Phys. **352** (2017), no. 1, 215–285.
- [7] M. A. Ebde and H. Zaag, *Construction and stability of a blow up solution for a nonlinear heat equation with a gradient term*, SeMA Journal **55** (2011), no. 1, 5–21.
- [8] A. Friedman, *Remarks on nonlinear parabolic equations.*, Proc. Sympos. Appl. Math **Vol. XVII** (1965), no. 17, 3–23.
- [9] ———, *Partial differential equations*, Robert E. Krieger Publishing Co **Vol. XVII** (1976), 3–23.
- [10] H. Fujita, *On the blowing up of solutions of the Cauchy problem for $u_t = \Delta u + u^{1+\alpha}$* , Journal of the Faculty of Science of the University of Tokyo **13** (1966), 109–124.
- [11] ———, *On some nonexistence and nonuniqueness theorems for nonlinear parabolic equations*, Proc. Symp. Pure Math. XVII Am. Math. Soc. **18** (1970), 105–113.
- [12] P. Groisman, J. Rossi, and H. Zaag, *On the dependence of the blow-up time with respect to the initial data in a semilinear parabolic problem* (2003).
- [13] A. Haraux and F. Weissler, *Non-uniqueness for a semilinear initial value problem*, Indiana University Mathematics Journal **31** (1982), no. 2, 167–189.
- [14] K. Hayakawa, *On nonexistence of global solutions of some semilinear parabolic differential equations*, Proceedings of the Japan Academy **49** (1973), no. 7, 503–505.
- [15] D. Henry, *Geometric theory of semilinear parabolic equations.* **840** (1985).
- [16] Miguel A. Herrero and Juan J. L. Velázquez, *Explosion de solutions d’équations paraboliques semilinéaires supercritiques*, C. R. Acad. Sci. Paris Sér. I Math. **319** (1994), no. 2, 141–145 (French, with English and French summaries).
- [17] P. G. Lemarié-Rieusset, *Sobolev multipliers, maximal functions and parabolic equations with a quadratic nonlinearity*, J. Funct. Anal. **274** (2018), no. 3, 659–694.
- [18] Y. Meyer, *Oscillating Patterns in Some Nonlinear Evolution Equations*, Mathematical Foundation of Turbulent Viscous Flows, 2006, pp. 101-187.
- [19] H. Matano and F. Merle, *Classification of type I and type II behaviors for a supercritical nonlinear heat equation*, J. Funct. Anal. **256** (2009), no. 4, 992–1064.
- [20] C. Miao, B. Yuan, and B. Zhang, *Strong solutions to the nonlinear heat equation in homogeneous Besov spaces.*, Nonlinear Analysis: Theory, Methods Applications **67** (2007), no. 5, 1329–1343.
- [21] S. Montgomery-Smith, *Finite time blow up for a Navier-Stokes like equation*, Proceedings of the American Mathematical Society **129** (2001), no. 10, 3025–3029.
- [22] B. Palais, *Blowup for nonlinear equations using a comparison principle in Fourier space*, Comm. Pure Appl. Math. **41** (1988), no. 2, 165–196.
- [23] S. Ru and J. Chen, *The blow-up solutions of the heat equations in $FL^1(\mathbb{R}^n)$* , J. Funct. Anal. **269** (2015), 1264–1288.
- [24] O. Sawada, *A remark on the Navier-Stokes flow with bounded initial data having a special structure*, Hokkaido Math. J. **43** (2014), 201–208.
- [25] R. Suzuki, *Critical Blow-Up for Quasilinear Parabolic Equations in Exterior Domains*, Tokyo Journal of Mathematics **12** (1996), no. 2, 397–409.

- [26] E. Terraneo, *Non-uniqueness for a critical non-linear heat equation*, Proceedings of the American Mathematical Society (2002).
- [27] F.B. Weissler, *Local existence and nonexistence for semilinear parabolic equations in L^p* , Indiana Univ. Math. J. **29** (1980), 79-102.
- [28] ———, *Existence and non-existence of global solutions for a semilinear heat equation*, Israel Journal of Mathematics **38** (1981), no. 1-2, 29–40.
- [29] T. Yoneda, *Ill-posedness of the 3D-Navier-Stokes equations in a generalized Besov space near.*, Journal of Functional Analysis **258** (2010), no. 10, 3376–3387.

L. BRANDOLESE: UNIVERSITÉ DE LYON, UNIVERSITÉ CLAUDE BERNARD LYON 1, INSTITUT CAMILLE JORDAN, UMR CNRS, 43 BD. DU 11 NOVEMBRE 1918, 69622 VILLEURBANNE CEDEX.

F. CORTEZ: ESCUELA POLITÉCNICA NACIONAL DEL ECUADOR, FACULTAD DE CIENCIAS, DEPARTAMENTO DE MATEMÁTICA, LADRÓN DE GUEVARA E11-253, QUITO-ECUADOR.