

LOCAL-IN-SPACE CRITERIA FOR BLOWUP IN SHALLOW WATER AND DISPERSIVE ROD EQUATIONS

LORENZO BRANDOLESE

ABSTRACT. We unify a few of the best known results on wave breaking for the Camassa–Holm equation (by R. Camassa, A. Constantin, J. Escher, L. Holm, J. Hyman and others) in a single theorem: a sufficient condition for the breakdown is that $u'_0 + |u_0|$ is strictly negative in at least one point $x_0 \in \mathbb{R}$. Such blowup criterion looks more natural than the previous ones, as the condition on the initial data is purely local in the space variable. Our method relies on the introduction of two families of Lyapunov functions. Contrary to McKean’s necessary and sufficient condition for blowup, our approach applies to other equations that are not integrable: we illustrate this fact by establishing new local-in-space blowup criteria for an equation modeling nonlinear dispersive waves in elastic rods.

1. INTRODUCTION

For many evolution equations, typical well-posedness results hold a priori only for short time intervals. A basic problem is to establish whether such intervals can be taken of arbitrary length. When it is not the case, one expects to find a maximal existence time $T^* < \infty$ and some spatial norms of the solution such that $\|u(t, \cdot)\|$ is finite for $t \in (0, T^*)$ and becomes unbounded as $t \uparrow T^*$. A blowup criterion is nothing but a condition on the initial data $u_0(x)$ implying the latter scenario. In this paper we will focus on blowup criteria of a very special nature: *local-in-space blowup criteria*. This notion requires the introduction of an equivalence relation in the class of initial data.

Let $x_0 \in \Omega$, where Ω is the spatial domain where the PDE is considered. We say that two initial data are “equivalent at x_0 ” if they are identical in a neighborhood of x_0 . A *local blowup criterion at x_0* is a condition on the equivalence class $[u_0]$, implying the finite time blowup of *any solution* arising from initial data equivalent to u_0 .

For evolution equations involving diffusive phenomena, blowup criteria are typically non local with respect to the space variable: indeed, it is usually possible to prevent the blowup of the solutions by perturbing the initial data only in some regions, leaving the data unchanged in the other regions.

For non-diffusive equations the situation is different. The simplest example is provided by the inviscid Burgers equation on the line, $u_t + uu_x = 0$. Applying the method of the characteristics shows that the condition for the formation of shock waves is that u_0 has a negative derivative in some point $x_0 \in \mathbb{R}$.

The goal of this paper is to establish the analogue of such elementary fact for more realistic models arising in one dimensional shallow water theory and for equations modeling

Key words and phrases. Camassa–Holm, rod equation, blowup, wave breaking, nonlinear waves, dispersive waves.

This research was partially supported by the ERC starting grant NuSiKiMo.

the propagation of dispersive waves in elastic rods. Such models are well-posed in Sobolev spaces $H^s(\mathbb{R})$, at least during a short time interval, for sufficiently large $s > 0$. Our blowup criterion will simply read

$$(1.1) \quad \inf_{\mathbb{R}} (u'_0 + \beta_\gamma |u_0|) < 0,$$

where β_γ is a constant depending on the physical parameter γ (related to the Finger deformation tensor) of the model. For the celebrated Camassa–Holm equation, such criterion boils down to

$$(1.2) \quad \inf_{\mathbb{R}} (u'_0 + |u_0|) < 0.$$

In the last 20 years, several hundreds of papers were devoted to study of the Camassa–Holm equation or its generalizations, and many of them addressed the issue of blowup (or wave breaking). At best of our knowledge, criteria (1.1)–(1.2) remained unnoticed, despite they were anticipated by many earlier results. In fact, we unify in this way some of the best known blowup criteria, including those established by R. Camassa, A. Constantin, J. Escher, L. Holm, J. Hyman, Y. Zhou, etc.). See Section 2.2 for a short survey of previous results and the relevant references.

A specific feature of the blowup criteria (1.1)–(1.2) is that they are local (in the sense of the previous definition) at the point x_0 where the negative infimum is achieved. This means that it is impossible to prevent the blowup without modifying u_0 *around the point* x_0 : perturbing u_0 only outside a neighborhood of x_0 may just help in delaying the formation of singularities.

A related physical interpretation is that fast oscillations will always lead to a blowup for such models (i.e., to the formation of a breaking wave or to the breaking of the rod in finite time), no matter how small is the region where the oscillations are present and how small are their amplitude.

The organization of the paper is straightforward. There is only one new theorem (Theorem 2.1). We state it and compare it with previously known results in Section 2. Section 3 is devoted to its proof and Section 4 to technical remarks and concluding observations.

2. THE MAIN RESULT

2.1. The compressible hyper-elastic rod equation. The propagation of nonlinear waves inside cylindrical hyper-elastic rods, assuming that the diameter is small when compared to the axial length scale, is described by the one dimensional equation

$$v_\tau + \sigma_1 v v_\xi + \sigma_2 v_{\xi\xi\tau} + \sigma_3 (2v_\xi v_{\xi\xi} + v v_{\xi\xi\xi}) = 0, \quad \xi \in \mathbb{R}, \tau > 0.$$

Such equation was derived by H.H. Dai [14]. Here $v(\tau, \xi)$ represents the radial stretch relative to a prestressed state, $\sigma_1 \neq 0$, $\sigma_2 < 0$ and $\sigma_3 \leq 0$ are physical constants depending by the material. The scaling transformations

$$\tau = \frac{3\sqrt{-\sigma_2}}{\sigma_1} t, \quad \xi = \sqrt{-\sigma_2} x,$$

with $\gamma = 3\sigma_3/(\sigma_1\sigma_2)$ and $u(t, x) = v(\tau, \xi)$, allow us to reduce the above equation to

$$(2.1) \quad u_t - u_{xxt} + 3uu_x = \gamma(2u_x u_{xx} + uu_{xxx}), \quad x \in \mathbb{R}, t > 0.$$

We complement this equation with vanishing boundary conditions at $\pm\infty$: such boundary conditions will be taken into account through an appropriate choice of the functional setting, guaranteeing the well-posedness of the Cauchy problem. The existence and the orbital stability of solitary waves for equation (2.1) is discussed e.g. in [13], [6], [15], [19]. In the case $\gamma = 1$, solitary waves are peaked solitons.

For $\gamma = 0$, the rod equation (2.1) reduces to the well-known BBM equation [1], modeling surface waves in a canal. In this case the solutions exist globally, meaning that the BBM equation can model permanent waves, but is unsuitable for describing breaking waves.

For $\gamma = 1$ (2.1) becomes the Camassa–Holm equation (CH), modeling long waves in shallow water. Such equation marked an important development in nonlinear dynamics and is of great current interest [4], [12], [16]. It admits strong solutions that exist globally and others that lead to a wave breaking in finite time. In this case equation (2.1) has a bi-hamiltonian structure and is integrable. The Camassa–Holm equation is thus much better understood than equation (2.1). Useful survey papers on such equation are [8] and [24].

We denote by

$$p(x) = \frac{1}{2}e^{-|x|}$$

the fundamental solution of the operator $1 - \partial_x^2$. Let $y = u - u_{xx}$ be the potential of u . We thus have $u = p * y$, and y satisfies

$$y_t + \gamma y_x u + 2\gamma y u_x + 3(1 - \gamma)u u_x = 0.$$

It is also convenient to rewrite the Cauchy problem associated with equation (2.1) in the following weak form:

$$(2.2) \quad \begin{cases} u_t + \gamma u u_x = -\partial_x p * \left(\frac{3-\gamma}{2} u^2 + \frac{\gamma}{2} u_x^2 \right), & t \in (0, T), x \in \mathbb{R}, \\ u(0, x) = u_0(x). \end{cases}$$

For any $\gamma \in \mathbb{R}$, the Cauchy problem for the rod equation is locally well-posed in H^s , when $s > 3/2$. More precisely, if $u_0 \in H^s(\mathbb{R})$, $s > 3/2$, then there exists a maximal time $0 < T^* \leq \infty$ and a unique solution $u \in C([0, T^*), H^s) \cap C^1([0, T^*), H^{s-1})$. Moreover, the solution u depends continuously on the initial data.

It is also known that u admits the invariants

$$E(u) = \int_{\mathbb{R}} (u^2 + u_x^2) dx$$

and

$$F(u) = \int_{\mathbb{R}} (u^3 + \gamma u u_x^2) dx.$$

In particular, the invariance of the Sobolev H^1 -norm of the solution implies that $u(x, t)$ remains uniformly bounded up to the time T^* . On the other hand, if $T^* < \infty$ then $\limsup_{t \rightarrow T} \|u(t)\|_{H^s} = \infty$ ($s > 3/2$) and the precise blowup scenario of strong solutions is the following:

$$(2.3) \quad T^* < \infty \iff \liminf_{t \rightarrow T^*} \left(\inf_{x \in \mathbb{R}} \gamma u_x(t, x) \right) = -\infty.$$

Applying the above result with $\gamma = 0$ one recovers the fact that solutions of the BBM do not blowup.

The above results are due to Constantin and Strauss [13]. They were previously established in the particular case $\gamma = 1$ corresponding to the Camassa–Holm equations (see e.g. [11]).

The main result of the present paper is the following blowup criterion for equation (2.2), in the case $1 \leq \gamma \leq 4$ (for $\gamma \notin [1, 4]$, see Section 4).

Theorem 2.1. *Let $1 \leq \gamma \leq 4$. There exists a constant $\beta_\gamma \geq 0$ (given by the explicit formula (3.5) below, see Figure 2.1) with the following property: Let T^* be the maximal time of the unique solution u of equation (2.2) in $C([0, T^*), H^3) \cap C^1([0, T^*), H^2)$ arising from $u_0 \in H^3(\mathbb{R})$. Assume that there exists $x_0 \in \mathbb{R}$ such that*

$$u'_0(x_0) < -\beta_\gamma |u_0(x_0)|,$$

then $T^* < \infty$.

The function $\gamma \mapsto \beta_\gamma$, defined for $\gamma \in [1, 4]$, is continuous, strictly decreasing on $[1, 3]$ and increasing on $[3, 4]$. Moreover, $\beta_1 = 1$, $\beta_3 = 0$ and $\beta_4 = \frac{1}{2}$.

Remark 2.2. It was observed in [13] that, for $\gamma = 3$, all non-zero solutions develop a singularity in finite time. This conclusion agrees with Theorem 2.1, since $\beta_3 = 0$ and any non-zero initial data $u_0 \in H^3$ must have a strictly negative derivative in some point.

Let us restate explicitly the result corresponding to the wave breaking for the Camassa–Holm equation ($\gamma = 1$):

Corollary 2.3. *Let T^* the maximal time of the unique solution $u \in C([0, T^*), H^3) \cap C^1([0, T^*), H^2)$ of the Camassa–Holm equation on \mathbb{R} ,*

$$u_t + uu_x = -\partial_x p * \left(u^2 + \frac{1}{2} u_x^2 \right),$$

starting from $u_0 \in H^3(\mathbb{R})$. If there exists $x_0 \in \mathbb{R}$ such that

$$u'_0(x_0) < -|u_0(x_0)|,$$

then $T^* < \infty$.

2.2. Earlier results on wave breaking for Camassa–Holm. In their pioneering paper [4], R. Camassa and L. Holm announced that if $u'_0(x_0)$ is negative with $|u'_0(x)|$ sufficiently large, then the solution of the Camassa–Holm equation must lead to a wave breaking. Several subsequent theorems confirmed their claim. The best known have been nicely surveyed by L. Molinet [24] and are the following. In the theorems below, the word “solution” must be understood as in Corollary 2.3.

Theorem 2.4 (Camassa, Holm, Hyman [5]). *The more precise formulation below is taken from [10]). If $u_0 \in H^3(\mathbb{R})$, u_0 is odd and $u'_0(0) < 0$, then the corresponding solution of the Camassa–Holm equation blows up in finite time.*




FIGURE 2.1. The plot of β_γ

The proof of the above theorem relies on the fact that the solution remains odd for all time, implying that $x \mapsto u_{xx}(t, x)$ is also odd and thus $u_{t,xx}(t, 0) = 0$ for all $t \in [0, T)$. One then deduces the differential inequality $\frac{d}{dt}[u_x(t, 0)] \leq -\frac{1}{2}u_x(t, 0)^2$, leading to the condition of blowup scenario.

Next theorem has been systematically adapted to the many different generalizations of the Camassa–Holm equation. It was anticipated by [4], [5], where one finds the idea of obtaining a differential inequality similar to the previous one for $m(t) = u_x(t, \bar{x}(t))$, where $\bar{x}(t) \in \mathbb{R}$ is a point where $u_x(\cdot, t)$ attains its minimum. The main technical difficulty consists in proving that the map $t \mapsto m(t)$ is a.e. differentiable. This subtle point was successfully addressed by A. Constantin and J. Escher. The resulting criterion reads as follows:

Theorem 2.5 (Constantin, Escher [11]). *Let $u_0 \in H^3(\mathbb{R})$ be such that, at some point $x_0 \in \mathbb{R}$, $u'(x_0) < -\|u_0\|_{H^1}/\sqrt{2}$. Then the solution of the Camassa–Holm equation blows up in finite time.*

A third interesting blow up criterion is provided by next theorem.

Theorem 2.6 (Constantin, [9]). *Let $u_0 \in H^3(\mathbb{R})$ be such that the associated potential $y_0 = u_0 - (u_0)_{xx}$ changes sign and satisfies, for some $x_0 \in \mathbb{R}$, $y(x) \geq 0$ for $x \leq x_0$ and $y_0(x) \leq 0$ for $x \geq x_0$. Then the solution of the Camassa–Holm equation blows up in finite time.*

Notice that these three criteria are *non-local* in the sense explained in the introduction. Moreover, despite a few common features, none of them implies the other two. On the other hand, Corollary 2.3 clearly unifies the three above results: The blowup criterion of Camassa, Holm and Hyman is considerably relaxed: for example, one can replace the antisymmetry condition on u_0 by the simpler requirement $u_0(0) = 0$. On the other hand,

it follows from Corollary 2.3 that solutions arising from small perturbations of a nonzero antisymmetric initial datum always blow up.

The criterion by Constantin and Escher follows immediately from the Sobolev embedding inequality

$$\|u\|_{L^\infty} \leq \|u_0\|_{H^1}/\sqrt{2}.$$

The blowup criterion of Constantin [9] is deduced from Corollary 2.3 applying the elementary identities (just apply integration by parts):

$$(2.4) \quad (u_0 - u'_0)(x) = e^{-x} \int_{-\infty}^x e^\xi y_0(\xi) \, d\xi,$$

$$(2.5) \quad (u_0 + u'_0)(x) = e^x \int_x^{+\infty} e^{-\xi} y_0(\xi) \, d\xi.$$

There is a fourth blowup criterion, due to Y. Zhou [25], that slightly extends Constantin's theorem. Zhou's theorem affirms that the solution of the Camassa–Holm equation blows up provided that there exists $x_1 \in \mathbb{R}$ such that

$$y(x_1) = 0, \quad \int_{-\infty}^{x_1} e^\xi y_0(\xi) \, d\xi > 0 \quad \text{and} \quad \int_{x_1}^{+\infty} e^{-\xi} y_0(\xi) \, d\xi < 0.$$

Using identities (2.4)-(2.5) shows that Corollary 2.1 encompasses also Zhou's criterion. In fact, the corollary essentially tells us that the above restriction $y(x_1) = 0$ can be dropped.

On the other hand, using some properties that are specific to the case $\gamma = 1$ (in particular the fact that the Camassa–Holm equation has a bi-hamiltonian structure and can be integrated), H. McKean succeeded in providing a necessary and sufficient condition for wave-breaking [23]. His theorem asserts that, if $u_0 \in H^3(\mathbb{R})$ and $y_0 \in L^1(\mathbb{R})$, then the solution of the Camassa–Holm equation will develop a singularity in finite time *if and only if* y_0 satisfies the following sign condition:

$$\exists x_1 < x_2 \quad \text{such that} \quad y_0(x_1) > 0 > y_0(x_2).$$

Notice that the condition $y_0 \in L^1(\mathbb{R})$ is not needed in Corollary 2.3. Even though in the case $\gamma = 1$ the result of Theorem 2.1 is not as precise as McKean's (because it provides a sufficient condition that is no longer necessary), its proof has the advantage of using in a less fundamental way the properties that are valid only for the Camassa–Holm equation. For this reason our theorem, unlike McKean's, remains valid also in the case $\gamma > 1$. Moreover, it seems hopeful that its proof can be adapted to a wide class of physically relevant equations arising in shallow water theory.

2.3. Main idea of the proof. As in [9], [21], the starting point is the analysis of the flow map $q(t, x)$, defined by

$$(2.6) \quad \begin{cases} q_t(t, x) = \gamma u(t, q(t, x)), & t \in (0, T), \, x \in \mathbb{R}, \\ q(0, x) = x. \end{cases}$$

One easily checks that $q \in C^1([0, T] \times \mathbb{R}, \mathbb{R})$ and $q_x(t, x) > 0$ for all $t \in (0, T)$ and $x \in \mathbb{R}$. In the case $\gamma = 1$, the potential $y = u - u_{xx}$ satisfies the fundamental identity

$$y(t, q(t, x))(q_x(t, x))^2 = y_0(x), \quad t \in [0, T], \quad x \in \mathbb{R} \quad (\gamma = 1),$$

implying that the zeros and the sign of y are invariant under the flow. This nice property of y is an essential feature of the Camassa–Holm. It implies a few remarkable algebraic identities (see [21]), and can be used not only to obtain wave breaking criteria, but also global existence results, [11], [9], [23].

For $\gamma \neq 1$ the zeros and the sign of y are no longer invariant of the flow. This explains why global existence results are much more difficult to establish in this case. Indeed, the above identity extends as follows (for $t \in [0, T)$ and $x \in \mathbb{R}$)

$$y(t, q(t, x))(q_x(t, x))^2 = y_0(x) + 3(\gamma - 1) \left(\int_0^t (uu_x)(s, q(s, x))q_x(x, s) \, d\xi \right) (q_x(t, x))^2.$$

Such formula is obtained from the previous one by applying the “variation of the constants”. It can also be checked directly by computing the total time derivative. But it turns out that this identity is not as useful as for $\gamma = 1$.

The essential idea of the proof of Theorem 2.1 is then to find two constants α and β such that, for all $x \in \mathbb{R}$, the functions

$$t \mapsto e^{\alpha q(t, x)}(\beta u - u_x)(t, q(t, x))$$

are monotone non-decreasing on $[0, T)$, and the functions

$$t \mapsto e^{-\alpha q(t, x)}(\beta u + u_x)(t, q(t, x))$$

are non-increasing. Such monotonicity properties will be convenient substitutes of McKean’s algebraic identities [21] established for the Camassa–Holm equation.

3. TWO FAMILIES OF LYAPUNOV FUNCTIONS

Proof of Theorem 2.1. As in [9], we shall look for a differential inequality for $\frac{d}{dt}u_x(t, q(t, x))$. Recalling that $\partial_x^2 p * f = p * f - f$, differentiating equation (2.2) with respect to the x variable yields

$$(3.1) \quad u_{tx} + \gamma uu_{xx} = \frac{3-\gamma}{2}u^2 - \frac{\gamma}{2}u_x^2 - p * \left(\frac{3-\gamma}{2}u^2 + \frac{\gamma}{2}u_x^2 \right).$$

For $0 \leq \gamma \leq 4$, let $\delta = \delta(\gamma)$ given by

$$(3.2) \quad \delta = \frac{\sqrt{\gamma}}{4}(\sqrt{12 - 3\gamma} - \sqrt{\gamma}).$$

Notice that $\delta \leq 0$ for $3 \leq \gamma \leq 4$. The above constant δ was characterized by Y. Zhou [26] as the best constant satisfying the inequality

$$p * \left(\frac{3-\gamma}{2}u^2 + \frac{\gamma}{2}u_x^2 \right) \geq \delta u^2.$$

But in fact we will need a more general estimate, established in the next lemma.

Lemma 3.1. *Let $0 \leq \gamma \leq 4$, $0 \leq \beta \leq 1$ and δ as in (3.2). Then*

$$(3.3) \quad (p \pm \beta \partial_x p) * \left(\frac{3-\gamma}{2}u^2 + \frac{\gamma}{2}u_x^2 \right) \geq \delta u^2.$$

Proof. We denote by $\mathbf{1}_{\mathbb{R}^+}$ and $\mathbf{1}_{\mathbb{R}^-}$ the characteristic functions of \mathbb{R}^+ and \mathbb{R}^- respectively. Let $a \in \mathbb{R}$:

$$\begin{aligned} (p\mathbf{1}_{\mathbb{R}^+}) * (a^2u^2 + u_x^2)(x) &= \frac{e^{-x}}{2} \int_{-\infty}^x e^{\xi} (a^2u^2 + u_x^2)(\xi) \, d\xi \\ &\geq ae^{-x} \int_{-\infty}^x e^{\xi} uu_x \, d\xi \\ &= \frac{au^2(x)}{2} - \frac{ae^{-x}}{2} \int_{-\infty}^x e^{\xi} u^2 \, d\xi \\ &= \frac{au^2(x)}{2} - a(p\mathbf{1}_{\mathbb{R}^+}) * (u^2)(x). \end{aligned}$$

This leads to

$$(p\mathbf{1}_{\mathbb{R}^+}) * ((a^2 + a)u^2 + u_x^2) \geq \frac{a}{2}u^2.$$

Choosing a to be the largest real root of the second order equation (a will be negative for $3 < \gamma \leq 4$)

$$a^2 + a = (3 - \gamma)/\gamma$$

we get $\gamma a = 2\delta$, hence

$$(p\mathbf{1}_{\mathbb{R}^+}) * \left(\frac{3-\gamma}{2}u^2 + \frac{\gamma}{2}u_x^2 \right) \geq \frac{\delta}{2}u^2.$$

The same computations also show that

$$(p\mathbf{1}_{\mathbb{R}^-}) * \left(\frac{3-\gamma}{2}u^2 + \frac{\gamma}{2}u_x^2 \right) \geq \frac{\delta}{2}u^2.$$

We have, both in the a.e. and the distributional sense,

$$\begin{aligned} p - \beta \partial_x p &= (1 - \beta)p\mathbf{1}_{\mathbb{R}^-} + (1 + \beta)p\mathbf{1}_{\mathbb{R}^+}, \\ p + \beta \partial_x p &= (1 + \beta)p\mathbf{1}_{\mathbb{R}^-} + (1 - \beta)p\mathbf{1}_{\mathbb{R}^+}. \end{aligned}$$

For $0 \leq \beta \leq 1$, taking the linear combination in the two last inequalities implies estimate (3.3). \square

The optimality of the constant δ in inequality (3.3) follows by the fact that for $0 < \gamma < 3$, the equality holds at the origin when $au = u_x$ in $(-\infty, 0)$ and $-au = u_x$ in $(0, \infty)$. Here a is the positive root of $a^2 + a = (3 - \gamma)/\gamma$. In other words, the choice $u(x) = e^{-a|x|}$ gives the equality in (3.3) for $x = 0$. When $3 < \gamma \leq 4$, then $a < 0$ and one chooses $u(x) = e^{a|x|}$.

Let $0 < T \leq T^*$. Recalling that $u \in C^1([0, T], H^2)$, we see that u and u_x are continuous on $[0, T) \times \mathbb{R}$ and $x \mapsto u(t, x)$ is Lipschitz, uniformly with respect to t in any compact time interval in $[0, T)$. Then the flow map $q(t, x)$ is well defined by (2.6) in the whole time interval $[0, T)$ and $q \in C^1([0, T) \times \mathbb{R}, \mathbb{R})$.

We have, for $0 < \gamma \leq 4$,

$$\begin{aligned} (3.4) \quad \frac{d}{dt} [u_x(t, q(t, x))] &= [u_{tx} + \gamma uu_{xx}](t, q(t, x)) \\ &= \frac{3-\gamma}{2}u^2 - \frac{\gamma}{2}u_x^2 - p * \left(\frac{3-\gamma}{2}u^2 + \frac{\gamma}{2}u_x^2 \right) \\ &\leq \left(\left(\frac{3-\gamma}{2} - \delta \right) u^2 - \frac{\gamma}{2}u_x^2 \right)(t, q(t, x)) \\ &= \frac{\gamma}{2}(\beta_\gamma^2 u^2 - u_x^2)(t, q(t, x)), \end{aligned}$$

where

$$\beta_\gamma^2 = \frac{3-\gamma}{\gamma} - \frac{2\delta}{\gamma}.$$

According to (3.2), then we set

$$(3.5) \quad \beta_\gamma = \left(-\frac{1}{2} + \frac{3}{\gamma} - \frac{\sqrt{12-3\gamma}}{2\sqrt{\gamma}} \right)^{1/2}.$$

This expression shows that $\gamma \mapsto \beta_\gamma$ is continuous, decreasing on $(0, 3]$, increasing for $\gamma \in [3, 4]$. Moreover, $\beta_1 = 1$ and $\beta_3 = 0$ and $\beta_4 = \frac{1}{2}$.

Next step is to find a good factorization of $(\beta_\gamma^2 u^2 - u_x^2)(t, q(t, x))$ ensuring some monotonicity properties for each factor. The obvious factorization $\beta_\gamma^2 u^2 - u_x^2 = (\beta_\gamma u - u_x)(\beta_\gamma u + u_x)$ is not the most interesting one, as it will be revealed by the next proposition.

This leads us to study the functions of the form:

$$A(t, x) = e^{\alpha q(t, x)}(\beta u - u_x)(t, q(t, x)),$$

and

$$B(t, x) = e^{-\alpha q(t, x)}(\beta u + u_x)(t, q(t, x)).$$

where $0 < \gamma \leq 4$, and α and β are real constants to be chosen later.

Computing the derivatives with respect to t using the definition of the flow map (2.6) gives, for $\alpha \in \mathbb{R}$ and $0 \leq \beta \leq 1$,

$$\begin{aligned} A_t(t, x) &= e^{\alpha q(t, x)} \left[\alpha \beta \gamma u^2 - \alpha \gamma u u_x + \beta (u_t + \gamma u u_x) - (u_{tx} + \gamma u u_{xx}) \right] \\ &= e^{\alpha q(t, x)} \left[\left(\alpha \beta \gamma - \frac{3-\gamma}{2} \right) u^2 + \frac{\gamma}{2} u_x^2 - \alpha \gamma u u_x + (p - \beta \partial_x p) * \left(\frac{3-\gamma}{2} u^2 + \frac{\gamma}{2} u_x^2 \right) \right] \\ &\geq e^{\alpha q(t, x)} \left[\left(\alpha \beta \gamma - \frac{3-\gamma}{2} + \delta \right) u^2 - \alpha \gamma u u_x + \frac{\gamma}{2} u_x^2 \right]. \end{aligned}$$

Here we used in the second equality equations (2.2) and (3.1), next the inequality provided by Lemma 3.1.

The condition on the discriminant

$$\Delta = \alpha^2 \gamma^2 - 2\gamma \left(\alpha \beta \gamma - \frac{3-\gamma}{2} + \delta \right) \leq 0$$

guarantees that the quadratic form inside the brackets is nonnegative. Minimizing Δ with respect to α gives $\alpha = \beta$. Choosing $\alpha = \beta$, the condition $\Delta \leq 0$ boils down to

$$\beta^2 \geq \frac{3-\gamma}{\gamma} - \frac{2\delta}{\gamma} = \beta_\gamma^2,$$

where β_γ is given by (3.5).

In particular, choosing $\alpha = \beta = \beta_\gamma$ (this is indeed possible when $1 \leq \gamma \leq 4$ but not when $0 < \gamma < 1$, as in the former case $0 \leq \beta \leq 1$ and Lemma 3.1 can be applied) we get

$$A_t(t, x) \geq 0, \quad \text{for all } t \in [0, T), x \in \mathbb{R}.$$

Similarly, for $\alpha \in \mathbb{R}$ and $0 \leq \beta \leq 1$,

$$\begin{aligned} B_t(t, x) &= -e^{-\alpha q(t, x)} \left[\left(\alpha \beta \gamma - \frac{3-\gamma}{2} \right) u^2 + \frac{\gamma}{2} u_x^2 + \alpha \gamma u u_x + (p + \beta \partial_x p) * \left(\frac{3-\gamma}{2} u^2 + \frac{\gamma}{2} u_x^2 \right) \right] \\ &\leq -e^{-\alpha q(t, x)} \left[\left(\alpha \beta \gamma - \frac{3-\gamma}{2} + \delta \right) u^2 + \alpha \gamma u u_x + \frac{\gamma}{2} u_x^2 \right]. \end{aligned}$$

The condition guaranteeing that the quadratic form inside the brackets is non-negative is identical to the previous one.

Summarizing, we established the following fundamental proposition:

Proposition 3.2. *Let $1 \leq \gamma \leq 4$, u as in Theorem 2.1 and β_γ as in (3.5). Set*

$$\begin{aligned} A(t, x) &= e^{\beta_\gamma q(t, x)} (\beta_\gamma u - u_x)(t, q(t, x)), \\ B(t, x) &= e^{-\beta_\gamma q(t, x)} (\beta_\gamma u + u_x)(t, q(t, x)). \end{aligned}$$

Then, for all $x \in \mathbb{R}$, the function $t \mapsto A(t, x)$ is monotonically increasing and $t \mapsto B(t, x)$ is decreasing.

It is then convenient to factorize

$$(\beta_\gamma^2 u^2 - u_x^2)(t, q(t, x)) = A(t, x)B(t, x).$$

This being achieved, the proof now can proceed like Constantin's [9]: from inequality (3.4) we get

$$(3.6) \quad \begin{aligned} \frac{d}{dt} [u_x(t, q(t, x))] &= \frac{\gamma}{2} (\beta_\gamma^2 u - u_x^2)(t, q(t, x)) \\ &= \frac{\gamma}{2} (AB)(t, x). \end{aligned}$$

Now let x_0 be such that $u'(x_0) < -\beta_\gamma |u(x_0)|$. We denote $g(t) = u_x(t, q(t, x_0))$, $A(t) = A(t, x_0)$ and $B(t) = B(t, x_0)$. Proposition 3.2 yields, for all $t \in [0, T)$,

$$A(t) \geq A(0) > 0 \quad \text{and} \quad B(t) \leq B(0) < 0$$

Thus, $AB(t) \leq AB(0) < 0$. Then we get, for all $t \in [0, T)$,

$$g'(t) \leq \frac{\gamma}{2} AB(t) \leq \frac{\gamma}{2} AB(0) < 0.$$

Assume, by contradiction, $T = \infty$. Let $\alpha_0 = \frac{\gamma}{2} (u'(0)^2 - \beta_\gamma^2 u_0^2)(x_0)$. Then $g(t) \leq g(0) - \alpha_0 t$. We can choose t_0 such that $g(0) - \alpha_0 t_0 \leq 0$ and $(g(0) - \alpha_0 t_0)^2 \geq \|u_0\|_{H^1}^2$. For $t \geq t_0$, using that $g(t) \leq g(t_0)$, and the conservation of the energy $E(u) = \|u(t)\|_{H^1}^2 = \|u_0\|_{H^1}^2$, we get

$$\begin{aligned} g'(t) &\leq \frac{\gamma}{2} A(t)B(t) = \frac{\gamma}{2} (\beta_\gamma^2 u^2 - u_x^2)(t, q(t, x_0)) \\ &\leq \frac{\gamma}{2} \left(\frac{\beta_\gamma^2}{2} \|u_0\|_{H^1}^2 - g(t)^2 \right) \\ &\leq -\frac{\gamma}{4} g(t)^2. \end{aligned}$$

This differential inequality implies that $u_x(t, q(t, x_0))$ must blow up in finite time. The condition of the blowup scenario (2.3) is fulfilled. \square

4. THE CASE $\gamma \notin [1, 4]$ AND EARLIER BLOW CRITERIA FOR THE ROD EQUATION

Experimental studies on a few hyper-elastic materials revealed that the parameter γ can range from -29.4770 to 3.4174, see [15]. The interval $[1, 4] \ni \gamma$ where Theorem 2.1 can be applied thus overlap, but not completely cover, the physically interesting cases.

Adapting to the elastic rod equation the proof of Theorem 2.5, Y. Zhou [26] established that for $0 < \gamma < 3$ a blowup occurs provided $\exists x_0 \in \mathbb{R}$ such that

$$(4.1) \quad |u'_0(x_0)| < -\frac{\beta_\gamma}{\sqrt{2}} \|u_0\|_{H^1}.$$

Theorem 2.1 thus improves Zhou's criterion in the range $1 \leq \gamma < 3$ and extends it to $3 \leq \gamma \leq 4$. On the other hand, Theorem 2.1 cannot be applied for $0 < \gamma < 1$. For reader's convenience, let us give a direct and simple proof of criterion (4.1), valid for $0 < \gamma \leq 4$. We only need to recall inequality (3.4)

$$\frac{d}{dt}[u_x(t, q(t, x))] \leq \frac{\gamma}{2}(\beta_\gamma^2 u^2 - u_x^2)(t, q(t, x)),$$

and the usual L^∞ -estimate

$$\|u\|_\infty \leq \frac{1}{\sqrt{2}}\|u\|_{H^1} = \frac{1}{\sqrt{2}}\|u_0\|_{H^1}.$$

Letting $g(t) = u_x(t, q(t, x_0))$, then we obtain the differential inequality

$$g'(t) \leq \frac{\gamma}{2}\left(\frac{\beta_\gamma^2}{2}\|u_0\|_{H^1}^2 - g(t)^2\right)$$

with the initial condition $g(0) = u'_0(x_0)$. Condition (4.1) ensures $g'(0) < 0$ and the blowup follows for $0 < \gamma \leq 4$ integrating this Riccati-type differential inequality .

Remark 4.1. The reason for excluding $\gamma < 1$ in Theorem 2.1 is that one would have $\beta_\gamma > 1$ (see Figure 2.1). In this case, it is not difficult to see that the result of Proposition 3.2 (implying that the condition $u'_0(x_0) < -\beta_\gamma|u_0(x_0)|$ is conserved by the flow) is no longer valid. The main reason is the lack of the inequalities

$$p \pm \beta \partial_x p \geq 0$$

when $\beta > 1$.

According to [13] and [15], equation (2.2) admits smooth and orbitally stable solitary waves solutions of the form $\phi_c(x - ct)$ for $\gamma < 1$ and orbitally stable peaked solitons $ce^{-|x-ct|}$ for $\gamma = 1$. As observed in [13], if $u(t, x) = \phi_c(x - ct)$ is a solution vanishing at infinity, then the profile ϕ_c must be a translate of $c\phi$, i.e.,

$$\phi_c(x) = c\phi(x - a), \quad x \in \mathbb{R},$$

where ϕ solves the differential equation

$$(\phi')^2 = \gamma\phi(\phi')^2 - \phi^3 + \phi^2.$$

Moreover, ϕ is positive, even and monotonically decreasing from its peak. In addition, it decays exponentially as $|x| \rightarrow \infty$, and the same for its derivatives. For $\gamma < 1$, the initial datum $u_0(x) = c\phi(x)$ gives rise to a global smooth solution $u_c = c\phi(x - ct)$. It follows from the last equation, the parity and the decay properties of ϕ , that we can find a large enough x_0 such that $u'(x_0, t) \simeq -|u(x_0, t)|$. This means that if one believes in the validity of a local-in-space blowup criterion of the form $u'_0(x_0) < -\alpha_\gamma|u_0(x_0)|$ also for $\gamma < 1$, then α_γ must satisfy $\alpha_\gamma \geq 1$.

The restriction $\gamma > 4$ might be purely technical as well. Of course, the expression of the coefficient β_γ computed in (3.5) in this case does not make sense, but one might conjecture that the statement of Theorem 2.1 holds with another coefficient α_γ .

Let us stress the fact that for $\gamma < 0$ or $\gamma > 4$ nonlocal blowup criteria are still available: for example, it was observed in [13], [26] that a blowup occurs when, for some x_0 ,

$$u'_0(x_0) > \frac{\sqrt{(\gamma-3)/\gamma}}{\sqrt{2}}\|u_0\|_{H^1}, \quad \text{if } \gamma < 0,$$

or

$$u'_0(x_0) < -\frac{\sqrt{(\gamma-3)/\gamma}}{\sqrt{2}} \|u_0\|_{H^1}, \quad \text{if } \gamma > 3.$$

This readily follows from equation (3.1). However, for $3 < \gamma \leq 4$, criterion (4.1) is slightly better, as $\beta_\gamma \leq \sqrt{(\gamma-3)/\gamma}$.

In this paper we did not discuss the continuation of the solution after the formation of a singularity. In general, there are several way to continue solutions beyond the wave breaking and it is possible to obtain weak solutions that conserve or that dissipate the energy $E(u)$. This issue is extensively studied e.g. in [2], [7], [18].

REFERENCES

- [1] T. B. Benjamin, J. L. Bona, and J. J. Mahony, *Model equations for long waves in nonlinear dispersive systems*, Philos. Trans. Roy. Soc. London Ser. A **272** (1972), no. 1220, 47–78.
- [2] Alberto Bressan and Adrian Constantin, *Global conservative solutions of the Camassa-Holm equation*, Arch. Ration. Mech. Anal. **183** (2007), no. 2, 215–239.
- [3] Lorenzo Brandolese, *Breakdown for the Camassa-Holm equation using decay criteria and persistence in weighted spaces*, Int. Math. Res. Not. **rnr218** (2011), 21 pages, DOI doi:10.1093/imrn/rnr218.
- [4] R. Camassa and L. Holm, *An integrable shallow-water equation with peaked solitons*, Phys. Rev. Letters **71** (1993), 1661–1664.
- [5] R. Camassa, L. Holm, and James M. Hyman, *A new integrable shallow-water equation*, Adv. Appl. Mech. **31** (1994), 1–33.
- [6] Robin Ming Chen, Yue Liu, and Zhijun Qiao, *Stability of solitary waves and global existence of a generalized two-component Camassa-Holm system*, Comm. Partial Differential Equations **36** (2011), no. 12, 2162–2188.
- [7] G. M. Coclite, H. Holden, and K. H. Karlsen, *Global weak solutions to a generalized hyperelastic-rod wave equation*, SIAM J. Math. Anal. **37** (2005), no. 4, 1044–1069 (electronic).
- [8] Adrian Constantin, *Geometrical methods in hydrodynamics*, Journées équations aux dérivées partielles (2001), Art. 2, 14p.
- [9] ———, *Existence of permanent and breaking waves for a shallow water equation: a geometric approach*, Ann. Inst. Fourier (Grenoble) **50** (2000), no. 2, 321–362.
- [10] Adrian Constantin and Joachim Escher, *Global existence and blow-up for a shallow water equation*, Ann. Scuola Norm. Pisa **26** (1998), no. 2, 303–328.
- [11] ———, *Wave breaking for nonlinear nonlocal shallow water equations*, Acta Math. **181** (1998), no. 2, 229–243.
- [12] Adrian Constantin and David Lannes, *The hydrodynamical relevance of the Camassa-Holm and Degasperis-Procesi equations*, Arch. Ration. Mech. Anal. **192** (2009), no. 1, 165–186.
- [13] Adrian Constantin and Walter A. Strauss, *Stability of a class of solitary waves in compressible elastic rods*, Phys. Lett. A **270** (2000), no. 3-4, 140–148.
- [14] H.-H. Dai, *Model equations for nonlinear dispersive waves in a compressible Mooney-Rivlin rod*, Acta Mech. **127** (1998), no. 1-4, 193–207.
- [15] H.-H. Dai and Y. Huo, *Solitary shock waves and other travelling waves in a general compressible hyperelastic rod*, R. Soc. Lond. Proc. Ser. A. Math. Phys. Eng. Sci. **456** (2000), 331–363.
- [16] Olivier Glass and Frank Sueur, *Smoothness of the flow map for low-regularity solutions of the Camassa-Holm equations*, Discrete and Continuous Dynamical System - A (to appear).
- [17] A. Alexandrou Himonas and Gerard Misiołek, *The Cauchy problem for an integrable shallow-water equation*, Differential Integral Equations **14** (2001), no. 7, 821–831.
- [18] Helge Holden and Xavier Raynaud, *Global conservative solutions of the generalized hyperelastic-rod wave equation*, J. Differential Equations **233** (2007), no. 2, 448–484.
- [19] Jonatan Lenells, *Traveling waves in compressible elastic rods*, Discrete Contin. Dyn. Syst. Ser. B **6** (2006), no. 1, 151–167 (electronic).

- [20] Yue Liu, *Global existence and blow-up solutions for a nonlinear shallow water equation*, Math. Ann. **335** (2006), no. 3, 717–735.
- [21] H.P. McKean, *Breakdown of a shallow water equation*, Asian J. Math. **2** (1998), no. 4, 867–874.
- [22] ———, *Correction to “Breakdown of a shallow water equation”*, Asian J. Math. **3** (1999).
- [23] ———, *Breakdown of the Camassa-Holm equation*, Comm. Pure Appl. Math. **57** (2004), no. 3, 416–418.
- [24] Luc Molinet, *On well-posedness results for Camassa-Holm equation on the line: a survey*, J. Nonlinear Math. Phys. **11** (2004), no. 4, 521–533.
- [25] Yong Zhou, *Wave breaking for a shallow water equation*, Nonlinear Anal. **57** (2004), no. 1, 137–152.
- [26] ———, *Local well-posedness and blow-up criteria of solutions for a rod equation*, Math. Nachr. **278** (2005), no. 14, 1726–1739.

L. BRANDOLESE: UNIVERSITÉ DE LYON ; UNIVERSITÉ LYON 1 ; CNRS UMR 5208 INSTITUT CAMILLE JORDAN, 43 BD. DU 11 NOVEMBRE, VILLEURBANNE CEDEX F-69622, FRANCE.

E-mail address: brandolese@math.univ-lyon1.fr

URL: <http://math.univ-lyon1.fr/~brandolese>