

Errata of the book  
*Analysis and Geometry of Markov Diffusion Operators*  
Springer, Grundlehren der mathematischen Wissenschaften  
Vol. 348 (2013)

Dominique Bakry, Ivan Gentil and Michel Ledoux

January 13, 2023

- Page 30, line 8: instead of  $\int (LP_t)^2 d\mu$  read  $\int (LP_t f)^2 d\mu$ , thanks to Kevin Tanguy.

- Page 35, line 10: instead of

$$\sum_{y \in E} \mu(x) L(x, y) = 0$$

read

$$\sum_{x \in E} \mu(x) L(x, y) = 0,$$

thanks to Michał Strzelecki.

- Page 36, line 4: instead of  $\hat{L}$  read  $K$ , thanks again to Michał Strzelecki.
- Page 39, line -1: instead of  $F(x, T)$  read  $F(T, x)$ , thanks to Michał Strzelecki.
- Page 42, line -4: the matrix  $\mathbf{g}$  is supposed to be also definite-positive, thanks to Michał Strzelecki.
- Page 49, line -13: instead of  $g = \psi'(f)$  read  $g = \psi(f)$ , thanks to Michał Strzelecki.
- Page 79, line -5: instead of

$$\sum_{i,j=1}^n (\partial_{ij} f)^2 \geq \frac{1}{n} \left( \sum_{i=1}^n \partial_i^2 f \right)^2$$

read

$$\sum_{i,j=1}^n (\partial_{ij}^2 f)^2 \geq \frac{1}{n} \left( \sum_{i=1}^n \partial_{ii}^2 f \right)^2,$$

thanks to Michał Strzelecki.

- Page 90, line 11: instead of

$$k_3(t, d) = \frac{1}{4\pi t} \frac{d}{\sinh(d)} \exp\left(-t - \frac{d^2}{4t}\right)$$

read (add a ( ))

$$k_3(t, d) = \frac{1}{(4\pi t)^{3/2}} \frac{d}{\sinh(d)} \exp\left(-t - \frac{d^2}{4t}\right)$$

- Page 129, line -1: instead of

$$\Gamma(f)(x) = \lim_{k \rightarrow \infty} \left( \frac{1}{2t_k} P_{t_k}(f^2)(x) - P_{t_k}(f)(x)^2 \right)$$

read

$$\Gamma(f)(x) = \lim_{k \rightarrow \infty} \frac{1}{2t_k} \left( P_{t_k}(f^2)(x) - P_{t_k}(f)^2(x) \right)$$

thanks to Michał Strzelecki.

- Page 152, line -7: Item (iii) has to be understood as follows, for any functions  $f_1, \dots, f_k \in \mathcal{A}$  and  $\Psi : \mathbb{R}^k \rightarrow \mathbb{R}$  a smooth function ( $\mathcal{C}^\infty$ ), then  $\Psi(f_1, \dots, f_k) \in \mathcal{A}$ .
- Page 156, line 17: instead of  $f - g$  read  $f + g$  (three times), thanks to Michał Strzelecki.
- Page 158, line 13: instead of

$$H(f)(g, h) = \frac{1}{2} \left[ \Gamma(g, \Gamma(f, h)) + \Gamma(h, \Gamma(f, g)) - \Gamma(f, \Gamma(g, h)) \right].$$

read (add a ( ))

$$H(f)(g, h) = \frac{1}{2} \left[ \Gamma(g, \Gamma(f, h)) + \Gamma(h, \Gamma(f, g)) - \Gamma(f, \Gamma(g, h)) \right].$$

- Page 170, line -6: remove  $L^*(f)$  at the beginning of the formula.
- Page 200, line -8 to the end of the page. Replace the paragraph by the following:

The second set (ii) of inequalities, without any boundary condition, appears as a consequence of (iii) by symmetrization and periodization (for  $f : [0, 1] \rightarrow \mathbb{R}$  arbitrary, define  $g : [-1, +1] \rightarrow \mathbb{R}$  by  $g(x) = f(x)$  for  $x \in [0, +1]$ ,  $g(x) = f(-x)$  for  $x \in [-1, 0]$ , and apply (iii) to  $g$  on the interval  $[-1, +1]$  after re-scaling).

Finally (i) is a consequence of (iii) by anti-symmetrization and periodization. For  $f : [0, 1] \rightarrow \mathbb{R}$  such that  $f(0) = f(1) = 0$ , define  $g : [-1, +1] \rightarrow \mathbb{R}$  by  $g(x) = f(x)$  for  $x \in [0, +1]$ ,  $g(x) = -f(-x)$  for  $x \in [-1, 0]$ . Then

$$\int_{[0,1]} f^2 dx = \int_{[-1,1]} g^2 \frac{dx}{2} - \left( \int_{[-1,1]} g \frac{dx}{2} \right)^2 \leq \frac{1}{\pi^2} \int_{[-1,1]} g'^2 \frac{dx}{2} = \frac{1}{\pi^2} \int_{[0,1]} f'^2 dx,$$

where (iii) has been applied to the probability measure  $1_{[-1,1]} \frac{dx}{2}$  with the optimal constant  $1/\pi^2$ . The function  $f(x) = \sin(\pi x)$  is an optimal function from a direct computation.

- Page 201, line -7: instead of

$$\int_K (f_\ell^2 - \frac{1}{\mu(K)} \int_K f_\ell d\mu)^2 d\mu,$$

read

$$\int_K (f_\ell - \frac{1}{\mu(K)} \int_K f_\ell d\mu)^2 d\mu,$$

(thanks to Arnak Dalalyan).

- Page 205, Proposition 4.6.4: instead of

$$C_{K \cup L} \leq \frac{\mu(K \cap L)}{\mu(K \cup L)} \max(C_K, C_L),$$

read

$$C_{K \cup L} \leq 2 \frac{\mu(K \cap L)}{\mu(K \cup L)} \max(C_K, C_L),$$

(thanks to Michał Strzelecki).

- Page 211, line 5: instead of  $\Gamma(P_t f) = O(t^{-1/2})$  read  $\sqrt{\Gamma(P_t f)} = O(t^{-1/2})$ .
- Page 240, line -1: instead of  $s = \int_E f d\mu$  read  $s = f$ , thanks to Michał Strzelecki.
- Page 249, Proposition 5.2.7: instead of  $\mathbb{E}_1, \mathbb{E}_2$  read  $E_1, E_2$ .
- Page 251, formula (5.3.2), read  $(q-1)^{k/2}$  instead of  $(q-1)^k$ , thanks to Max Fathi.
- Page 263, line 11: instead of  $\Lambda^{q-1}(s)$  in the LHS, read

$$\frac{q^2}{q'} \Lambda^{q-1}(s) \Lambda'(s),$$

moreover the function  $q$  is decreasing, thanks to Michał Strzelecki.

- Page 267, line -7: instead of

$$\frac{d(x, y)}{2t}$$

in the RHS, read

$$\frac{d(x, y)}{2\sqrt{t}},$$

thanks to Michał Strzelecki.

- Page 298, line -8: instead of

$$P_t(f \log f) - P_t f \log P_t f \leq t \Delta P_t f + \frac{n}{2} \log(1 - \frac{2t P_t(f \Delta(\log f))}{n P_t f}),$$

read

$$P_t(f \log f) - P_t f \log P_t f \leq t \Delta P_t f + \frac{n}{2} P_t f \log(1 - \frac{2t P_t(f \Delta(\log f))}{n P_t f}).$$

- Page 298. The proof of Theorem 6.7.3 can be simplified as follows.

Let  $f$  be a nonnegative function and let, as usual, for  $s \in [0, t]$

$$\Lambda(s) = P_s(P_{t-s}f \log P_{t-s}f).$$

As already observed,

$$\Lambda'(s) = P_s(P_{t-s}f \Gamma(\log P_{t-s}f)),$$

$$\Lambda''(s) = 2P_s(P_{t-s}f \Gamma_2(\log P_{t-s}f))$$

and the  $CD(0, n)$  condition yields the inequality (6.7.6) page 300,

$$\Lambda''(s) \geq \frac{2}{nP_{t-s}f} [LP_{t-s}f - \Lambda'(s)]^2.$$

Now, letting  $\varphi(s) = \Lambda(s) - sLP_{t-s}f$ , the previous inequality can be reformulated as,

$$\varphi''(s) \geq \frac{2}{nP_{t-s}f} (\varphi'(s))^2, \quad s \in [0, t].$$

In other words, the map

$$[0, t] \ni s \mapsto \exp\left(-\frac{2}{nP_{t-s}f} \varphi(s)\right)$$

is concave.

Then the two inequalities hold true:

$$-\frac{2}{nP_{t-s}f} \varphi'(t) \exp\left(-\frac{2}{nP_{t-s}f} \varphi(t)\right) \leq \frac{\exp\left(-\frac{2}{nP_{t-s}f} \varphi(t)\right) - \exp\left(-\frac{2}{nP_{t-s}f} \varphi(0)\right)}{t} \leq -\frac{2}{nP_{t-s}f} \varphi'(0) \exp\left(-\frac{2}{nP_{t-s}f} \varphi(0)\right).$$

The first inequality can be written as

$$P_t \left( \frac{\Gamma(f)}{f} \right) - LP_{t-s}f + \frac{n}{2t} P_{t-s}f \geq \frac{n}{2t} P_{t-s}f \exp\left(-\frac{2}{nP_{t-s}f} (\varphi(0) - \varphi(t))\right),$$

which is a reformulation of inequality (6.7.4), and the second one can be written as

$$-\frac{\Gamma(P_{t-s}f)}{P_{t-s}f} + LP_{t-s}f + \frac{n}{2t} P_{t-s}f \geq \frac{n}{2t} P_{t-s}f \exp\left(-\frac{2}{nP_{t-s}f} (\varphi(t) - \varphi(0))\right),$$

which is a reformulation of inequality (6.7.5). We recover the Li-Yau inequality since the exponential is positive.

- Page 301, line 11: instead of

$$\Lambda''(s) \geq \frac{2[LP_{t-s}f - \Lambda'(s)]^2}{nP_{t-s}f} + \rho \Lambda'(s),$$

read

$$\Lambda''(s) \geq \frac{2[LP_{t-s}f - \Lambda'(s)]^2}{nP_{t-s}f} + 2\rho \Lambda'(s).$$

- Page 308, additional information on Theorem 6.8.3. For all the computations explained on page 309, the extremal function  $f$  has to satisfy some properties.

First, from the identity

$$\int (f^{q-1} - (1 + \epsilon)f)u d\mu = C\mathcal{E}(f, u),$$

we get

$$\int f^{q-1}u d\mu = C \int f \left( \frac{1 + \epsilon}{C}u - Lu \right) d\mu.$$

That is, if  $R_\lambda(u) = g$  with  $\lambda = \frac{1+\epsilon}{C}$ , the equality becomes

$$\int (R_\lambda(f^{q-1}) - Cf)g d\mu = 0.$$

This equation implies back that

$$f = \frac{1}{C}R_\lambda(f^{q-1})$$

and then,  $f \in \mathcal{D}(L)$ .

It is proved that  $f$  is bounded from above and below (by a strictly positive constant). From the equation satisfied by  $f$ , we know that  $Lf$  is also bounded. To apply the various integration by parts formula, we need to prove that for any constant  $a \in \mathbb{R}$ ,  $f^a \in \mathcal{D}(L)$ . One way to prove it is to show that  $\Gamma(f)$  is a bounded function.

From the first formula page 312, we have

$$f = \frac{1}{C}R_\lambda(f^{q-1}),$$

which implies that

$$\sqrt{\Gamma(f)} \leq \frac{1}{C} \int_0^\infty e^{-\lambda t} \sqrt{\Gamma(P_t(f^{q-1}))} dt.$$

Now, since the model satisfies the  $CD(0, \infty)$  condition and  $f^{q-1}$  is a bounded function, Inequality 4.7.7 page 211 implies that

$$\sqrt{\Gamma(P_t(f^{q-1}))} \leq \frac{\|f^{q-1}\|_\infty}{\sqrt{t}}, \quad t > 0.$$

The two previous inequalities imply that  $\Gamma(f)$  is a bounded function.

- Page 315, formula (6.9.2): instead of  $\hat{L}$ , read  $\hat{L}(f)$ .
- Page 317, line 12: instead of  $\nabla W(f)$ , read  $\Gamma(W, f)$ .
- Page 318, line 13: instead of  $\mu$ , read  $\mu_g$ .
- Page 321, Proposition 6.9.6. The proposition and the proof have to be replaced by the following (see also [1] for a more developed proof).

**Proposition 6.9.6** *Let  $d\mu = e^{-W}d\mu_{\mathfrak{g}}$  and  $\alpha \in \mathbb{R}$ , then*

$$S_{\alpha}(\mu, \Gamma) = \gamma_n(\alpha)[sc_{\mathfrak{g}} - \alpha\Delta_{\mathfrak{g}}W + \beta_n(\alpha)\Gamma(W)]$$

*is  $n$ -conformal invariant where*

$$\beta_n(\alpha) = \frac{\alpha(n - 2n_0 + 2) - 2(n_0 - 1)}{2(n - n_0)}$$

*and*

$$\gamma_n(\alpha) = \frac{n - 2}{4(n_0 - 1) - 2\alpha(n - n_0)}.$$

**Proof**

$\triangleleft$  It is enough to check that  $S_{\alpha}(\mu, \Gamma)$  satisfies the condition (6.9.1). The measure  $\mu$  is transformed to  $\hat{\mu} = c^{-n}\mu$ , and  $\Gamma$  to  $\hat{\Gamma} = c^2\Gamma$ . From the previous computations,  $sc_{\mathfrak{g}}$  becomes

$$\hat{sc}_{\mathfrak{g}} = c^2[sc_{\mathfrak{g}} + (n_0 - 1)(2\Delta_{\mathfrak{g}}\tau - (n_0 - 2)\Gamma(\tau))],$$

$W = -\log \frac{d\mu}{d\mu_{\mathfrak{g}}}$  becomes

$$\hat{W} = -\log \frac{d\hat{\mu}}{d\hat{\mu}_{\mathfrak{g}}} = -\log \frac{c^{-n}d\mu}{c^{-n_0}d\mu_{\mathfrak{g}}} = -\log c^{n_0-n} \frac{d\mu}{d\mu_{\mathfrak{g}}} = W + (n - n_0)\tau,$$

and finally,  $\Delta_{\mathfrak{g}}$  becomes

$$\hat{\Delta}_{\mathfrak{g}} = c^2[\Delta_{\mathfrak{g}} - (n_0 - 2)\Gamma(\tau, \cdot)].$$

So,

$$\begin{aligned} S_{\alpha}(c^{-n}\mu, c^2\Gamma) &= c^2\gamma_n(\alpha) \left[ sc_{\mathfrak{g}} + [2(n_0 - 1) - \alpha(n - n_0)]\Delta_{\mathfrak{g}}(\tau) \right. \\ &\quad + [\beta_n(\alpha)(n - n_0)^2 - (n_0 - 1)(n_0 - 2) + \alpha(n_0 - 2)(n - n_0)]\Gamma(\tau) \\ &\quad \left. - \alpha\Delta_{\mathfrak{g}}(W) + [\alpha(n_0 - 2) + 2\beta_n(\alpha)(n - n_0)]\Gamma(\tau, W) + \beta_n(\alpha)\Gamma(W) \right]. \end{aligned}$$

It has to be equal to

$$c^2 \left[ \gamma_n(\alpha)[sc_{\mathfrak{g}} - \alpha\Delta_{\mathfrak{g}}(W) + \beta_n(\alpha)\Gamma(W)] + \frac{n-2}{2} \left( \Delta_{\mathfrak{g}}(\tau) - \Gamma(W, \tau) - \frac{n-2}{2}\Gamma(\tau) \right) \right].$$

On can check the values of  $\gamma_n(\alpha)$  and  $\alpha_n(\alpha)$  proposed do the job.  $\triangleright$

- Page 338, line -1: instead of  $I(u)$ , read  $I_{\mu, F}(u)$ .
- Page 364, line -6: The sentence starting by *In the finite measure case...* is not correct. It has to be replaced by the following one: In the finite measure case, the tight Nash inequality (3.2.3), p. 281, corresponds to a function  $\Phi$  which is the inverse function of  $(1, +\infty) \ni x \mapsto (x^{1+2/n} - x)/C$ .

- Page 372, line -5: instead of  $e^{-C/t}$ , read  $e^{-t/C}$ .
- Page 373, line -13: instead of  $w(x) = p(x)^{1/2}(1+x^2)^{-\beta}$ , read  $w(x) = p(x)^{-1/2}(1+x^2)^{-\beta}$  (thanks to Persi Diaconis).
- Page 425, Theorem 8.6.3: the set  $A_{d_t}$  should be here the  $d_t$ -closed neighborhood of  $A$  instead of the open one ( $A_{d_t} = \{x \in E; d(x, A) \leq d_t\}$  instead of  $A_{d_t} = \{x \in E; d(x, A) < d_t\}$ ).
- Page 448, line -7: (the line before formula (9.3.5)) the integration is w.r.t. the measure  $u^{1-1/n}dx$  instead of  $udx$  (thanks to Emanuel Milman).
- Page 464, formula (9.7.4) should be

$$W_2^2(P_t f \mu, P_t g \mu) \leq W_2^2(f \mu, g \mu) + 2n(\sqrt{t} - \sqrt{s})^2,$$

thanks to Luigia Ripani.

- Page 516, in the formula (C.6.5) the last term should be

$$H(f_i)(f_j, f_l)$$

instead of

$$H(f_i)(f_i, f_l)$$

thanks to François Bolley.

- Page 514, line -2: instead of wrapped product, read, of course, warped products !

## References

- [1] L. Dupaigne, I. Gentil, S. Zugmeyer. A conformal geometric point of view on the Caffarelli-Kohn-Nirenberg inequality. Preprint 2021.