

Recall that a simple random walk on a graph is called *recurrent* if it returns to the starting point with probability 1, and *transient* otherwise.

**Exercise 1.** *General knowledge*

Let  $G$  be a general graph (hence locally finite for the scope of this course: every vertex has finite degree), let  $v$  be a vertex of  $G$  and  $(S_n)_{n \geq 0}$  be a simple random walk starting at  $v$ . We denote by  $\mathbb{P}_v$  the corresponding probability measure.

1. Explain what a simple random walk on  $G$  is.

**Solution.** A Markov process which jumps at each time, independantly from the past, uniformly to one of its neighbours.

2. Prove that  $(S_n)_{n \geq 0}$  is recurrent if and only if

$$\sum_{n=0}^{\infty} \mathbb{P}_v(S_n = v) = \infty.$$

**Solution.** We first prove that a simple random walk on  $G$  is recurrent if and only if the probability of return to the origin is 1, i.e. if and only if the expected number of returns to the origin is  $\infty$ . If  $\tau_v^n$  is the stopping time for the  $n^{\text{th}}$  visit at  $v$ , we have

$$\mathbb{P}(\tau_v^n < \infty) = \mathbb{P}(\tau_v^n < \infty | \tau_v^{n-1} < \infty) \mathbb{P}(\tau_v^{n-1} < \infty) = \mathbb{P}(\tau_v^1 < \infty) \mathbb{P}(\tau_v^{n-1} < \infty),$$

where we used the Markov property in the last equality, hence by recurrence :  $\mathbb{P}(\tau_v^n < \infty) = \mathbb{P}(\tau_v^1 < \infty)^n$ .  
If  $N_v$  is the number of visits at  $v$ , we also have the identity

$$N_v = \sum_{n=1}^{\infty} \mathbf{1}_{\{\tau_v^n < \infty\}}$$

where  $\mathbf{1}_{\{\tau_v^n < \infty\}} = 1$  if  $\tau_v^n < \infty$  holds and 0 if not. Therefore on the one hand we have

$$\mathbb{E}_v(N_v) = \sum_{n=1}^{\infty} \mathbb{P}(N_v \geq n) = \sum_{n=1}^{\infty} \mathbb{P}(\tau_v^n < \infty) = \sum_{n=1}^{\infty} \mathbb{P}(\tau_v^1 < \infty)^n = \frac{\mathbb{P}(\tau_v^1 < \infty)}{1 - \mathbb{P}(\tau_v^1 < \infty)}.$$

(In a shorter way, one can say that  $N_v$  is a geometric variable of parameter  $\mathbb{P}(\tau_v^1 = \infty)$ .)

And on the other hand we have

$$\mathbb{E}(N_v) = \mathbb{E} \left[ \sum_{k=0}^{\infty} \mathbf{1}_{\{S_k = v\}} \right] = \sum_{n=0}^{\infty} \mathbb{P}_v(S_n = v).$$

This allows to conclude that  $\sum_{n=0}^{\infty} \mathbb{P}_v(S_n = v) = \infty$  if and only if  $\mathbb{P}(\tau_v^1 < \infty) = 1$ .

3. Let us suppose that  $G$  is connected and  $v, w$  are vertices of  $G$ .
  - (a) Show that the simple random walk on  $G$  is recurrent when started from  $v$  if and only if it is recurrent when started from  $w$ .

**Solution.** Since  $G$  is connected (and locally finite), the probability that  $(S_n)_{n \geq 0}$  goes to  $w$  before going back to  $v$  is strictly positive. Suppose  $E_k$  is the event that our walk started from  $v$  passes through  $w$  during the  $k$ -th excursion from  $v$  before coming back the  $(k+1)$ -th time. By the Markov property,  $E_k$ 's are independent and has the same non-zero probability. Therefore, an infinite number of  $E_k$  happens: with probability 1 the walk from  $v$  passes through  $w$  infinitely many times.

Let us consider the random walk after it goes through  $w$  for the first time. Because of the previous discussion, it visits  $w$  an infinite number of times, but it has also (by the Markov property) the same law as the random walk which starts at  $w$ . So we conclude that the simple random walk on  $G$  is recurrent when started from  $w$ .

**Solution.** Other solution : we know that there exists an integer  $k$  such that we can go from  $v$  to  $w$  in  $k$  steps. Then for any  $n \geq 2k$ , if we consider the paths which go from  $w$  to  $v$  in  $k$  steps, then do  $n - 2k$  steps and come back to  $v$  then  $k$  steps to come back to  $w$ , we get:

$$\mathbb{P}_w(S_n = w) \geq \mathbb{P}_w(S_k = v) \mathbb{P}_v(S_{n-2k} = v) \mathbb{P}_v(S_k = w)$$

which after summation gives :

$$\sum_n \mathbb{P}_w(S_n = w) \geq \mathbb{P}_w(S_k = v) \sum_n \mathbb{P}_v(S_{n-2k} = v) \mathbb{P}_v(S_k = w) = \infty$$

which allows to conclude.

- (b) Show that if the simple random walk  $(S_n)_{n \geq 0}$  on  $G$  is recurrent when started from  $v$ , then for any vertex  $w$  of  $G$ ,  $\mathbb{P}_v(\exists n, S_n = w) = 1$  and  $\mathbb{P}_w(\exists n, \tilde{S}_n = v) = 1$  where  $(\tilde{S}_n)_{n \geq 0}$  is a simple random walk on  $G$  starting at  $w$ .

**Solution.** The first assertion was proven in the proof of the previous point 3.(a). For the second, by the result of 3.(a),  $w$  is also recurrent, thus we can apply the first assertion and permuting the role of  $v$  and  $w$ : this gives us the second assertion.

4. Show that a simple random walk on a finite graph is recurrent.

**Solution.** We can restrict ourself to the case where  $G$  is finite and connected. The walk must be somewhere in  $G$  at any time  $n$  thus  $\sum_{v \in G} \mathbb{1}(S_n = v) = 1$ , and therefore

$$\sum_{n \in \mathbb{N}} \sum_{v \in G} \mathbb{1}(S_n = v) = \infty.$$

Yet,  $\sum_{n \in \mathbb{N}} \sum_{v \in G} \mathbb{1}(S_n = v) = \sum_{v \in G} \sum_{n \in \mathbb{N}} \mathbb{1}_{S_n=v} = \sum_{v \in G} N_v = \infty$ , where  $N_v$  is the number of visits to  $v$ . Since  $G$  is finite, there exists  $w$  such that  $N_w = \infty$  with a positive probability. If we consider the random walk conditionned to visit  $w$  (which is possible since we know that  $N_w = \infty$  and therefore  $\mathbb{P}(\exists n, S_n = w) > 0$ ), the law of the walk after the first visit to  $w$  is the simple random walk which starts at  $w$ . It visits  $w$  infinitely many times as the simple random walk is recurrent when started from  $w$  and because of the part 3. the simple random walk is recurrent when started from any vertex of  $G$ .

5. Show that a simple random walk  $(S_n)_{n \geq 0}$  on  $\mathbb{Z}^d$  is recurrent if and only if

$$\sum_{n=0}^{\infty} \mathbb{P}_{\vec{0}}(S_{2n} = \vec{0}) = \infty.$$

**Solution.** This is due to the fact that for any integer  $n$ ,  $\sum_{i=0}^d S_n^i$  has the same parity as  $n$ . Thus  $S_k = \vec{0}$  can be true only if  $k \in 2\mathbb{N}$ . One concludes using the point 2.

### Exercise 2. Universality of the recurrence for random walks on $\mathbb{Z}$

Consider a random walk on  $\mathbb{Z}$  defined using identically independant jumps :  $S_n = Z_1 + \dots + Z_n$  ( $Z_i$  are i.i.d.  $\mathbb{Z}$ -valued random variables). Let us suppose that  $Z_1$  satisfies  $\mathbb{E}(|Z_1|) < \infty$ .

1. Prove that if  $\mathbb{E}(Z_1) \neq 0$  then  $S_n$  is transient.
2. What is the derivative of  $\phi(t) = \mathbb{E}(e^{itZ_1})$  at 0 ? Give the Taylor expansion of  $\phi(t)$  at 0 at order 1.
3. Using the previous point, prove that if  $Z_1$  is symmetric ( $-Z_1$  has the same law as  $Z_1$ ) then  $S_n$  is recurrent.  
*Hint: use the derivation using the Fourier transform as seen in the lesson.*

**Solution.**

1. Without loss of generality, suppose  $\mathbb{E}(Z_1) = \mu > 0$ . By the law of large numbers  $\frac{S_n}{n} \xrightarrow{a.s.} \mathbb{E}(Z_1)$ . Therefore,  $\forall \epsilon > 0$ ,

$$\mathbb{P}(\exists N > 0, \forall n > N, S_n < n(\mu - \epsilon)) \geq \mathbb{P}\left(\exists N > 0, \forall n > N, \left|\frac{S_n}{n} - \mu\right| < \epsilon\right) = 1$$

Therefore, taking  $\epsilon = \frac{\mu}{2}$  gives  $\mathbb{P}(\exists N > 0, \forall n > N, S_n < \frac{n\mu}{2}) = 1$ . Hence  $S_n$  almost surely only visits any integer finitely many times.

2. Using the derivation under the integral (which holds because  $\mathbb{E}(|Z_1|) < \infty$ ) we have  $\phi'(0) = i\mathbb{E}(Z_1)$  and so the Taylor expansion is

$$\phi(x) = 1 + ix\mathbb{E}(Z_1) + o(x).$$

3. For this question, we refer to the lesson for the whole solution. Let  $P_1$  be the law of the jumps of  $S_n$  (thus the law of  $Z_1$ ). Using the lesson, we know that

- (a) in order to show that  $S_n$  is recurrent, we prove that  $\sum_k r^k \mathbb{P}(X_k = 0) \xrightarrow{r \rightarrow 1} \infty$ ,
- (b)  $\sum_k r^k \mathbb{P}(X_k = 0) = \sum_k r^k \mathcal{F}^{-1}(\mathcal{F}(P_k))(0)$ , where  $P_k(\cdot) = \mathbb{P}(S_k = \cdot)$ , and  $\mathcal{F}$  and  $\mathcal{F}^{-1}$  are respectively the Fourier and the inverse fourier transform,
- (c)  $\hat{P}_k = \hat{P}_1^{*k} = \left(\hat{P}_1\right)^k$  where  $\hat{f} = \mathcal{F}(f)$ ,
- (d)  $\hat{P}_1(\xi) = \mathcal{F}(\mathbb{P}(Z_1 = \cdot))(\xi) = \sum_n e^{in\xi} \mathbb{P}(Z_1 = n) = \mathbb{E}(e^{iZ_1\xi})$
- (e) thus  $\sum_k r^k \mathbb{P}(X_k = 0) = \frac{1}{2\pi} \int_{-\pi}^{\pi} \sum_k r^k \mathbb{E}(e^{i\xi Z_1})^k = \frac{1}{2\pi} \int_{-\pi}^{\pi} \frac{1}{1 - r\mathbb{E}(e^{i\xi Z_1})} d\xi$ .
- (f) As  $\sum_k r^k \mathbb{P}(X_k = 0)$  is real we automatically have  $\frac{1}{2\pi} \int_{-\pi}^{\pi} \frac{1}{1 - r\mathbb{E}(e^{i\xi Z_1})} d\xi \in \mathbb{R}$ . Furthermore, as  $Z_1$  is symmetric, the intergrand  $\mathbb{E}(e^{i\xi Z_1}) \in \mathbb{R}$ . Therefore the dampening factor  $r$  does imply convergence, justifying the previous swapping of the intergral and sum.
- (g) The only possibility for the divergence of this integral (as  $r \rightarrow 1$ ) is when  $\mathbb{E}(e^{i\xi Z_1}) = 1$ , that is when  $\xi = 0$ . Actually, we only need to understand the nature of the integral :

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} \frac{1}{1 - \mathbb{E}(e^{i\xi Z_1})} d\xi.$$

Let us remark that  $\mathbb{E}(e^{i\xi Z_1}) \leq 1$  thus the integrand is positive : we only need to consider the integral

$$\frac{1}{2\pi} \int_0^{\pi} \frac{1}{1 - \mathbb{E}(e^{i\xi Z_1})} d\xi.$$

Using Point 2 of this exercise, this integral is of the form

$$\frac{1}{2\pi} \int_0^{\pi} \frac{1}{1 - 1 - i\xi\mathbb{E}(Z_1) + o(\xi)} d\xi = \frac{1}{2\pi} \int_0^{\pi} \frac{1}{o(\xi)} d\xi$$

where we recall that the integrand is positive. This allows us to conclude : there exists  $\epsilon > 0$  such that for any  $\xi \leq \epsilon$ ,  $0 \leq o(\xi) \leq \xi$  thus  $\frac{1}{o(\xi)} \geq \frac{1}{\xi}$  on  $[0, \epsilon]$  and thus

$$\frac{1}{2\pi} \int_0^{\pi} \frac{1}{o(\xi)} d\xi \geq \frac{1}{2\pi} \int_0^{\epsilon} \frac{1}{\xi} = \infty.$$