

RIEMANNIAN CURL IN CONTACT GEOMETRY

SOFIANE BOUARROUDJ AND VALENTIN OVSIENKO

ABSTRACT. We consider a contact manifold with a pseudo-Riemannian metric and define a contact vector field intrinsically associated to this pair of structures. We call this new differential invariant the contact Riemannian curl. We show that the contact Riemannian curl vanishes if the metric is of constant curvature and the contact structure is defined by a Killing 1-form; the contact Riemannian curl has a strong similarity with the Schwarzian derivative since it depends only on the projective equivalence class of the metric. For the Laplace-Beltrami operator on a contact manifold, the contact Riemannian curl is proportional to the subsymbol defined in [CO12]. We also show that the contact Riemannian curl vanishes on the (co)tangent bundle over a Riemannian manifold. This implies that the corresponding subsymbol of the Laplace-Beltrami operator is identically zero.

1. INTRODUCTION

The principal object of this paper is related to the notion of *invariant differential operator*, i.e., the operators commuting with the action of the group of diffeomorphisms. The notion of *differential invariant* is perhaps the oldest notion of differential geometry. The best known example is perhaps the curvature in all its avatars. The domain to which belongs the present work was initiated by Veblen [Veb22] who started a systematic study of invariant differential operators on smooth manifolds. The theory was intensively studied in the 80's in the context of Gelfand-Fuchs cohomology; see [Fu86, GLS02] and references therein.

We consider a smooth manifold M equipped simultaneously with a contact structure and a pseudo-Riemannian metric. We will present a construction of a contact vector field corresponding to these two structures; this vector field will be called the *contact Riemannian curl*. Our construction will be coordinate free and invariant with respect to the action of the group of contact diffeomorphisms, i.e., the contact Riemannian curl is a differential invariant. Moreover, our goal is to define this differential invariant in a “most symmetric” way, so that it is also invariant with respect to natural equivalence relations.

One of the equivalence relations we consider is as follows. Two metrics are called *projectively equivalent* (or geodesically equivalent) if they have the same non-parametrized geodesics. i.e., their Levi-Civita connections are projectively equivalent connections. It turns out that the constructed contact Riemannian curl is invariant with respect to this equivalence relation. This invariance implies, in particular, that the contact Riemannian curl of a metric of constant scalar curvature and contact structure defined by a Killing 1-form vanishes. Invariance of the contact Riemannian curl relative projectively equivalence of metrics makes this notion quite similar to that of classical *Schwarzian derivative* (see [BO00, OT05, B06] for various multi-dimensional generalizations of the Schwarzian derivative). We investigate this relation in more details.

Among the main properties of the contact Riemannian curl that we investigate is its relation to the Laplace-Beltrami operator. Differential operators on contact manifolds were studied from the geometric point of view the recent work [CO12], where the notion of *subsymbol* of a differential operator on a contact manifold was introduced. The subsymbol of a differential operator is a tensor

Key words and phrases. Contact geometry, Riemannian geometry, differential invariants.

field of degree lower than that of the principal symbol. Note that the subsymbol is not well-defined for an arbitrary manifold, one needs a contact structure to obtain an invariant definition. For a given second order differential operators, the subsymbol is just a contact vector field. In the present paper, we consider the *Laplace-Beltrami operator* associated to an arbitrary metric on a contact manifold and calculate its subsymbol. It turns out that it is proportional to the contact Riemannian curl.

We also apply our general construction to a particularly interesting example of a manifold that has natural contact and Riemannian structures, namely to the spherical (or projectivized) cotangent bundle ST^*M over a Riemannian manifold (M, g) . The manifold ST^*M is equipped with the canonical lift of the metric g . We show that the contact Riemannian curl, and therefore the subsymbol of the Laplace-Beltrami operator, is identically zero in this case. Let us mention that the projectivization of the cotangent bundle over a Riemannian manifold M , as well as the sphere bundle ST^*M , is an example of a “real-complex” manifold whose local invariants were recently introduced and computed in [BGLS12].

In the end of the paper, we provide several concrete examples of the contact Riemannian curl. For instance, we calculate it for the 3-ellipsoid equipped with the conformally flat metric introduced in [Tab99] and intensively used in [MT01, DV11].

We believe that the study of differential invariants of a pair (a Riemannian metric, a contact structure) is worth a systematic study.

2. CONTACT GEOMETRY AND TENSOR FIELDS

Contact geometry is an old classical subject, that can be viewed as an odd-dimensional version of symplectic geometry. Let M be a contact manifold and $\dim(M) = 2\ell + 1$, we will always assume that $\ell \geq 1$. Unlike a symplectic structure in symplectic geometry, a contact structure on M is not defined by a differential form. A contact structure is defined by a *conformal class* of a 1-form θ such that $d\theta$ is a 2-form of rank 2ℓ , called a *contact form*. It is important that a contact form is not intrinsically associated with a contact structure.

A contact diffeomorphism (a contact vector field) is a diffeomorphism (a vector field) preserving the contact structure. It does not necessarily preserve a given contact form. The space of all contact vector fields can be identified with the space of smooth functions, but this correspondence depends on the choice of a contact form; see [Arn89].

In this section, we recall several standard facts of contact geometry — those of contact structure and contact vector fields — using somewhat unconventional notation of [OT05, Ovs06] which are among our main references. The contact structure is defined as a tensor field which is a *weighted contact form*. Contact vector fields are in one-to-one correspondence with weighted densities of weight $-\frac{1}{\ell+1}$.

2.1. Weighted densities. A weighted density is a standard object in differential geometry. In order to make the definitions intrinsic, we recall here this notion.

Let M be a manifold of dimension n . For any $\lambda \in \mathbb{R}$, we denote by $\Lambda^n T^*M^{\otimes \lambda}$ the line bundle of homogeneous complex valued functions of weight λ on the determinant bundle $\Lambda^n TM$. The space $\mathcal{F}_\lambda(M)$ of smooth sections of $\Lambda^n T^*M^{\otimes \lambda}$ with complex coefficients is called the space of *weighted densities* of weight λ , (or λ -densities for short).

Example 2.1.1. If the manifold M is orientable and if ω is a volume form, then $\phi\omega^\lambda$, where $\phi \in C^\infty(M)$, is a λ -density.

The space $\mathcal{F}_\lambda(M)$ has the structure of a module over the Lie algebra $\text{Vect}(M)$ of all smooth vector fields on M . We denote by Div the divergence operator associated with a volume form ω

on M . That is, $L_X(\omega) = \text{Div}(X)\omega$. The action of a vector fields reads as follows:

$$(1) \quad L_X^\lambda(\phi\omega^\lambda) = (X(\phi) + \lambda\text{Div}(X)\phi)\omega^\lambda,$$

for every vector field X and $\phi \in C^\infty(M)$.

2.2. Contact manifolds. A smooth manifold M is called *contact* if it is equipped with a completely non-integrable distribution

$$\mathcal{D} \subset TM$$

of codimension 1. The distribution \mathcal{D} is called a *contact distribution*; the hyperplane $\mathcal{D}_x \subset T_xM$ is called a *contact hyperplane* for every point $x \in M$. Contact structure on M exists only if M is odd-dimensional.

A usual way to define a contact structure is to chose a (locally defined) differential 1-form θ on M such that $\mathcal{D} = \ker \theta$. Such a 1-form is called a *contact form*. The complete non-integrability of the distribution \mathcal{D} is equivalent to the fact that

$$\text{vol} := \theta \wedge (d\theta)^\ell$$

is a (locally defined) volume form; equivalently, the 2-form $d\theta$ is a non-degenerate on the contact hyperplanes \mathcal{D}_x of \mathcal{D} . However, there is no canonical choice of a contact form. Moreover, the above definition is not invariant.

A diffeomorphism $f : M \rightarrow M$ is a *contact diffeomorphism* if f preserves \mathcal{D} . If θ is a contact form corresponding to the contact distribution \mathcal{D} and f is a contact diffeomorphism, then f does not necessarily preserve θ , more precisely, $f^*\theta = F_f\theta$, where F_f is a function.

We refer to [Arn89, Bla10] for excellent textbooks on contact geometry.

2.3. The contact tensor. We are ready to introduce the main notion of this section.

Definition 2.3.1. Given a contact form θ , let the *contact tensor field* be

$$(2) \quad \Theta = \theta \otimes \text{vol}^{-\frac{1}{\ell+1}},$$

where $\text{vol}^{-\frac{1}{\ell+1}}$ is a weighted density of weight $-\frac{1}{\ell+1}$, and tensor product is performed over $C^\infty(M)$.

Proposition 2.3.2. *The tensor field Θ is globally defined on a contact manifold M , it is independent of the choice of a contact form, and it is invariant with respect to the contact diffeomorphisms.*

Proof. Let F be a non-vanishing function and consider the contact form $F\theta$. The corresponding volume form is $F\theta \wedge (d(F\theta))^\ell = F^{\ell+1}\theta \wedge (d\theta)^\ell$. Therefore, the contact tensor fields defined by Eq. (2), corresponding to the contact forms θ and $F\theta$, coincide. Hence, Θ is globally defined and invariant with respect to contact diffeomorphisms. \square

A contact structure is intrinsically defined by the corresponding contact tensor.

Example 2.3.3. Local coordinates $(x^1, \dots, x^\ell, y^1, \dots, y^\ell, z)$ on M are often called the *Darboux coordinates* if the contact structure can be represented by the 1-form

$$\theta_{\text{Dar}} = dz + \frac{1}{2} \sum_{i=1}^{\ell} (x^i dy^i - y^i dx^i).$$

The corresponding volume form is then the standard one:

$$\text{vol} = (-1)^{\frac{\ell(\ell-1)}{2}} \ell! dx^1 \wedge \dots \wedge dx^\ell \wedge dy^1 \wedge \dots \wedge dy^\ell \wedge dz.$$

A contact structure has no local invariants, therefore Darboux coordinates always exist in the vicinity of every point; see [Arn89] (and [GL07] for a simple algebraic proof).

2.4. Contact vector fields. A *contact vector field* on a contact manifold M is a vector field that preserves the contact distribution. This is usually expressed in terms of contact forms: a vector field X is contact if, for every contact form θ , the Lie derivative $L_X\theta$ is proportional to θ :

$$(3) \quad L_X\theta = \frac{1}{\ell+1} \operatorname{Div}(X)\theta.$$

In terms of the contact tensor (2), we have the following corollary of Proposition 2.3.2.

Corollary 2.4.1. *A vector field X is contact if and only if it preserves the contact tensor:*

$$L_X\Theta = 0.$$

Let $\mathcal{K}(M)$ denote the space of all smooth contact vector fields on M . This space has a Lie algebra structure, it is also a module over the group of contact diffeomorphisms. The following observation can be found in [Ovs06, CO12].

Proposition 2.4.2. *As a module over the group of contact diffeomorphisms, the space $\mathcal{K}(M)$ is isomorphic to the space of weighted densities $\mathcal{F}_{-\frac{1}{\ell+1}}(M)$.*

Proof. The space of contact forms is isomorphic to $\mathcal{F}_{\frac{1}{\ell+1}}(M)$. Indeed, this follows from Proposition 2.3.2 and from Eq. (3). The statement then follows from the fact that there is a natural $C^\infty(M)$ -valued pairing between the spaces of contact vector fields and of contact forms: $(X, \theta) \mapsto \theta(X)$. \square

Remark 2.4.3. The above proposition means that, unlike the symplectic geometry, the notion of *contact Hamiltonian* should be understood as a weighted density and not as a function. However, in the Darboux coordinates, the correspondence between the elements of $\mathcal{K}(M)$ and $\mathcal{F}_{-\frac{1}{\ell+1}}(M)$ becomes the usual correspondence between contact vector fields and functions (see [Arn89]):

$$X_{\phi\omega^{-\frac{1}{\ell+1}}} = \sum_{i=1}^{\ell} (\partial_{x^i}(\phi)\partial_{y^i} - \partial_{y^i}(\phi)\partial_{x^i}) + \frac{1}{2}\partial_z(\phi)\mathcal{E} + (\phi - \frac{1}{2}\mathcal{E}(\phi))\partial_z,$$

where

$$\mathcal{E} = \sum_{i=1}^{\ell} (x^i\partial_{x^i} + y^i\partial_{y^i})$$

is the Euler vector field.

Example 2.4.4. If $\dim M = 3$, contact vector fields are identified with $-\frac{1}{2}$ -densities; if $\dim M = 5$, then $\mathcal{K}(M) \cong \mathcal{F}_{-\frac{1}{3}}(M)$, etc. Note also that, in the one-dimensional case, the contact structure is trivial and every vector field is contact, one then has $\operatorname{Vect}(M) \cong \mathcal{F}_{-1}(M)$.

2.5. Another definition of weighted densities on contact manifolds. In presence of a contact structure defined by a contact form θ , it is natural to express elements of any rank 1 bundle, for example, weighted densities, in terms of powers of θ :

$$\phi \operatorname{vol}^{\frac{\lambda}{\ell+1}} \longleftrightarrow \phi\theta^\lambda,$$

where as above ϕ is a smooth function.

Remark 2.5.1. The notation $\phi\theta^\lambda$ is adopted in many works by physicists (see also [Ovs90, GLS01]). In this notation, many formulas simplify. For instance, if X is a contact vector field, then the corresponding contact Hamiltonian is $\phi\theta^{-1}$, where the function ϕ is simply the evaluation $\phi = \theta(X)$.

2.6. The Poisson algebra of weighted densities. The space, $\mathcal{F} = \bigcup_{\lambda} \mathcal{F}_{\lambda}$, of all weighted densities on M can be endowed with a structure of a Poisson algebra (see [Arn89, OT05, Ovs06]):

$$\{.,.\} : \mathcal{F}_{\lambda} \otimes \mathcal{F}_{\mu} \rightarrow \mathcal{F}_{\lambda+\mu+\frac{1}{\varepsilon+1}}.$$

The explicit formula in Darboux coordinates is as follows:

$$\{\phi \omega^{\lambda}, \psi \omega^{\mu}\} = \left(\sum_{i=1}^n (\partial_{x^i} \phi \partial_{y^i} \psi - \partial_{x^i} \psi \partial_{y^i} \phi) + \partial_z \phi (\mu \psi + \mathcal{E} \psi) - \partial_z \psi (\lambda \phi + \mathcal{E} \phi) \right) \omega^{\lambda+\mu+\frac{1}{\varepsilon+1}}.$$

The subspace $\mathcal{F}_{-\frac{1}{\varepsilon+1}}(M)$ is a Lie subalgebra of \mathcal{F} isomorphic to $\mathcal{K}(M)$. The Poisson bracket of $-\frac{1}{\varepsilon+1}$ -densities precisely corresponds to the Lie derivative:

$$X_{\{\phi, \psi\}} = L_{X_{\phi}}^{-\frac{1}{\varepsilon+1}}(\psi),$$

where $\phi = \phi \omega^{-\frac{1}{\varepsilon+1}}$, $\psi = \psi \omega^{-\frac{1}{\varepsilon+1}}$.

2.7. The invariant splitting. The full space of vector fields $\text{Vect}(M)$ splits into direct sum

$$\text{Vect}(M) = \mathcal{K}(M) \oplus \mathcal{T}an(M),$$

where $\mathcal{T}an(M)$ is the space of vector fields tangent to the contact distribution, i.e., $\theta(Y) = 0$ for every contact form θ and every $Y \in \mathcal{T}an(M)$. Such vector fields are called *tangent vector fields*. Unlike $\mathcal{K}(M)$, the space $\mathcal{T}an(M)$ is not a Lie algebra, but a $\mathcal{K}(M)$ -module.

The above splitting is invariant with respect to the group of contact diffeomorphisms. In particular, there is an invariant projection

$$(4) \quad \pi : \text{Vect}(M) \rightarrow \mathcal{K}(M),$$

that will be very useful.

3. THE CONTACT RIEMANNIAN CURL AND ITS PROPERTIES

In this section, we introduce our main notion, a contact vector field corresponding to a metric and a contact structure. We also study its main properties, such as projective invariance and relation to the multi-dimensional Schwarzian derivative.

3.1. Covariant derivative. Let us assume now that M is endowed with a pseudo-Riemannian metric g . We denote the Levi-Civita connection on M by ∇ , and the Christoffel symbols by Γ_{ij}^k . The *covariant derivative*, also denoted by ∇ , is the linear map that can be defined for arbitrary space of tensor fields, $\mathcal{T}(M)$:

$$\nabla : \mathcal{T}(M) \rightarrow \mathcal{T}(M) \otimes \Omega^1(M).$$

It is written in the form $\nabla(t) = \nabla_i(t) dx^i$, and therefore it suffices to define the partial derivatives ∇_i .

The covariant derivative of vector fields and differential 1-forms is given, in local coordinates, by the well-known formulas

$$\nabla_i (V^j \partial_j) = \left(\partial_i V^j + \Gamma_{ik}^j V^k \right) \partial_j, \quad \nabla_i (\beta_j dx^j) = (\partial_i \beta_j - \Gamma_{ij}^k \beta_k) dx^j,$$

respectively, where $\partial_i = \partial/\partial x^i$. Here and below summation over repeated indices is understood (Einstein's notation); see [DNF92]. The covariant derivative then extended to every tensor fields by Leibniz rule.

For instance, the covariant derivative of weighted densities is defined in local coordinates by the following formula:

$$\nabla_i (\phi \omega^{\lambda}) = \left(\partial_i \phi - \lambda \Gamma_{ij}^j \phi \right) \omega^{\lambda},$$

that we will extensively use throughout the paper.

3.2. The main definition. The main notion introduced in this paper is the following one.

Definition 3.2.1. (a) For every pseudo-Riemannian metric g on a contact manifold M , we define a weighted density of degree $-\frac{1}{\ell+1}$:

$$(5) \quad A_{g,\Theta} := \langle g, \nabla\Theta \rangle,$$

in local coordinates, $A_{g,\Theta} := g^{ij}\nabla_i\Theta_j$.

(b) We call the contact vector field $X_{A_{g,\Theta}}$ with contact Hamiltonian $A_{g,\Theta}$ the *contact Riemannian curl of g* .

Note that the quantity $A_{g,\Theta}$ is, indeed, a weighted density of degree $-\frac{1}{\ell+1}$, so that it has a meaning of contact Hamiltonian; see Proposition 2.4.2.

It will be useful to have an explicit expression for $A_{g,\Theta}$ in local coordinates.

Proposition 3.2.2. *In local coordinates, such that $\Theta = \theta \otimes \text{vol}^{-\frac{1}{\ell+1}}$, one has*

$$(6) \quad A_{g,\Theta} = g^{ij} \left(\partial_i\theta_j - \left(\Gamma_{ij}^k - \frac{1}{2(\ell+1)} (\delta_i^k\Gamma_{jr}^r + \delta_j^k\Gamma_{ir}^r) \right) \theta_k \right) \text{vol}^{-\frac{1}{\ell+1}},$$

where δ_i^k is the Kronecker symbol.

Proof. This can be obtained directly from Definition 3.2.1 and the expression of the covariant derivative of a weighted density. \square

Remark 3.2.3. It follows from the intrinsic definition (5) that the local expression (6) is actually invariant with respect to the action of the group of contact diffeomorphisms. The formula (6) remains unchanged for any choice of local coordinates. It is also independent of the choice of the contact form.

3.3. Projective invariance. Let us recall a fundamental notion of projectively equivalent connections due to Cartan [Car24].

A *projective connection* is an equivalence class of symmetric affine connections giving the same non-parameterized geodesics. The *symbol of a projective connection* is given by the expression

$$\Pi_{ij}^k := \Gamma_{ij}^k - \frac{1}{n+1} (\delta_i^k\Gamma_{lj}^l + \delta_j^k\Gamma_{il}^l),$$

where n is the dimension; see [KN64]. Note that in our case, $n = 2\ell + 1$.

The simplest properties of a projective connection are as the following.

- (1) Two affine connections, ∇ and $\tilde{\nabla}$, are projectively equivalent if and only if their symbols Π_{ij}^k and $\tilde{\Pi}_{ij}^k$ coincide.
- (2) Equivalently, ∇ and $\tilde{\nabla}$ are projectively equivalent if and only if there exists a 1-form β such that

$$\tilde{\Gamma}_{ij}^k = \Gamma_{ij}^k + \delta_j^k\beta_i + \delta_i^k\beta_j.$$

It turns out that the contact curl of a metric depends only on the projective equivalence class of the Levi-Civita connection.

The following statement is perhaps the most important property of the contact Riemannian curl that makes it similar to the Schwarzian derivative.

Theorem 3.3.1. *If g and \tilde{g} are two metrics whose Levi-Civita connections are projectively equivalent, then $A_{g,\Theta} = A_{\tilde{g},\Theta}$.*

Proof. The coordinate formula (6) for $A_{g,\Theta}$ becomes can be written as follows:

$$A_{g,\Theta} = g^{ij} (\partial_i \theta_j - \Pi_{ij}^k \theta_k).$$

This expression depends only on the projective class of the Levi-Civita connection and implies projective invariance. \square

Let $[g]$ denote the class of geodesically equivalent metrics, let $[\nabla]$ denote the corresponding projective connection. The above theorem means that the projective Riemannian curl actually depends only on $[g]$ and not on the metric itself.

Remark 3.3.2. Geodesically equivalent metrics is a very classical subject of Riemannian geometry that goes back to Beltrami, Levi-Civita, Weyl, and Cartan. We refer to the classical book [Eis97] for a survey. The subject is still very active, see [BKM09] and references therein.

3.4. Projectively flat connections, metrics of constant curvature and Killing contact forms. It is now natural to investigate projectively flat case.

A connection ∇ on M is called *projectively flat* if, in a neighborhood of every point, there exist local coordinates, often called *adapted coordinates*, such that the geodesics are straight lines in these coordinates.

A connection is projectively flat if in adapted coordinates, the symbols Π_{ij}^k vanish identically. Note also that projectively flat connections admit a (local) action of the group $SL(n+1, \mathbb{R})$, in other words, adapted coordinates admit linear-fractional changes.

The classical Beltrami theorem states that *the Levi-Civita connection of a Riemannian metric is projectively flat if and only if the metric has a constant sectional curvature*. This fact allows us to obtain an important consequence of Theorem 3.3.1.

Let us recall the notion of Killing differential forms that goes back to Yano [Yan52]. A 1-form $\beta = \beta_i(x) dx^i$ is said to be a *Killing form* if

$$\nabla_i \beta_j + \nabla_j \beta_i = 0.$$

Recall also a more common notion of Killing vector field. A vector field $V = V^i(x) \partial_i$ is said to be a *Killing vector field* if

$$L_V g = 0$$

Every Killing 1-form can be obtained from a Killing vector field by lowering indices: $\beta = \langle g, V \rangle$; i.e., $\beta_i = g_{ij} V^j$ in local coordinates.

Corollary 3.4.1. *If g is a metric of constant sectional curvature and if the contact structure is defined by a contact 1-form θ which is a Killing form with respect to a metric from the projective class $[g]$, then $A_{g,\Theta} = 0$.*

Proof. Assume that the Levi-Civita connection corresponding to g is projectively flat. In this case, there exist local coordinates for which $\Pi_{ij}^k \equiv 0$, and therefore

$$A_{g,\Theta} = g^{ij} \partial_i \theta_j.$$

If, furthermore, $\partial_i \theta_j + \partial_j \theta_i = 0$ for all i, j , then $A_{g,\Theta}$ vanishes identically since the tensor g^{ij} is symmetric. The equation $\partial_i \theta_j + \partial_j \theta_i = 0$ means that θ is a Killing form with respect to the flat metric which is projectively equivalent to g .

The corollary then follows from Theorem 3.3.1. \square

Example 3.4.2. The Darboux form in Example 2.3.3 is a Killing form with respect to the flat metric. Note that in other works, especially in those on analytical mechanics, another normal form of the contact form is often used: $dz + \sum_{1 \leq i \leq \ell} x^i dy^i$. (Over fields of characteristic 2, only this latter form can be used, see [Leb10].) However, this is not a Killing form with respect to the flat metric.

3.5. Contact equivariance. Consider the action of the group of contact diffeomorphisms. It immediately follows from the intrinsic (i.e., invariant) definition (5) of $A_{g,\Theta}$ of that the map $g \mapsto A_{g,\Theta}$ from the space of metrics to that of $-\frac{1}{\ell+1}$ -densities commutes with this action:

$$(7) \quad A_{f^*g,\Theta} = f^*(A_{g,\Theta}).$$

From this fact and Corollary 3.4.1, we deduce the following statement.

Corollary 3.5.1. *If a metric \tilde{g} is contactomorphic to a metric g of constant sectional curvature and if the contact structure is defined by a contact 1-form θ which is a Killing form with respect to a metric from the projective class $[g]$, then $A_{\tilde{g},\Theta} = 0$.*

3.6. Action of the full group of diffeomorphisms. Let us consider the action of the group of all diffeomorphisms. It turns out that this action is related to a quite remarkable 1-cocycle.

Recall that the space of connections is an affine space associated with the space of $(2,1)$ -tensor fields, i.e., given two connections, ∇ and $\tilde{\nabla}$, the difference $\nabla - \tilde{\nabla}$ is a well-defined $(2,1)$ -tensor field. This allows one to define a 1-cocycle on the group of all diffeomorphisms. If f is an arbitrary, not necessarily contact, diffeomorphism, we set:

$$C(f) := f^*\nabla - \nabla,$$

where ∇ is an arbitrary fixed connection, choice of which changes C by a coboundary¹.

Let ∇ and $\tilde{\nabla}$ be two connections on M . The difference of the projective equivalence classes $[\nabla] - [\tilde{\nabla}]$ can be understood as a traceless $(2,1)$ -tensor field. Therefore, a projective connection on M leads to the following 1-cocycle on the group of all diffeomorphisms:

$$\mathfrak{T}(f) = f^*[\nabla] - [\nabla]$$

which vanishes on (locally) projective diffeomorphisms. In local coordinates,

$$\mathfrak{T}(f)_{ij}^k := f^*\Pi_{ij}^k - \Pi_{ij}^k,$$

where Π_{ij}^k are the projective Christoffel symbols².

Proposition 3.6.1. *If $f : M \rightarrow M$ is an arbitrary diffeomorphism, then*

$$(8) \quad f^*(A_{g,\Theta}) - A_{f^*g,\Theta} = f^*\langle g, \nabla\Theta \rangle - \langle f^*g, \nabla\Theta \rangle + \langle f^*g \otimes \Theta, \mathfrak{T}(f) \rangle.$$

Proof. Let us first clarify the notation. Since $\mathfrak{T}(f)$ is a $(2,1)$ -tensor field, the pairing $\langle g \otimes \Theta, \mathfrak{T}(f) \rangle$ is well-defined. Furthermore, taking into account the weight of the contact tensor Θ , it follows that $\langle g \otimes \Theta, \mathfrak{T}(f) \rangle$ is a weighted density of weight $-\frac{1}{\ell+1}$.

In local coordinates and using Proposition 3.2.2, we have

$$\begin{aligned} A_{f^*g,\Theta} &= (f^*g)^{ij} (\partial_i\theta_j - f^*\Pi_{ij}^k \theta_k) \\ &= (f^*g)^{ij} (\partial_i\theta_j - (f^*\Pi_{ij}^k - \Pi_{ij}^k) \theta_k) - (f^*g)^{ij} \Pi_{ij}^k \theta_k \\ &= (f^*g)^{ij} (\partial_i\theta_j - \mathfrak{T}(f)_{ij}^k \theta_k) - (f^*g)^{ij} \Pi_{ij}^k \theta_k \\ &= (f^*g)^{ij} (\partial_i\theta_j - \Pi_{ij}^k \theta_k) - (f^*g)^{ij} \mathfrak{T}(f)_{ij}^k \theta_k \\ &= (f^*g)^{ij} \nabla_i(\Theta_j) - (f^*g)^{ij} \mathfrak{T}(f)_{ij}^k \theta_k \\ &= \langle f^*g, \nabla(\Theta) \rangle - \langle f^*g \otimes \Theta, \mathfrak{T}(f) \rangle. \end{aligned}$$

It remains to notice that $f^*(A_{g,\Theta}) = f^*\langle g, \nabla(\Theta) \rangle$. Proposition 3.6.1 is proved. \square

¹Note also that the cocycle C provides a universal way to construct representatives of non-trivial classes of the Gelfand-Fuchs cohomology; see [Gel70].

²The 1-cocycle \mathfrak{T} is often considered as a higher-dimensional analog of the Schwarzian derivative; see [OT05]. If ∇ is projectively flat, then the group $\text{SL}(n+1, \mathbb{R})$ of (local) symmetries of $[\nabla]$ is precisely the kernel of \mathfrak{T} .

4. THE SUBSYMBOL OF THE LAPLACE-BELTRAMI OPERATOR

In this section, we explain the relation of the Riemannian curl to the classical Laplace-Beltrami operator. Let us mention that study of differential operators on contact manifolds is a classical subject; see a recent work [vE10] and references therein.

4.1. Differential operators and diffeomorphism action. Let M be an arbitrary smooth manifold and $\mathcal{D}_{\lambda,\mu}(M)$ be the space of linear differential operators acting on the space of weighted densities:

$$T : \mathcal{F}_\lambda(M) \rightarrow \mathcal{F}_\mu(M).$$

The space $\mathcal{D}_{\lambda,\mu}(M)$ is naturally a module over the group of diffeomorphisms, the module structure being dependent of the weights λ and μ . For $k \in \mathbb{N}$, let $\mathcal{D}_{\lambda,\mu}^k(M)$ be the space of linear differential operators of order $\leq k$. The spaces $\mathcal{D}_{\lambda,\mu}^k(M)$ define a filtration on $\mathcal{D}_{\lambda,\mu}(M)$ invariant with respect to the group of diffeomorphisms.

Recall the classical notion of *symbol* (or the *principal symbol*) of a differential operator of order k . It is defined as the image of the projection

$$\sigma : \mathcal{D}_{\lambda,\mu}(M) \rightarrow \mathcal{D}_{\lambda,\mu}^k(M) / \mathcal{D}_{\lambda,\mu}^{k-1}(M).$$

Observe that, in the particular case $\lambda = \mu$, the quotient space $\mathcal{D}_{\lambda,\lambda}^k(M) / \mathcal{D}_{\lambda,\lambda}^{k-1}(M)$ can be identified with the space of symmetric contravariant tensor fields of degree k on M .

We will be especially interested in the space $\mathcal{D}_{\lambda,\lambda}^2(M)$ of 2-nd order operators acting on λ -densities; a systematic study of this space viewed as a module over the group of diffeomorphisms was initiated in [DO97].

4.2. The subsymbol of a second order differential operator. In [CO12], the space of differential operators on a contact manifold was studied as a module over the group of contact diffeomorphisms. It was proved that there exists a notion of *subsymbol* which is a tensor field of degree lower than that of the principal symbol.

For a 2-nd order differential operator, the subsymbol is just a contact vector field. More precisely, for every λ , there exists a linear map (which is unique up to a constant factor)

$$s\sigma : \mathcal{D}_{\lambda,\lambda}^2(M) \rightarrow \mathcal{K}(M),$$

invariant with respect to the action of the group of contact diffeomorphisms. The image $s\sigma(T)$ was called the *subsymbol* of the operator T . We will need the explicit formula for the subsymbol of a given second order differential operator.

If M is a contact manifold, then every operator $T \in \mathcal{D}_{\lambda,\lambda}^2(M)$ can be written (in many different ways) in the form:

$$(9) \quad T = L_{X_{\phi_1}}^\lambda \circ L_{X_{\phi_2}}^\lambda + L_{X_{\phi_3}}^\lambda \circ L_{Y_1}^\lambda + L_{Y_2}^\lambda \circ L_{Y_3}^\lambda + L_{X_{\phi_4}}^\lambda + L_{Y_4}^\lambda + F,$$

where each Y_i is a vector field tangent to the contact distribution, X_ϕ is the contact vector field with the contact Hamiltonian $\phi \in \mathcal{F}_{-\frac{1}{\ell+1}}(M)$, the Lie derivative L^λ is defined by Eq. (1), and F denotes the operator of multiplication by a function.

The explicit expression for the subsymbol of differential operator (9) is as follows (see [CO12]):

$$(10) \quad s\sigma(T) = \frac{1}{2} [X_{\phi_1}, X_{\phi_2}] - \left(\frac{\ell+1}{\ell+2}\right) \left(\lambda - \frac{1}{2}\right) X_{\tilde{L}_{Y_1}(\phi_3)} + \frac{1}{2} \pi [Y_2, Y_3] + X_{\phi_4},$$

where $\tilde{L}_Y(\phi) := L_Y^{-\frac{1}{\ell+1}}(\phi)$ denotes the Lie derivative of a $-\frac{1}{\ell+1}$ -density ϕ along the vector field Y , and $\pi : \text{Vect}(M) \rightarrow \mathcal{K}(M)$ is defined in (4).

Remark 4.2.1. Although it seems almost impossible, the map $s\sigma$ defined by (10) is well-defined. In other words, it is independent of the choice of the vector fields in the representation (9) of the operator T . This can be checked directly by rewriting it in local coordinates, see formula (12) below. Since the expression (10) is written using invariant terms, it commutes with the action of contact diffeomorphisms. Note also that the existence of such a map is indigenous to contact geometry. There is no similar map commuting with the full group of diffeomorphisms, except for the principal symbol.

4.3. The Laplace-Beltrami operator on the space of weighted densities. The classical Laplace-Beltrami operator acting on the space of smooth functions is defined as follows

$$\Delta_{\mathbf{g}} = g^{ij} \nabla_i \nabla_j.$$

This operator is completely determined by the metric \mathbf{g} .

We will go to a more general framework and consider the generalized Laplace-Beltrami operator acting on the space of weighted densities:

$$\Delta_{\mathbf{g}}^{\lambda} : \mathcal{F}_{\lambda} \rightarrow \mathcal{F}_{\lambda},$$

defined by $\Delta_{\mathbf{g}}^{\lambda} = g^{ij} \nabla_i^{\lambda} \nabla_j^{\lambda}$. The explicit formula of this operator is as follows:

$$\Delta_{\mathbf{g}}^{\lambda}(\phi \omega^{\lambda}) = \left(g^{ij} \nabla_i \nabla_j(\phi) + \frac{n^2 \lambda(\lambda - 1)}{(n - 1)(n + 2)} R \phi \right) \omega^{\lambda},$$

where R is the scalar curvature (see [DO01], Proposition 5.2).

4.4. Calculating the subsymbol of the Laplace-Beltrami operator. Recall that M is a contact manifold and $n = 2\ell + 1$. It turns out that the contact Riemannian curl of a given metric \mathbf{g} is proportional to the subsymbol of the Laplace-Beltrami operator associated with \mathbf{g} . This property can be considered as an equivalent definition of the contact Riemannian curl.

Theorem 4.4.1. *One has*

$$(11) \quad s\sigma(\Delta_{\mathbf{g}}^{\lambda}) = \binom{\ell+1}{\ell+2} (2\lambda - 1) X_{A_{\mathbf{g}}, \Theta}.$$

Proof. The proof is essentially a direct computation.

Let us choose local Darboux coordinates. Every 2-nd order differential operator can be written in these coordinates as:

$$\begin{aligned} T = & T_{2,0,0} \partial_z^2 + T_{1,i,0} \partial_z \partial_{x_i} + T_{1,0,i} \partial_z \partial_{y_i} + T_{0,ij,0} \partial_{x_i} \partial_{x_j} + T_{0,i,j} \partial_{x_i} \partial_{y_j} + T_{0,0,ij} \partial_{y_i} \partial_{y_j} \\ & + T_{1,0,0} \partial_z + T_{0,i,0} \partial_{x_i} + T_{0,0,i} \partial_{y_i} + T_{0,0,0}. \end{aligned}$$

The coordinate formula of the subsymbol was calculated in [CO12]:

$$(12) \quad \begin{aligned} s\sigma(T) = & \frac{1+2\lambda(\ell+1)}{\ell+2} \left(\partial_z (T_{2,0,0} - \frac{1}{2} y_i T_{1,i,0} + \frac{1}{2} x_i T_{1,0,i}) \right. \\ & + \partial_{x_i} (T_{1,i,0} - \frac{1}{2} y_j T_{0,ij,0} + \frac{1}{2} x_j T_{0,i,j}) \\ & \left. + \partial_{y_i} (T_{1,0,i} + \frac{1}{2} x_j T_{0,0,ij} - \frac{1}{2} y_j T_{0,j,i}) \right) \\ & + T_{1,0,0} - \frac{1}{2} y_i T_{0,i,0} + \frac{1}{2} x_i T_{0,0,i}. \end{aligned}$$

One can check that this is exactly the same formula as (10).

The expression of the generalized Laplace-Beltrami operator Δ^{λ} in local coordinates was calculated in [DO01], the result is:

$$\Delta_{\mathbf{g}}^{\lambda} = g^{ij} \partial_i \partial_j - (g^{jk} \Gamma_{jk}^i + 2\lambda g^{ij} \Gamma_{jk}^k) \partial_i + (0 - \text{th order coefficients}).$$

Let us combine the above two formulas. We obtain $\text{sc}(\Delta_g^\lambda) = X_\phi$, where ϕ is a weighted density of the form

$$(13) \quad \phi = \left(\left(1 - \frac{1+2\lambda(\ell+1)}{\ell+2} \right) g^{jk} \Gamma_{jk}^t \theta_t + \left(2\lambda - \frac{1+2\lambda(\ell+1)}{\ell+2} \right) \Gamma_{ij}^j g^{it} \theta_t \right) \text{vol}^{-\frac{1}{\ell+1}},$$

and X_ϕ is the corresponding contact vector field.

Finally, taking into account the fact that $g^{ij} \partial_i(\theta_j) = 0$, for the Darboux form θ , the expression (13), after collecting the terms, coincides with $\left(\frac{\ell+1}{\ell+2} \right) (2\lambda - 1) A_{g, \Theta}$. \square

Corollary 4.4.2. *For a generic metric, $\text{sc}(\Delta_g^\lambda) = 0$ if and only if $\lambda = \frac{1}{2}$.*

Remark 4.4.3. In differential geometry it is known that the space of half-densities and the space of differential operators $\mathcal{D}_{\frac{1}{2}, \frac{1}{2}}(M)$ acting on them play a very special role. In our context, the space of half-densities appears naturally.

5. COTANGENT LIFT AND THE GEODESIC SPRAY

In this section, we calculate the contact Riemannian curl on the unit sphere bundle STM over a Riemannian manifold (M, g) . The manifold STM is a classical example of contact manifold, and, furthermore, it is equipped with the canonical lift of the metric. We prove that the contact Riemannian curl vanishes in this case.

Recall that the classical *geodesic spray* is the Hamiltonian vector field on TM with Hamiltonian $H(x, y) = g_{ij}(x) y^i y^j$, where y^i are coordinates on the fibers; the restriction of this vector field to STM is an intrinsically defined contact vector field. It would be therefore too optimistic to expect another vector field in this situation.

5.1. The coordinates on STM . Let (M, g) be any Riemannian manifold of dimension n . The Riemannian geometry of the sphere bundle STM was studied in [Tah69], we will be using the notation of that work.

Denote by (x^h) a local coordinate system in M and (y^k) the Cartesian coordinates in the tangent space $T_x M$ at the point x in M . The coordinates (x^h, y^h) are local coordinates on the tangent bundle on TM . The unit sphere bundle STM is a hypersurface of the tangent bundle $T(M)$, singled out as the level surface of the Hamiltonian of the geodesic spray

$$H(x, y) = 1$$

with respect to the metric $g(x)$ in the tangent space $T_x(M)$ at every point x .

5.2. Statement of the main result. The Riemannian metric g on M has a canonical lift to STM that will be denoted by \bar{g} . The main result of this section is as follows.

Theorem 5.2.1. *The contact Riemannian curl on STM is identically zero.*

In order to prove this theorem, we will need explicit formulas for the contact structure and the canonical Riemannian metric on STM .

5.3. The contact structure of the sphere bundle STM . The sphere bundle STM is represented by parametric equations:

$$x^h = x^h, \quad x^{\bar{h}} = y^h = y^h(x^i, u^\kappa),$$

where u^κ are local coordinates on the sphere.³ The tangent vectors $B_\alpha^A = \partial x^A / \partial u^\alpha$ of STM in $T(M)$ are given by

$$(14) \quad \begin{aligned} B_i^h &= \delta_i^h, & B_\lambda^h &= 0, \\ B_i^{\bar{h}} &= \partial_i y^h, & B_\lambda^{\bar{h}} &= \partial_\lambda y^h. \end{aligned}$$

The square matrix $\begin{pmatrix} B_\alpha^A \\ C^A \end{pmatrix}$, where $C^i = 0$ and $C^{\bar{i}} = y^i$, is invertible at each point x in M . Its inverse will be the matrix $(B^{\alpha A}, C_A)$, given by the equations:

$$(15) \quad \begin{aligned} B^h_i &= \delta_i^h, & B^h_{\bar{i}} &= 0, \\ B^\kappa_i &= -B_i^{\bar{h}} B^\kappa_{\bar{h}}, & B^\kappa_{\bar{i}} &, \end{aligned}$$

and $C_A = \begin{pmatrix} C_i \\ C_{\bar{i}} \end{pmatrix}$, where $C_{\bar{i}} = g_{ih} y^h$ and $C_i = \Gamma_{rs}^h g_{hi} y^r y^s$. The next formulas can be deduced from Eqs. (14), (15), and are useful for what follows

$$\begin{aligned} B_\lambda^{\bar{h}} B^\kappa_{\bar{h}} &= \delta_\lambda^\kappa, & y^h B^\kappa_{\bar{h}} &= 0, & B_\lambda^{\bar{h}} B^{\lambda_{\bar{i}}} + y^h C_{\bar{i}} &= \delta_i^h, \\ B_\lambda^{\bar{h}} C_{\bar{h}} &= 0, & y^h C_{\bar{h}} &= 1. \end{aligned}$$

The Riemannian metric identifies the tangent bundle $T(M)$ and the cotangent bundle $T^*(M)$, and hence induces a 1-form θ on $T(M)$, called the *Liouville form*, which in local coordinates reads as follows:

$$\theta = g_{ij} y^j dx^i,$$

Denote by $\bar{\theta}$ the restriction of the 1-form θ to the sphere bundle STM . It is as follows:

$$\bar{\theta}_\alpha = \theta_A B_\alpha^A.$$

Eq. (14) imply that $\bar{\theta}_i = g_{ij} y^j$ and $\bar{\theta}_\kappa = 0$.

Lemma 5.3.1. *The form $\bar{\theta}$ defines a contact structure on STM . The volume form associated with it reads (up to a factor) as:*

$$\Omega dx^1 \wedge \dots \wedge dx^n \wedge du^{n+1} \wedge \dots \wedge du^{2n-1},$$

where $\Omega = \det(B_\alpha^A, C^A) \det(g_{ij})$.

Proof. This is well known, see [Tah69], and can also be checked by a direct computation. \square

5.4. The Riemannian metric on STM . The Riemannian metric g on M can be extended to a Riemannian metric \bar{g} on the sphere bundle STM . Explicitly, \bar{g} is given by (cf. [Tah69]):

$$\begin{aligned} \bar{g}_{ji} &= g_{ji} + g_{ts} (\nabla_j y^t) (\nabla_i y^s), \\ \bar{g}_{\mu i} &= g_{ts} (\partial_\mu y^t) (\nabla_i y^s), \\ \bar{g}_{\mu \lambda} &= g_{ji} (\partial_\mu y^j) (\partial_\lambda y^i). \end{aligned}$$

³Following [Tah69], we will adopt the following index gymnastics. Capital Latin letters such as A, B, \dots run from 1 to $2n$. Small latin letters such as i, j, \dots run from 1 to n . Barred Latin indices such as \bar{i}, \bar{j}, \dots run from $n+1$ to $2n$. Some of the Greek letters such as α, β, \dots run from 1 to $2n-1$. Some other Greek letters such as κ, λ, \dots run from $n+1$ to $2n-1$.

The inverse of \bar{g} is given by

$$\begin{aligned}\bar{g}^{ji} &= g^{ji}, \\ \bar{g}^{\lambda h} &= -g^{hl}(\nabla_l y^i)B^\lambda_{\bar{i}}, \\ \bar{g}^{\lambda\kappa} &= (g^{ih} + g^{ts}(\nabla_t y^i)(\nabla_s y^h))B^\lambda_{\bar{i}}B^\kappa_{\bar{h}}.\end{aligned}$$

The Christoffel symbols associated with this metric are given by

$$\begin{aligned}\bar{\Gamma}_{ji}^h &= \Gamma_{ij}^h + \frac{1}{2}(R_{rsj}{}^h y^r \nabla_i y^s + R_{rsi}{}^h y^r \nabla_j y^s), \\ \bar{\Gamma}_{\mu i}^h &= \frac{1}{2}R_{rsi}{}^h y^r B_\mu^{\bar{s}}, \\ \bar{\Gamma}_{\mu\lambda}^h &= 0, \\ \bar{\Gamma}_{ji}^\kappa &= (\nabla_j \nabla_i y^h + \frac{1}{2}R_{rji}{}^h y^r - \frac{1}{2}R_{rij}{}^h y^r - \frac{1}{2}(R_{rsj}{}^l y^r \nabla_i y^s + \frac{1}{2}R_{rsi}{}^l y^r \nabla_j y^s) \nabla_l y^h) B^\kappa_{\bar{h}}, \\ \bar{\Gamma}_{\mu j}^\kappa &= (\partial_\mu \nabla_i y^h - \frac{1}{2}R_{rsi}{}^l y^r B_\mu^{\bar{s}} \nabla_l y^h) B^\kappa_{\bar{h}}, \\ \bar{\Gamma}_{\mu\lambda}^\kappa &= (\partial_\mu \partial_\lambda y^h) B^\kappa_{\bar{h}}.\end{aligned}$$

5.5. Proof of Theorem 5.2.1. We are ready to prove the main result of this section.

Lemma 5.5.1. *We have*

$$(16) \quad \begin{aligned}y^i \partial_i(\Omega) &= -y^i B_\lambda^{\bar{h}} \partial_i(B^\lambda_{\bar{h}}) \Omega + 2y^i \Gamma_i \Omega - y^i y^h y^m g_{hm} \Gamma_{ir}^h \Omega, \\ y^l (\nabla_l y^i) B^\lambda_{\bar{i}} \partial_\lambda(\Omega) &= -y^l (\nabla_l y^i) \partial_\lambda(B^\lambda_{\bar{i}}) \Omega, \\ y^l (\nabla_l y^k) (\partial_\mu \partial_\lambda y^j) B^\lambda_{\bar{k}} B^\mu_{\bar{j}} &= -y^l (\nabla_l y^k) \partial_\lambda(B^\lambda_{\bar{k}}), \\ \partial_\lambda (\nabla_l y^h) B^\lambda_{\bar{h}} y^l &= -(\partial_l B^\lambda_{\bar{h}}) B_\lambda^{\bar{h}} y^l + \Gamma_{li}^i y^l - \Gamma_{rs}^h g_{ih} y^i y^r y^s.\end{aligned}$$

Proof. The first and the second lines of (16) follow from the fact that

$$\begin{aligned}B^\lambda_{\bar{k}} \partial_\mu (B_\lambda^{\bar{j}}) &= -\partial_\mu (B^\lambda_{\bar{k}}) B_\lambda^{\bar{j}} - \partial_\mu (y^j y^m) g_{mk}, \\ y^i \partial_i B_\lambda^{\bar{j}} &= -y^i B_{\lambda}^{\bar{h}} B_\kappa^{\bar{j}} \partial_i (B^\kappa_{\bar{h}}) - y^i y^j y^m g_{hm} \Gamma_{il}^h B_\lambda^{\bar{l}},\end{aligned}$$

and the property of the determinant. The third line of (16) follows from the fact that

$$B_\lambda^{\bar{h}} B^\lambda_{\bar{i}} + y^h C_{\bar{i}}^h = \delta_i^h$$

and applying to it the partial derivative ∂_μ . The fourth line of (16) follows when we substitute the covariant derivative $\nabla_l y^h = \partial_l y^h + \Gamma_{li}^h y^i$ and use the third equation. \square

By definition,

$$A_{\bar{g}, \Theta} = \bar{g}^{ih} \bar{\nabla}_i (\theta_h \Omega^{-\frac{1}{n}}) + \bar{g}^{\lambda h} \bar{\nabla}_\lambda (\theta_h \Omega^{-\frac{1}{n}}) + \bar{g}^{h\lambda} \bar{\nabla}_h (\theta_\lambda \Omega^{-\frac{1}{n}}) + \bar{g}^{\lambda\kappa} \bar{\nabla}_\lambda (\theta_\kappa \Omega^{-\frac{1}{n}}).$$

The last two summands vanish because $\theta_\kappa = 0$. Let us compute the first two summands separately. Applying the covariant derivative $\bar{\nabla}$, we get

$$\begin{aligned}\bar{g}^{ih} \bar{\nabla}_i (\theta_h \Omega^{-\frac{1}{n}}) &= \partial_i y^i \Omega^{-\frac{1}{n}} + y^i \partial_i (\Omega^{-\frac{1}{n}}) - R_{qrsi} y^q y^r (\nabla_h y^s) g^{ih} \Omega^{-\frac{1}{n}} \\ &\quad + (1 + \frac{1}{n}) \Gamma_{ij}^j y^i \Omega^{-\frac{1}{n}} + \frac{1}{n} (\frac{1}{2} R_{rsih} y^r y^l (\nabla_l y^s) + \partial_\lambda (\nabla_l y^h) B^\lambda_{\bar{h}} y^l) \Omega^{-\frac{1}{n}}.\end{aligned}$$

Similarly,

$$\begin{aligned}\bar{g}^{\lambda h} \bar{\nabla}_\lambda (\theta_h \Omega^{-\frac{1}{n}}) &= -\partial_i y^i \Omega^{-\frac{1}{n}} - \Gamma_{ij}^j y^i \Omega^{-\frac{1}{n}} - (\nabla_i y^k) B^\lambda_{\bar{k}} y^i \partial_\lambda (\Omega^{-\frac{1}{n}}) \\ &\quad + \frac{1}{2} R_{rskh} y^r y^s g^{hl} (\nabla_l y^k) \Omega^{-\frac{1}{n}} - \frac{1}{2n} (\nabla_i y^k) y^i R_{rkh}{}^{\bar{h}} y^r \Omega^{-\frac{1}{n}} \\ &\quad + -\frac{1}{n} y^l (\nabla_l y^k) (\partial_\lambda \partial_\mu y^j) B^\lambda_{\bar{k}} B^\mu_{\bar{j}} \Omega^{-\frac{1}{n}}.\end{aligned}$$

By collecting the terms and using Lemma 5.5.1, we finally obtain:

$$A_{\tilde{g},\Theta} = -\frac{1}{2} (R_{ilsj} y^i y^l (\nabla_h y^s) g^{jh}) \text{vol}^{-\frac{1}{n}} \equiv 0,$$

since the curvature tensor R_{ilsj} is antisymmetric in two first indices.

Theorem 5.2.1 is proved.

6. EXAMPLES

We finish the paper with concrete examples of Riemannian curl for the 3-dimensional sphere (with two natural metrics) and the 3-dimensional ellipsoid with the standard metric.

6.1. The sphere S^3 . Consider the sphere S^3 in the standard symplectic space \mathbb{R}^4 . It is endowed with the natural contact structure that can be defined by the contact form

$$\theta = dz + xdy - ydx,$$

where x, y and z are affine coordinates on S^3 . More precisely, if p_1, p_2, q^1, q^2 are symplectic Darboux coordinates on \mathbb{R}^4 , then $x = \frac{p_1}{q^2}$, $y = \frac{q^1}{q^2}$, $z = -\frac{p_2}{q^2}$.

The restriction of the Euclidean metric to the sphere S^3 takes the following form:

$$\begin{aligned} g_{S^3} = & F \left((y^2 + z^2 + 1) dx^2 + (x^2 + z^2 + 1) dy^2 + (x^2 + y^2 + 1) dz^2 \right. \\ & \left. - 2xy dx dy - 2xz dx dz - 2yz dy dz \right), \end{aligned}$$

where $F = \frac{1}{(x^2 + y^2 + z^2 + 1)^2}$.

Let us also consider another, conformally equivalent, metric on S^3 :

$$\tilde{g}_{S^3} := \left(\frac{x^2 + y^2 + z^2 + 1}{\frac{x^2}{a} + \frac{y^2}{b} + \frac{z^2}{c} + 1} \right) g_{S^3},$$

which appeared in the context of integrable systems in [Tab99, MT01], see also [DV11].

Proposition 6.1.1. *The following results hold.*

- (i) *In the case of the “round” metric g_{S^3} , we have $A_{g_{S^3}} = 0$;*
- (ii) *In the case of the metric \tilde{g}_{S^3} , we have $A_{\tilde{g}_{S^3}} = \frac{5}{2} \left(\left(\frac{1}{b} - \frac{1}{a} \right) xy + \left(\frac{1}{c} - 1 \right) z \right)$.*

Proof. Part (i) follows from Corollary 3.4.1. Part (ii) can be obtained by a straightforward computation using Eq. (13). \square

6.2. The case of the ellipsoid $E^3(a, b, c)$. Consider the 3-dimensional ellipsoid endowed with the standard metric

$$g_{E^3_{a,b,c}} = g_{x,x} dx^2 + g_{y,y} dy^2 + g_{z,z} dz^2 + g_{x,y} dx dy + g_{x,z} dx dz + g_{y,z} dy dz,$$

where

$$\begin{aligned}
g_{x,x} &= \frac{(b^2y^2 + c^2z^2 + 1)^2 + a^4x^2(y^2 + z^2 + 1)}{((ax)^2 + (by)^2 + (cz)^2 + 1)^2}, \\
g_{y,y} &= \frac{(a^2x^2 + c^2z^2 + 1)^2 + b^4y^2(x^2 + z^2 + 1)}{((ax)^2 + (by)^2 + (cz)^2 + 1)^2}, \\
g_{z,z} &= \frac{(a^2x^2 + b^2y^2 + 1)^2 + c^4z^2(x^2 + y^2 + 1)}{((ax)^2 + (by)^2 + (cz)^2 + 1)^2}, \\
g_{x,y} &= -2xy \frac{a^4x^2 - a^2(z^2(b^2 - c^2) + b^2 - 1) + b^2(b^2y^2 + c^2z^2 + 1)}{((ax)^2 + (by)^2 + (cz)^2 + 1)^2}, \\
g_{x,z} &= -2xz \frac{a^4x^2 - a^2(y^2(c^2 - b^2) + c^2 - 1) + c^2(b^2y^2 + c^2z^2 + 1)}{((ax)^2 + (by)^2 + (cz)^2 + 1)^2}, \\
g_{y,z} &= -2yz \frac{b^4y^2 - b^2(x^2(c^2 - a^2) + c^2 - 1) + c^2(a^2x^2 + c^2z^2 + 1)}{((ax)^2 + (by)^2 + (cz)^2 + 1)^2}.
\end{aligned}$$

Proposition 6.2.1. *We have*

$$\begin{aligned}
A_{g_{E^3_{a,b,c}}} &= a^4(a^2 - b^2)(b^2 + 2c^2 + 2)x^3y + b^4(a^2 - b^2)(a^2 + 2c^2 + 2)xy^3 \\
&\quad + (a^2 - b^2)c^4(2 + a^2 + b^2 + c^2)xyz^2 \\
&\quad - a^4(c^2 - 1)(a^2 + 2b^2 + c^2 + 1)x^2z - b^4(c^2 - 1)(2a^2 + b^2 + c^2 + 1)y^2z \\
&\quad - c^4(c^2 - 1)(2a^2 + 2b^2 + 1)z^3 + (a^2 - b^2)(a^2 + b^2 + 2c^2 + 1)xy \\
&\quad - (c^2 - 1)(2a^2 + 2b^2 + c^2)z.
\end{aligned}$$

Proof. Straightforward computation using Eq. (13). \square

Acknowledgments. We thank Dimitry Leites and Christian Duval their interest in this work and careful reading of preliminary versions of it. We are also grateful to Charles Conley, Eugene Ferapontov and Serge Tabachnikov for a number of fruitful discussions. The first author was partially supported by the Grant NYUAD 063. The second author was partially supported by the Grant PICS05974 “PENTAFRIZ”, of CNRS.

REFERENCES

- [Arn89] V.I. ARNOLD, *Mathematical methods of classical mechanics*. Second edition. Graduate Texts in Mathematics, **60**, Springer-Verlag, New York, 1989.
- [Bla10] D. BLAIR, *Riemannian geometry of contact and symplectic manifolds*. Second edition. Progress in Mathematics, 203. Birkh user Boston, Inc., Boston, MA, 2010.
- [BKM09] A. BOLSINOV, V. KIOSAK, V. MATVEEV, A Fubini theorem for pseudo-Riemannian geodesically equivalent metrics. *J. Lond. Math. Soc. (2)* **80** (2009), 341–356.
- [B06] S. BOUARROUDJ, Projective and conformal Schwarzian derivatives and cohomology of Lie algebras vector of fields related to differential operators, *Int. Jour. Geom. Methods. Mod. Phys.* **3** (2006), 667–696.
- [BGLS12] S. BOUARROUDJ, P. GROZMAN, D. LEITES, I. SHCHEPOCHKINA,, Minkowski superspaces and superstrings as almost real-complex supermanifolds, *Theor. and Mathem. Physics*, **173** (2012), 1687–1708; [arXiv:1010.4480](https://arxiv.org/abs/1010.4480).
- [BO00] S. BOUARROUDJ, V. OVSIENKO, Schwarzian derivative related to modules of differential operators on a locally projective manifold. *Banach Cent. Publ.*, **51**, 2000.
- [Car24] E. CARTAN, Sur les vari t s   connexion projective. *Bull. Soc. Math. Fr.* **52** (1924), 205–241.
- [CO12] C. CONLEY, V. OVSIENKO, Linear differential operators on contact manifolds, [arXiv:1205.6562](https://arxiv.org/abs/1205.6562).

- [DNF92] B.A. DUBROVIN, A.T. FOMENKO, S.P. NOVIKOV, Modern geometry methods and applications. Part I. The geometry of surfaces, transformation groups, and fields. Second edition. Graduate Texts in Mathematics, **93**, Springer-Verlag, New York, 1992.
- [DO97] C. DUVAL, V. OVSIENKO, Space of second-order linear differential operators as a module over the Lie algebra of vector fields. *Adv. Math.* **132** (1997), no. 2, 316–333.
- [DO01] C. DUVAL, V. OVSIENKO, Conformally equivariant quantum Hamiltonians. *Selecta Math.* (N.S.) **7** (2001), no. 3, 291–320.
- [DV11] C. DUVAL, G. VALENT, A new integrable system on the sphere and conformally equivariant quantization. *J. Geom. Phys.* **61** (2011), no. 8, 1329–1347.
- [Eis97] L.P. EISENHART, Riemannian Geometry. Eighth edition. Princeton University Press, Princeton, N.J. 1997.
- [Fu86] D.B. FUKS, Cohomology of Infinite Dimensional Lie Algebras, Plenum, New York, 1986.
- [Gel70] I.M. GELFAND, The cohomology of infinite-dimensional Lie algebras: some questions of integral geometry, *Proc. Internat. Cong. Math.* (Nice 1970), Gauthier-Villars, Paris 1971, Vol. 1, pp. 95–111.
- [GL07] P. GROZMAN, D. LEITES, Nonholonomic Riemann and Weyl tensors for flag manifolds *Theoret. Math. Phys.* **153** (2007), 1511–1538.
- [GLS01] P. GROZMAN, D. LEITES, I. SHCHEPOCHKINA, Lie superalgebras of string theories, *Acta Math. Vietnam.* **26** (2001), 27–63.
- [GLS02] P. GROZMAN, D. LEITES, I. SHCHEPOCHKINA, Invariant operators on supermanifolds and standard models. Multiple facets of quantization and supersymmetry, 508–555, World Sci. Publ., River Edge, NJ, 2002; arXiv:math/0202193.
- [KN64] S. KOBAYASHI, T. NAGANO, On projective connections. *J. Math. Mech.* **13** (1964), 215–235.
- [Leb10] S. A. LEBEDEV, Analogs of the orthogonal, Hamiltonian, Poisson, and contact Lie superalgebras in characteristic 2. *J. Nonlinear Math. Phys.*, **17**, Special issue in memory of F. Berezin, (2010), 217–251.
- [MT01] V. MATVEEV, P. TOPALOV, Quantum integrability of Beltrami-Laplace operator as geodesic equivalence, *Math. Z.* **238** (2001), 833–866.
- [Ovs90] V. OVSIENKO, Contact analogues of the Virasoro algebra *Funct. Anal. Appl.* **24** (1990), 306–314.
- [Ovs06] V. OVSIENKO, Vector fields in the presence of a contact structure, *Enseign. Math.* (2) **52** (2006), no. 3-4, 215–229.
- [OT05] V. OVSIENKO, S. TABACHNIKOV, Projective differential geometry old and new. From the Schwarzian derivative to the cohomology of diffeomorphism groups. Cambridge Tracts in Mathematics, 165. Cambridge University Press, Cambridge, 2005.
- [Tab99] S. TABACHNIKOV, Projectively equivalent metrics, exact transverse line fields and the geodesic spray on the ellipsoid, *Comment. Math. Helv.* **74** (1999), 306–321.
- [Tah69] Y. TASHIRO, On contact structures of tangent sphere bundles, *Tôhoku Math. Journal.* **21** (1969), 117–143.
- [vE10] E. VAN ERP, The Atiyah-Singer index formula for subelliptic operators on contact manifolds, Part I, *Ann. Math.* (2) **171** (2010), no. 3, 1647–1681.
- [Veb22] O. VEULEN, Projective and affine geometry of paths, *Proc. Nat. Acad. Sci. USA* **8** (1922), 347–352.
- [Yan52] K. YANO, Some remarks on tensor fields and curvature. *Ann. of Math.* (2) **55**, (1952), 328–347.

DIVISION OF SCIENCE AND MATHEMATICS, NEW YORK UNIVERSITY ABU DHABI, PO BOX 129188, UNITED ARAB EMIRATES

E-mail address: sofiane.bouarroudj@nyu.edu

CNRS, INSTITUT CAMILLE JORDAN, UNIVERSITÉ CLAUDE BERNARD LYON I, 21 AVENUE CLAUDE BERNARD, 69622 VILLEURBANNE CEDEX, FRANCE

E-mail address: ovsienko@math.univ-lyon1.fr