

CAPACITY OF THE RANGE OF RANDOM WALK ON $\mathbb{Z}^4$

BY AMINE ASSELAH, BRUNO SCHAPIRA AND PERLA SOUSI

Université Paris-Est, Aix-Marseille Université and University of Cambridge

We study the scaling limit of the capacity of the range of a random walk on the integer lattice in dimension four. We establish a strong law of large numbers and a central limit theorem with a non-Gaussian limit. The asymptotic behaviour is analogous to that found by Le Gall in '86 [*Comm. Math. Phys.* **104** (1986) 471–507] for the *volume* of the range in dimension two.

CONTENTS

1. Introduction	1447
2. Preliminaries	1453
2.1. Notation and standard estimates	1453
2.2. A decomposition formula for the capacity	1456
3. Strong law of large numbers	1457
3.1. Preliminaries	1458
3.2. First and second moment estimates of the cross terms	1459
3.3. Strong law of large numbers	1460
4. Existence and definition of the limiting term	1462
4.1. Carleman's condition	1462
4.2. The limiting term	1465
5. Intersection and nonintersection probabilities	1466
6. Joint convergence in law of the cross terms	1473
6.1. Preliminaries	1474
6.2. Localisation of one cross term	1476
6.3. Moments of χ	1482
6.4. Proof of Proposition 6.1	1489
7. Central limit theorem	1491
Appendix	1492
Acknowledgements	1495
References	1495

1. Introduction. This paper is devoted to the study of the capacity of the range of a simple random walk in dimension four. The point of view we adopt is that the capacity is a *hitting probability*. More precisely, the capacity of a set is proportional to the probability a random walk *sent from infinity* hits the set. Then the capacity of the range of a random walk is cast into a problem of *intersection*

Received January 2017; revised March 2018.

MSC2010 subject classifications. Primary 60F05; secondary 60G50.

Key words and phrases. Capacity, Green kernel, law of large numbers, central limit theorem.

of paths, and dimension four is *critical* in view of classical results of Dvoretzky, Erdős and Kakutani [14] establishing that the paths of two independent Brownian motions do not intersect if, and only if, dimension is four or larger.

The capacity of a set $A \in \mathbb{Z}^4$ can also be viewed as an escape probability. Indeed, let $\mathbb{P}_x$ be the law of a simple random walk starting at x , let G_d be the discrete Green's function and let H_A and H_A^+ stand respectively for the hitting time of a finite set A and the return time in A . Then

$$(1.1) \quad \text{Cap}(A) = \sum_{x \in A} \mathbb{P}_x(H_A^+ = \infty) = \lim_{\|y\| \rightarrow \infty} \frac{\mathbb{P}_y(H_A < \infty)}{G_d(0, y)}.$$

One easily passes from one representation in (1.1) to the other using the last passage decomposition formula; see (2.8) below.

Denote by $\{S(n), n \in \mathbb{N}\}$ a simple random walk in $\mathbb{Z}^4$. For two integers m, n , the range $\mathcal{R}[m, n]$ (or simply $\mathcal{R}_n$ when $m = 0$) in the time period $[m, n]$ is defined as

$$\mathcal{R}[m, n] = \{S(m), \dots, S(n)\}.$$

Our first result is a strong law of large numbers for $\text{Cap}(\mathcal{R}_n)$.

THEOREM 1.1. *Let S be a simple random walk in $\mathbb{Z}^4$. Almost surely,*

$$\lim_{n \rightarrow \infty} \frac{\log n}{n} \cdot \text{Cap}(\mathcal{R}_n) = \frac{\pi^2}{8}.$$

Our second result is a central limit theorem for $\text{Cap}(\mathcal{R}_n)$, and requires more notation: G denotes the continuous Green's function, $\xrightarrow{(d)}$ convergence in law and $(\beta_s, s \geq 0)$ a standard 4-dimensional Brownian motion.

THEOREM 1.2. *Let S be a simple random walk in $\mathbb{Z}^4$. Then, as n goes to infinity*

$$\frac{(\log n)^2}{n} \cdot (\text{Cap}(\mathcal{R}_n) - \mathbb{E}[\text{Cap}(\mathcal{R}_n)]) \xrightarrow{(d)} -\frac{\pi^4}{4} \cdot \gamma_G([0, 1]^2),$$

where $\gamma_G([0, 1]^2)$ is formally defined as

$$(1.2) \quad \gamma_G([0, 1]^2) = \int_0^1 \int_0^1 G(\beta_s, \beta_t) ds dt - \mathbb{E} \left[\int_0^1 \int_0^1 G(\beta_s, \beta_t) ds dt \right].$$

Moreover, $\gamma_G([0, 1]^2)$ is nondegenerate, and non-Gaussian, since for some $\lambda \in \mathbb{R}$,

$$(1.3) \quad \mathbb{E}[\exp(\lambda \gamma_G([0, 1]^2))] = \infty.$$

REMARK 1.3. Although both terms appearing in the definition (1.2) of $\gamma_G([0, 1]^2)$ have infinite expectation, we make sense, in Section 4, of $\gamma_G([0, 1]^2)$ as an L^2 -random variable following Le Gall's approach used to define the self-intersection local time; see [27–29]. We also prove there that it has some infinite exponential moment.

REMARK 1.4. Theorem 1.2 shows that the capacity of the range in dimension 4 behaves in the same way as the size of the range in dimension 2 as shown by Le Gall in '86 [28]:

$$\frac{(\log n)^2}{n} \cdot (|\mathcal{R}_n| - \mathbb{E}[|\mathcal{R}_n|]) \xrightarrow{(d)} -\pi^2 \cdot \gamma([0, 1]^2),$$

where $\gamma([0, 1]^2)$ is defined formally via

$$\gamma([0, 1]^2) = \int_0^1 \int_0^1 \delta_{(0)}(\beta_s - \beta_t) ds dt - \mathbb{E} \left[\int_0^1 \int_0^1 \delta_{(0)}(\beta_s - \beta_t) ds dt \right],$$

where β is a standard 2-dimensional Brownian motion.

As a corollary of our result, we obtain the asymptotic behaviour of the variance of $\text{Cap}(\mathcal{R}_n)$.

COROLLARY 1.5. *Let S be a simple random walk in $\mathbb{Z}^4$. Then*

$$\lim_{n \rightarrow \infty} \frac{(\log n)^4}{n^2} \text{Var}(\text{Cap}(\mathcal{R}_n)) = \frac{\pi^8}{16} \cdot \mathbb{E}[(\gamma_G([0, 1]^2))^2].$$

We note that the limiting term in this corollary is nonzero.

Seen as a normalised hitting probability, the capacity of the range of a walk is an object which appeared in disguised form in topics linked with *intersection of paths of random walks*. This latter topic grew already large in the nineties, as Lawler's '91 book [24] testifies. One reason for that is the many diverse sources of motivation: (i) quantum field theories with a seminal insight of Symanzik [35], and with contributions by Lawler [22, 23], Aizenman [1], Felder and Fröhlich [16], to name a few (see the book [17] for a historical account and references therein), (ii) probability and the non-Markovian model of *self-avoiding walk*, with contributions from Brydges and Spencer [6], Madras and Slade [32] and Lawler [21], (iii) conformal field theories, and the intersection exponents relations in dimension two and with contributions by Duplantier and Kwon [11], Duplantier [10], Burdzy and Lawler [7], and Lawler, Schramm and Werner [26] and references therein.

The models studied can be either discrete random walks, or their continuous counterpart, the Wiener sausages. In the mid-nineties, Aizenman [1], Albeverio and Zhou [2], Pemantle, Peres and Shapiro [33] and Khoshnevisan [20] proposed useful methods and estimates for the intersection of two Wiener sausages. These estimates were important in understanding *how small* was the trace of a Brownian motion. In 2004, van den Berg, Bolthausen and den Hollander [37] studied the upward deviations for the volume of intersection of two Wiener sausages, and established a large deviations principle. More recently, Erhard and Poisat [15] also established large deviations estimates for the capacity of a Wiener sausage.

Recently, there has been a revival of problems linked with *intersection of paths*. The model of random interlacements was invented by Sznitman [36] initially to

study the trace of a walk, living in d -dimensional torus of side N for a time N^d . Sznitman introduced a measure on infinite paths on the infinite lattice of random walks whose probability of avoiding any given set is proportional to the exponential of minus its Newtonian capacity. Recently, Rath and Sapozhnikov [34] and Chang and Sapozhnikov [9] established moments and deviation bounds for the capacity of the union of ranges of paths. In [4], we observed that the precise two-sided nonintersection bounds of Lawler [24] yield in dimension four,

$$(1.4) \quad \lim_{n \rightarrow \infty} \frac{\log n}{n} \mathbb{E}[\text{Cap}(\mathcal{R}_n)] = \frac{\pi^2}{8}.$$

Soon after this, Chang [8] obtained a sharp estimate on the second moment:

$$\lim_{n \rightarrow \infty} \frac{\mathbb{E}[\text{Cap}(\mathcal{R}_n)^2]}{\mathbb{E}[\text{Cap}(\mathcal{R}_n)]^2} = 1,$$

implying a weak law of large numbers. Chang [8] also established a fluctuation result in dimension three by coupling the walk and the Brownian motion:

$$(1.5) \quad \frac{\text{Cap}(\mathcal{R}_n)}{\sigma \sqrt{n}} \xrightarrow{(d)} \text{Cap}(\beta[0, 1]),$$

with σ some renormalising constant and $\beta[0, 1]$ the trace of a 3-dimensional Brownian motion between time 0 and 1. In [4], we also proved a standard central limit theorem in dimension larger than or equal to 6 (with a standard $\sqrt{n}$ normalising factor, and a Gaussian limit), while the law of large numbers had already been obtained in dimension 5 and larger by Jain and Orey [18], almost 50 years ago. A striking correspondence emerges: all these results for the capacity of the range are analogous to results for the volume of the range (see [13, 19, 28]), but only after dropping space dimension by two units to go from capacity to volume of the range. The remaining open issue is the central limit theorem for the capacity of the range in dimension 5.

Recently, van den Berg, Bolthausen and den Hollander [38] studied the torsional rigidity of the complement of a Wiener sausage, as a way to probe the shape of the sausage. In order to obtain leading asymptotics for the torsional rigidity, one needs a law of large numbers for the capacity of a Wiener sausage, which is now proved in dimension four in our companion paper [5]. Our Theorem 1.1 establishes these asymptotics for the discrete model, and thus prepares the study of torsional rigidity for random walk.

Our own motivation for studying the capacity of the range of a random walk comes from studying a random walk conditioned on being localised during a time-period $[0, N]$ in a ball of volume of order N [3]. In this regime, the localised walk necessarily intersects often its own path, and one of the main technical estimates in [3] concerns the event of visiting a set Λ made up of nonoverlapping balls of fixed radius. We establish that visiting each ball, making up Λ , the same number of times is related to the capacity of Λ . This allowed us to obtain rough estimates

on the capacity of the range of a localised walk, and convinced us that the capacity of the range was a relevant object to consider.

Heuristics. We wish now to explain at a heuristic level the scaling of the capacity in $d = 4$, as well as the reason for our CLT. Along the way, we present a simple decomposition formula for the capacity of two finite sets, and highlight the connection with the volume of the range in $d = 2$.

Let us start by explaining in simple terms why the scaling of the capacity of the range is $n/\log n$ in dimension 4. Consider (1.1) where $A = \mathcal{R}_n$, and observe that it is enough to consider the site x on the boundary of a ball, say of radius R , containing the set A : a walker coming from infinity basically spreads uniformly on the boundary of such a ball when it hits it, and (1.1) is almost correct when considering x uniformly distributed on the boundary of ball of radius R and normalised by $G_d(0, R)$, since $G_d(0, y)/G_d(0, R)$ is the probability of eventually hitting the ball of radius R when starting at y . Now, during a time-period $[0, n]$, the walk typically stays in a ball of radius $R = \sqrt{n}$, and we consider R of this order. *We need therefore to estimate the probability that two independent walks starting at a distance $\sqrt{n}$ meet.* More precisely, we need to estimate

$$\mathbb{P}_{0,x}(\mathcal{R}[0, n] \cap \tilde{\mathcal{R}}[0, \infty) \neq \emptyset) \quad \text{with } \|x\| \sim \sqrt{n}.$$

To estimate this intersection event, Lawler [24] counts the number of times the two paths intersect. Its expectation is expressed as a product of the probability the two walks meet times the mean number of meetings after the first one, that is when they start from the same point. Then simple computations give the following when $\|x\| \sim \sqrt{n}$:

$$\mathbb{E}_{0,0}[|\mathcal{R}[0, n] \cap \tilde{\mathcal{R}}[0, \infty)|] \asymp \log n \quad \text{and} \quad \mathbb{E}_{0,x}[|\mathcal{R}[0, n] \cap \tilde{\mathcal{R}}[0, \infty)|] \asymp 1.$$

Then the order of the probability of intersection is obtained by taking the ratio of the two previous quantities, and dividing by $G_d(0, R)$ which is of order $1/R^2$. This is established rigorously in Section 4.3 of Lawler [24]. The scaling for the capacity follows, at least heuristically.

A simple and key observation of Le Gall [28], using the symmetry of the increments and translation invariance of the lattice, is that the range $\mathcal{R}[0, 2n]$ translated by $S(n)$, is the union of two independent ranges: $\mathcal{R}_n^1 = \mathcal{R}[0, n] - S(n)$ and $\mathcal{R}_n^2 = \mathcal{R}[n, 2n] - S(n)$, which yields by the exclusion-inclusion formula

$$(1.6) \quad |\mathcal{R}_n^1 \cup \mathcal{R}_n^2| = |\mathcal{R}_n^1| + |\mathcal{R}_n^2| - |\mathcal{R}_n^1 \cap \mathcal{R}_n^2|.$$

This is the starting block of Le Gall's proof. Our starting point is that the capacity of the range obeys a decomposition formula which plays exactly the same role as the exclusion-inclusion formula does for the volume of the range.

PROPOSITION 1.6. *Let A and B be two finite subsets of $\mathbb{Z}^d$. We have*

$$(1.7) \quad \text{Cap}(A \cup B) = \text{Cap}(A) + \text{Cap}(B) - \chi(A, B) - \chi(B, A) + \varepsilon(A, B),$$

where

$$\chi(A, B) = \sum_{y \in A} \sum_{z \in B} \mathbb{P}_y(H_{A \cup B}^+ = \infty) G_d(y, z) \mathbb{P}_z(H_B^+ = \infty),$$

and $0 \leq \varepsilon(A, B) \leq \text{Cap}(A \cap B)$.

REMARK 1.7. The equalities (1.6) and (1.7) explain the striking correspondence between asymptotics for the volume and the capacity of the range. As long as the term ε in (1.7) is innocuous, the order of magnitude of the cross term $\chi(\mathcal{R}_n^1, \mathcal{R}_n^2)$ in dimension $d + 2$ is shown to be the same as the order of magnitude of the intersection term $|\mathcal{R}_n^1 \cap \mathcal{R}_n^2|$ in dimension d .

REMARK 1.8. We intend to apply (1.7) with $A = \mathcal{R}_n^1$ and $B = \mathcal{R}_n^2$. In dimension 4, it turns out that $\varepsilon(A, B)$ is innocuous. One classical inequality $\text{Cap}(A \cup B) \leq \text{Cap}(A) + \text{Cap}(B) - \text{Cap}(A \cap B)$ (see Proposition 2.3.4 of Lawler [24]) misses the χ terms in (1.7) which dominate the fluctuations.

As in the CLT proof for the volume of the range in dimension 2 [28], we iterate (1.7) and write the capacity of the range as the sum of a rescaled self-similar part, consisting of a sum of independent and (almost) identically distributed terms, plus a sum of cross terms. Our proofs establish that for the law of large numbers it is the self-similar part which dominates, the cross terms being of smaller order than $n / \log n$, while for the central limit theorem, it is the opposite situation: the fluctuations of the self-similar part are negligible compared to those of the cross terms of order $n / \log^2 n$. This striking phenomenon is exactly the same as the one Le Gall discovered some 30 years ago, when dealing with the volume of the range [28].

We are now in a position to shed some light on the form of our CLT. We consider $\chi(\mathcal{R}_n^1, \mathcal{R}_n^2)$ and expect, as Theorem 1.1 essentially teaches, that typically and to leading order for $x \in \mathcal{R}_n^1$ and $y \in \mathcal{R}_n^2$

$$\mathbb{P}_x(H_{\mathcal{R}_n^1 \cup \mathcal{R}_n^2}^+ = \infty) \sim \frac{\pi^2}{8} \cdot \frac{1}{\log n} \quad \text{and} \quad \mathbb{P}_y(H_{\mathcal{R}_n^2}^+ = \infty) \sim \frac{\pi^2}{8} \cdot \frac{1}{\log n}.$$

Note that in dimension higher than four, typically for $x \in \mathcal{R}[0, n]$, the probability $\mathbb{P}_x(H_{\mathcal{R}[0, n]}^+ = \infty)$ is rather of order 1. Our key technical estimates is then to make the escape events into local events (in a space scale much smaller than $\sqrt{n}$), and thus transform the intersection term in a term looking to leading order like

$$(1.8) \quad \left(\frac{\pi^2}{8} \cdot \frac{1}{\log n} \right)^2 \cdot \sum_{x \in \mathcal{R}_n^1} \sum_{y \in \mathcal{R}_n^2} G_d(x, y).$$

The expression (1.8), in conjunction with the decomposition (1.7) which we iterate, explains heuristically the form (1.2).

The rest of the paper is organised as follows. In Section 2, we start by recalling known estimates on Green's kernel, and deriving useful simple estimates on random walks. Then we present the proof of Proposition 1.6. The strong law of large numbers is established in Section 3 after a rough second estimate is obtained for the cross term. Section 4 studies the limiting object in the CLT. Section 5 presents our nonintersection events—Proposition 5.2 which generalises Lawler's Theorem 5.1. Section 6 presents the asymptotics for the cross term obtained by the method of moments. Section 7 establishes the CLT based on estimates of Section 6, and the recursive use of the decomposition. Finally, the Appendix gathers computations linked with Section 5.

2. Preliminaries.

2.1. Notation and standard estimates. We mostly use the symbol S to denote a random walk, and use both notation S_k and $S(k)$ to denote its position at time k . When $0 \leq a \leq b$ are real numbers, $\mathcal{R}[a, b]$ denotes $\mathcal{R}([a], [b])$, where $[x]$ stands for the integer part of x . By convention, if $a > b$, then $\mathcal{R}[a, b]$ is the empty set. We also write $\mathcal{R}_a$ for $\mathcal{R}[0, [a]]$, and $S(n/2)$ for $S([n/2])$.

For positive functions f, g we write $f \sim g$ if $f(n)/g(n) \rightarrow 1$ as $n \rightarrow \infty$. We also write $f(n) \lesssim g(n)$ if there exists a constant $c > 0$ such that $f(n) \leq cg(n)$ for all n , and $f(n) \gtrsim g(n)$ if $g(n) \lesssim f(n)$. Finally, we use the notation $f(n) \asymp g(n)$ if both $f(n) \lesssim g(n)$ and $f(n) \gtrsim g(n)$.

For $\alpha > 0$, and $n \geq 2$, we note $n_\alpha := n \cdot (\log n)^{-\alpha}$.

The Euclidean norm of $x \in \mathbb{Z}^4$ is denoted $\|x\|$, and the Euclidean ball of center x and radius r is denoted $\mathcal{B}(x, r)$. We denote by $\mathbb{P}_x$ the law of a simple random walk starting from x , and simply write $\mathbb{P}$ when $x = 0$. Likewise $\mathbb{P}_{x, x'}$ denotes the law of two independent random walks starting from x and x' , and similarly when there are more walks. Recall that H_A denotes the hitting time of a set A , and we abbreviate this in H_x when A is reduced to a single point $x \in \mathbb{Z}^d$.

We write

$$p_k(x, y) = \mathbb{P}_x(S_k = y).$$

The function p_k is symmetric in x and y , and one has $p_k(x, y) = p_k(0, y - x) := p_k(y - x)$. Define

$$(2.1) \quad f_k(x) = \frac{8}{\pi^2 k^2} \exp\left(-2 \frac{\|x\|^2}{k}\right).$$

A well-known estimate (see Proposition 2.1.2(b) in [25]) shows that for some positive constants c and C

$$(2.2) \quad \forall k \geq 1, \quad \mathbb{P}\left(\max_{\ell \leq k} \|S_\ell\| \geq r\right) \leq C \cdot e^{-c \cdot r^2/k}.$$

Furthermore, for any fixed $\alpha < 2/3$, one has for all $k \geq 1$ and x with $p_k(x) > 0$ and $\|x\| \leq k^\alpha$ (see Proposition 1.2.5 in [24]):

$$(2.3) \quad p_k(x) = f_k(x)(1 + \mathcal{O}(k^{3\alpha-2})).$$

One deduces in particular the following useful estimate:

$$(2.4) \quad \mathbb{P}(\|S_k\|^2 \leq k/R) = \mathcal{O}(R^{-2}).$$

The discrete Green's function G_d is defined by

$$G_d(x, y) = \sum_{k \geq 0} \mathbb{P}_x(S_k = y) \quad \text{and} \quad \frac{G_d(x, y)}{G_d(0, 0)} = \mathbb{P}_x(H_y < \infty).$$

We also write $G_d(x) = G_d(0, x)$, and recall that G_d is symmetric, and satisfies $G_d(x, y) = G_d(y - x)$.

The continuous Green's function $G(x, y)$ is also symmetric and satisfies $G(x, y) = G(0, y - x) =: G(y - x)$. It is defined for $z \in \mathbb{R}^4$ nonzero, by

$$(2.5) \quad G(z) = \frac{1}{2\pi^2} \cdot \frac{1}{\|z\|^2}.$$

These two functions are linked by the relation (see Theorem 4.3.1 in [25]): for $x \in \mathbb{Z}^4$

$$(2.6) \quad G_d(x) = 4G(x) + \mathcal{O}\left(\frac{1}{1 + \|x\|^4}\right).$$

We also use the following (see Proposition 6.5.1 and 6.5.2 in [25]): there exists a constant $C > 0$, such that for all x and $r > 0$,

$$(2.7) \quad \mathbb{P}_x(H_{B(0,r)} < \infty) \leq C \cdot \frac{r^2}{1 + \|x\|^2}.$$

Finally, we prove two useful estimates on the heat kernel $p_k(x)$:

CLAIM 2.1. *Let $x \in \mathbb{Z}^d$ and $k \in \mathbb{N}$ be such that $\sqrt{k} \leq \|x\| \leq k^{3/5}$. Then there exists a positive constant C (independent of x) so that for all $i \leq k$ we have*

$$p_i(x) \leq C f_k(x).$$

PROOF. Suppose first that $i \leq k^{1-\varepsilon}$, for some $\varepsilon > 0$ to be fixed later. Then one can use (2.2) which gives

$$p_i(x) \leq \mathbb{P}(\|S_i\| \geq \|x\|) \lesssim \exp\left(-c \frac{\|x\|^2}{i}\right) \lesssim \exp\left(-\frac{c}{2} k^\varepsilon - 2 \frac{\|x\|^2}{k}\right) \lesssim f_k(x),$$

using for the third inequality that for k large enough, $\|x\|^2/i \geq \max(k^\varepsilon, (4/c) \cdot \|x\|^2/k)$.

Suppose next that $k^{1-\varepsilon} \leq i \leq k/2$. Now choose ε such that $3/(5(1-\varepsilon)) < 2/3$. Then one can use the local CLT (2.3), which gives

$$p_i(x) \lesssim \frac{1}{i^2} \exp\left(-\frac{2\|x\|^2}{i}\right) = \frac{\|x\|^4}{i^2} \exp\left(-\frac{\|x\|^2}{i}\right) \cdot \frac{1}{\|x\|^4} \exp\left(-\frac{\|x\|^2}{i}\right).$$

Using that the function $y^2 e^{-cy}$ is upper bounded by a constant, the assumption that $i \leq k/2$ and $\|x\| \geq \sqrt{k}$, we obtain

$$p_i(x) \lesssim \frac{1}{k^2} \exp\left(-\frac{2\|x\|^2}{k}\right)$$

and this completes the proof in the case $i \leq k/2$.

Suppose finally that $k/2 \leq i \leq k$. Then we conclude the proof

$$p_i(x) \lesssim \frac{1}{i^2} \exp\left(-\frac{2\|x\|^2}{i}\right) \lesssim \frac{1}{k^2} \exp\left(-\frac{2\|x\|^2}{k}\right). \quad \square$$

Recall next the notation $n_\alpha = n/(\log n)^\alpha$.

CLAIM 2.2. *Let $i, k, n \in \mathbb{N}$ and $x, z \in \mathbb{Z}^4$ satisfy $k \geq n_\alpha$, $\|x\| \leq \sqrt{n}(\log n)^2$, $i \leq k/(\log n)^\beta$ and $\|z\| \leq \sqrt{i}(\log n)^\gamma$ with α, β and γ satisfying $\beta > \alpha + 4$ and $4 + 2\gamma + \alpha - \beta < 0$. Then we have as $n \rightarrow \infty$*

$$f_{k-i}(x-z) = f_k(x) \cdot (1 + o(1)).$$

PROOF. First, k and i depend on n , and as $n \rightarrow \infty$

$$\frac{1}{k^2} = \frac{1}{(k-i)^2} \cdot (1 + o(1)).$$

We next turn to the exponential terms in the expression for f_k . We have

$$\begin{aligned} \frac{\|x\|^2}{k} - \frac{\|x-z\|^2}{(k-i)} &= \frac{(k-i)\|x\|^2 - k(\|x\|^2 - 2\langle x, z \rangle + \|z\|^2)}{k(k-i)} \\ &= \frac{2k\langle x, z \rangle - k\|z\|^2 - i\|x\|^2}{k(k-i)}. \end{aligned}$$

It suffices to prove that this last expression tends to 0 as $n \rightarrow \infty$. By the assumption,

$$\frac{i\|x\|^2}{k(k-i)} \lesssim (\log n)^{\alpha+4-\beta} \quad \text{and} \quad \frac{k\|z\|^2}{k(k-i)} \lesssim (\log n)^{2\gamma-\beta},$$

and since $\alpha + 4 < \beta$ and $2\gamma < \beta$ they both tend to 0. Finally, by Cauchy–Schwarz we get

$$\frac{|\langle x, z \rangle|}{k-i} \leq \frac{\|x\|\|z\|}{k-i} \lesssim (\log n)^{2+\gamma-\frac{\alpha}{2}-\frac{\beta}{2}} \rightarrow 0 \quad \text{as } n \rightarrow \infty,$$

again by using the assumption on α, β and γ and this completes the proof. $\square$

2.2. *A decomposition formula for the capacity.* Recall the last passage decomposition formula (see for instance Proposition 4.6.4 in [25]): for any finite set $A \subseteq \mathbb{Z}^d$ and $x \notin A$,

$$(2.8) \quad \mathbb{P}_x(H_A < \infty) = \sum_{y \in A} G_d(x, y) \cdot \mathbb{P}_y(H_A^+ = \infty).$$

We also recall two well-known formulas for the capacity of a finite set $A \subseteq \mathbb{Z}^d$. First,

$$(2.9) \quad \text{Cap}(A) = \lim_{\|x\| \rightarrow \infty} \frac{\mathbb{P}_x(H_A < \infty)}{G_d(x, 0)},$$

and for any $y \in A$,

$$(2.10) \quad \frac{\mathbb{P}_y(H_A^+ = \infty)}{\text{Cap}(A)} = \lim_{\|x\| \rightarrow \infty} \mathbb{P}_x(S(H_A) = y | H_A < \infty).$$

The first formula is obtained through the last passage decomposition formula (2.8) and the definition of the capacity (1.1), and the second is Theorem 2.1.3 of Lawler's book [24].

PROOF OF PROPOSITION 1.6. Consider two finite subsets A and B of $\mathbb{Z}^d$. One has

$$(2.11) \quad \begin{aligned} & \mathbb{P}_x(H_{A \cup B} < \infty) \\ &= \mathbb{P}_x(H_A < \infty) + \mathbb{P}_x(H_B < \infty) - \mathbb{P}_x(H_A < \infty, H_B < \infty) \\ &= \mathbb{P}_x(H_A < \infty) + \mathbb{P}_x(H_B < \infty) - \mathbb{P}_x(H_A \leq H_B < \infty) \\ & \quad - \mathbb{P}_x(H_B \leq H_A < \infty) + \mathbb{P}_x(H_A = H_B < \infty). \end{aligned}$$

Consider now the term $\mathbb{P}_x(H_A \leq H_B < \infty)$. Conditioning on the possible hitting point in A and using the Markov property yield:

$$\begin{aligned} & \mathbb{P}_x(H_A \leq H_B < \infty) \\ &= \sum_{y \in A} \mathbb{P}_x(S(H_{A \cup B}) = y, H_{A \cup B} < \infty) \mathbb{P}_y(H_B < \infty) \\ &= \mathbb{P}_x(H_{A \cup B} < \infty) \sum_{y \in A} \mathbb{P}_x(S(H_{A \cup B}) = y | H_{A \cup B} < \infty) \mathbb{P}_y(H_B < \infty). \end{aligned}$$

Then use (2.9) and (2.10) to obtain

$$\lim_{x \rightarrow \infty} \frac{\mathbb{P}_x(H_A \leq H_B < \infty)}{G_d(0, x)} = \sum_{y \in A} \mathbb{P}_y(H_{A \cup B}^+ = \infty) \mathbb{P}_y(H_B < \infty).$$

Finally, by using the last passage formula (2.8), we get the desired limit

$$\lim_{x \rightarrow \infty} \frac{\mathbb{P}_x(H_A \leq H_B < \infty)}{G_d(0, x)} = \sum_{y \in A} \sum_{z \in B} \mathbb{P}_y(H_{A \cup B}^+ = \infty) G_d(y, z) \mathbb{P}_z(H_B^+ = \infty).$$

By symmetry one also has

$$\lim_{x \rightarrow \infty} \frac{\mathbb{P}_x(H_B \leq H_A < \infty)}{G_d(0, x)} = \sum_{y \in B} \sum_{z \in A} \mathbb{P}_y(H_{A \cup B}^+ = \infty) G_d(y, z) \mathbb{P}_z(H_A^+ = \infty).$$

We also obtain the existence of the limit $\varepsilon(A, B)$ of $\mathbb{P}_x(H_A = H_B < \infty)/G_d(0, x)$, as $x \rightarrow \infty$, since in (2.11) all other ratios converge. To conclude, note that

$$\mathbb{P}_x(H_A = H_B < \infty) \leq \mathbb{P}_x(H_{A \cap B} < \infty),$$

which gives $\varepsilon(A, B) \leq \text{Cap}(A \cap B)$. $\square$

We will apply successively the decomposition of Proposition 1.6. To this end, we define for $i \geq 1$ and $j \leq 2^i$,

$$\mathcal{R}_n^{(i,j)} := \mathcal{R}[(j-1)2^{-i}n, j2^{-i}n].$$

PROPOSITION 2.3. *Fix $p \in \mathbb{N}$. Then we have*

$$\text{Cap}(\mathcal{R}_n) = \sum_{j=1}^{2^p} \text{Cap}(\mathcal{R}_n^{(p,j)}) - \sum_{i=1}^p \sum_{j=1}^{2^{i-1}} \chi_n(i, j) + \varepsilon_n,$$

where $\mathbb{E}[\varepsilon_n^2] = \mathcal{O}((\log n)^2)$ and

$$\chi_n(i, j) := \chi(\mathcal{R}_n^{(i, 2^{j-1})}, \mathcal{R}_n^{(i, 2^j)}) + \chi(\mathcal{R}_n^{(i, 2^j)}, \mathcal{R}_n^{(i, 2^{j-1})}).$$

PROOF. The proof follows directly by applying repeatedly Proposition 1.6 to $\mathcal{R}_n$. Moreover, from Proposition 1.6 we have that in every subdivision the term ε is upper bounded by the size of the intersection of two independent ranges. A straightforward calculation shows that this has expectation $\log n$ (see, for instance, [24], Section 3.4). The bound on the second moment follows from [31], Lemma 3.1. Since we are only taking a finite sum, the result follows. $\square$

3. Strong law of large numbers. In this section, we prove Theorem 1.1. The main part of the proof consists in obtaining good bounds on the first and second moments of the cross term $\chi_n(1, 1)$ appearing in the decomposition formula of the capacity. More precisely, we show that

LEMMA 3.1. *The first and second moments of $\chi_n(1, 1)$ satisfy*

$$(3.1) \quad \mathbb{E}[\chi_n(1, 1)] \lesssim n \cdot \frac{\log \log n}{(\log n)^2} \quad \text{and}$$

$$(3.2) \quad \mathbb{E}[\chi_n(1, 1)^2] \lesssim n^2 \cdot \frac{(\log \log n)^2}{(\log n)^4}.$$

Then, in Section 3.3, we deduce the strong law of large numbers by using our decomposition of the capacity, Proposition 2.3. Section 3.1 is devoted to some preliminary facts needed for the proof of Lemma 3.1.

3.1. *Preliminaries.* We first recall a standard fact, which directly follows from (2.6): the mean time a walk spends in a ball of radius R is of order R^2 . More precisely,

$$(3.3) \quad \sum_{k \in \mathbb{N}} \mathbb{P}(\|S(k)\| \leq R) = \sum_{x \in \mathcal{B}(0, R)} G_d(0, x) = \mathcal{O}(R^2).$$

Next, we present a lemma which is needed in the second moment estimate, and which deals with intersecting the trace of a path $\mathcal{R}_n$ by two independent walks starting far apart. The proof follows basically from estimates of Lawler [24]. Recall that $n_\alpha = n/(\log n)^\alpha$.

LEMMA 3.2. *Let $\alpha > 0$. Consider three independent walks S, S^1, S^2 starting at the origin, and let $x, y \in \mathbb{Z}^d$ with $\|x\|^2, \|y\|^2 \geq n_\alpha$. There is a constant $C > 0$, such that for n large enough,*

$$(3.4) \quad \begin{aligned} & \mathbb{P}((x + \mathcal{R}^1[0, \infty)) \cap \mathcal{R}_n \neq \emptyset, (y + \mathcal{R}^2[0, \infty)) \cap \mathcal{R}_n \neq \emptyset) \\ & \leq C \cdot \left(\frac{\log \log n}{\log n} \right)^2. \end{aligned}$$

PROOF. The proof consists in showing that even if the two events considered in (3.4) are not independent, one can still dissociate them. Consider the two random times

$$\sigma_x = \inf\{k : S(k) \in x + \mathcal{R}^1[0, \infty)\} \quad \text{and} \quad \sigma_y = \inf\{k : S(k) \in y + \mathcal{R}^2[0, \infty)\}.$$

Note that σ_x (resp., σ_y) is independent of S^2 (resp., of S^1). We can express the nonintersection event in terms of σ_x and σ_y :

$$\begin{aligned} & \mathbb{P}((x + \mathcal{R}^1[0, \infty)) \cap \mathcal{R}_n \neq \emptyset, ((y + \mathcal{R}^2[0, \infty)) \cap \mathcal{R}_n \neq \emptyset) \\ & \leq \mathbb{P}(\sigma_x \leq \sigma_y \leq n) + \mathbb{P}(\sigma_y \leq \sigma_x \leq n). \end{aligned}$$

By symmetry, it is enough to deal with the first probability on the right-hand side. Conditioning on S^1 and σ_x , we get

$$\begin{aligned} \mathbb{P}(\sigma_x \leq \sigma_y \leq n) & \leq \mathbb{E}[\mathbb{1}(\sigma_x \leq n) \mathbb{P}_{S(\sigma_x)}(\mathcal{R}[0, n - \sigma_x] \cap (y + \mathcal{R}^2[0, \infty)) \neq \emptyset)] \\ & \leq \mathbb{E}[\mathbb{1}(\sigma_x \leq n) \mathbb{P}_{S(\sigma_x)}(\mathcal{R}[0, n] \cap (y + \mathcal{R}^2[0, \infty)) \neq \emptyset)]. \end{aligned}$$

Note that since $\|y\|^2 \geq n_\alpha$, we have using (2.7),

$$\mathbb{P}(\sigma_x \leq n, \|S(\sigma_x) - y\|^2 \leq n_{\alpha+2}) \leq \mathbb{P}(H_{\mathcal{B}(y, \sqrt{n_{\alpha+2}})} < \infty) \lesssim \frac{1}{(\log n)^2}.$$

Now, on the event $\{\sigma_x \leq n, \|S(\sigma_x) - y\|^2 \geq n_{\alpha+2}\}$, [24], Theorem 4.3.3, shows that

$$\mathbb{P}_{S(\sigma_x)}((y + \mathcal{R}^2[0, \infty)) \cap \mathcal{R}[0, n] \neq \emptyset) \lesssim \frac{\log \log n}{\log n}.$$

Thus by another application of [24], Theorem 4.3.3, we get

$$\mathbb{P}(\sigma_x \leq \sigma_y \leq n) \lesssim \mathbb{P}(\sigma_x \leq n) \cdot \frac{\log \log n}{\log n} \lesssim \left(\frac{\log \log n}{\log n} \right)^2,$$

and this completes the proof. $\square$

3.2. First and second moment estimates of the cross terms.

PROOF OF LEMMA 3.1. Note first that by reversibility of the walk, $\chi_n(1, 1)$ is equal in law to $\chi(\mathcal{R}_{n/2}, \tilde{\mathcal{R}}_{n/2}) + \chi(\tilde{\mathcal{R}}_{n/2}, \mathcal{R}_{n/2})$, with $\mathcal{R}$ and $\tilde{\mathcal{R}}$ the ranges of two independent walks S and $\tilde{S}$. By symmetry (and Cauchy–Schwarz), it is enough to bound the first and second moments of $\chi(\mathcal{R}_{n/2}, \tilde{\mathcal{R}}_{n/2})$. However, to avoid annoying factors $1/2$ everywhere, we bound the term $\chi(\mathcal{R}_n, \tilde{\mathcal{R}}_n)$. Recall that for any finite sets A and B , we have by definition of $\chi(A, B)$ and using also the last exit formula (2.8):

$$\chi(A, B) = \sum_{y \in A} \mathbb{P}_y(H_{A \cup B}^+ = \infty) \cdot \mathbb{P}_y(H_B < \infty).$$

Even though the first moment bound (3.1) follows from (3.2) by using Jensen's inequality, it is interesting to include a direct proof of (3.1). Indeed, it serves as a warmup for the proof of the second moment estimate.

So let us prove (3.1). For this we need to consider two additional independent random walks S^1 and $\tilde{S}^1$ starting from the origin and also independent of S and $\tilde{S}$. Denote their ranges by $\mathcal{R}^1$ and $\tilde{\mathcal{R}}^1$, respectively. Also, we use the shorthand $\mathcal{R}$ for $\mathcal{R}[0, \infty)$ and $\mathcal{R}_+$ for $\mathcal{R}[1, \infty)$. We start with a bound on $\chi(\mathcal{R}_n, \tilde{\mathcal{R}}_n)$, which is a sum of n terms

$$\begin{aligned} & \sum_{k=1}^n \mathbb{1}_{\{S_k \notin \mathcal{R}_{k-1}\}} \mathbb{P}((S_k + \mathcal{R}_+^1) \cap (\mathcal{R}_n \cup \tilde{\mathcal{R}}_n) = \emptyset, (S_k + \tilde{\mathcal{R}}^1) \cap \tilde{\mathcal{R}}_n \neq \emptyset | S, \tilde{S}) \\ & \leq \sum_{k=1}^n \mathbb{P}((S_k + \mathcal{R}_+^1) \cap \mathcal{R}_n = \emptyset | S) \cdot \mathbb{P}((S_k + \tilde{\mathcal{R}}^1) \cap \tilde{\mathcal{R}}_n \neq \emptyset | S, \tilde{S}). \end{aligned}$$

Taking expectation on both sides and choosing any $\alpha > 2$, we bound $\mathbb{E}[\chi(\mathcal{R}_n, \tilde{\mathcal{R}}_n)]$ as follows:

$$\begin{aligned} & \sum_{k=1}^n \sum_{\|x\|^2 \geq n_\alpha} \mathbb{P}((x + \mathcal{R}_+^1) \cap \mathcal{R}_n = \emptyset, S_k = x) \cdot \mathbb{P}((x + \tilde{\mathcal{R}}^1) \cap \tilde{\mathcal{R}}_n \neq \emptyset) + \mathcal{O}(n_\alpha) \\ & \lesssim \sum_{k=1}^n \mathbb{P}((S_k + \mathcal{R}_+^1) \cap \mathcal{R}_n = \emptyset) \cdot \frac{\log \log n}{\log n} + \mathcal{O}(n_\alpha) \\ & \lesssim \sum_{k=n_\alpha}^{n-n_\alpha} \mathbb{P}((S_k + \mathcal{R}_+^1) \cap \mathcal{R}_n = \emptyset) \cdot \frac{\log \log n}{\log n} + \mathcal{O}(n_\alpha) \lesssim n \cdot \frac{\log \log n}{(\log n)^2}, \end{aligned}$$

where we used (3.3) at the first line and Lawler's results [24], Theorem 3.5.1 and 4.3.3, at the second and third lines. This proves (3.1).

Let $\{S, \tilde{S}, S^1, S^2, \tilde{S}^1, \tilde{S}^2\}$ be independent random walks all starting at the origin. Fix $\alpha > 4$, and n and introduce the following event $\mathcal{A}_k = \{\|S(k)\|^2 \geq n_\alpha\}$. Now using (3.3) at the second line and Lemma 3.2 at the fourth line, we bound the second moment of $\chi(\mathcal{R}_n, \tilde{\mathcal{R}}_n)$ by

$$\begin{aligned}
 & \sum_{k_1=1}^n \sum_{k_2=1}^n \mathbb{P}((S_{k_i} + \mathcal{R}_+^i) \cap \mathcal{R}_n = \emptyset, (S_{k_i} + \tilde{\mathcal{R}}^i) \cap \tilde{\mathcal{R}}_n \neq \emptyset, \forall i = 1, 2) \\
 & \leq \sum_{k_1, k_2} \mathbb{P}(\mathcal{A}_{k_i}, (S_{k_i} + \mathcal{R}_+^i) \cap \mathcal{R}_n = \emptyset, (S_{k_i} + \tilde{\mathcal{R}}^i) \cap \tilde{\mathcal{R}}_n \neq \emptyset, \forall i = 1, 2) \\
 & \quad + \mathcal{O}(n \cdot n_\alpha) \\
 & \leq \sum_{k_1, k_2} \sum_{x_1, x_2} \mathbb{P}(\mathcal{A}_{k_i}, S_{k_i} = x_i, (x_i + \mathcal{R}_+^i) \cap \mathcal{R}_n = \emptyset, \forall i = 1, 2) \\
 & \quad \times \mathbb{P}((x_i + \tilde{\mathcal{R}}^i) \cap \tilde{\mathcal{R}}_n \neq \emptyset, \forall i = 1, 2) + \mathcal{O}(n \cdot n_\alpha) \\
 & \lesssim \sum_{k_1, k_2} \mathbb{P}(\mathcal{A}_{k_i}, (S_{k_i} + \mathcal{R}_+^i) \cap \mathcal{R}_n = \emptyset, \forall i = 1, 2) \cdot \frac{\log^2 \log n}{\log^2 n} + \mathcal{O}(n \cdot n_\alpha) \\
 & \lesssim \sum_{n_\alpha \leq k_1 \leq k_2 - n_\alpha \leq n - 2n_\alpha} \mathbb{P}\left(\bigcap_{i=1,2} (S_{k_i} + \mathcal{R}_+^i) \cap \mathcal{R}_n = \emptyset\right) \\
 & \quad \times \frac{\log^2 \log n}{\log^2 n} + \mathcal{O}(n \cdot n_\alpha).
 \end{aligned}$$

We deal now with the nonintersection terms for which we removed the space constraints but we added time constraints. From the walk S , we form two walks stemming from position S_m with $m = [(k_1 + k_2)/2]$. One walk goes backward, and is denoted S^3 , and another one goes forward and is denoted S^4 . Translating the origin to S_m we obtain using [24], Theorem 3.5.1,

$$\begin{aligned}
 & \mathbb{P}((S_{k_i} + \mathcal{R}_+^i) \cap \mathcal{R}_n = \emptyset, \forall i = 1, 2) \\
 & \leq \mathbb{P}((S_{m/2}^3 + \mathcal{R}_+^1) \cap \mathcal{R}^3[0, m] = \emptyset) \\
 & \quad \times \mathbb{P}((S_{m/2}^4 + \mathcal{R}_+^2) \cap \mathcal{R}^4[0, n - m] = \emptyset) = \mathcal{O}\left(\frac{1}{(\log n)^2}\right),
 \end{aligned}$$

which proves (3.2). $\square$

3.3. Strong law of large numbers. We are now ready for the proof of Theorem 1.1. The first step is to use the dyadic decomposition of the capacity to produce self-similar *independent* terms at a smaller scale. If this scale is well chosen,

then the result of the previous section shows that the cross terms $\chi_n(i, j)$ are negligible, by a simple application of Chebyshev's inequality. On the other hand, using Borel–Cantelli and Chebyshev's inequality again, one can show that the self-similar part converges almost surely, at least along some subsequence growing sub-exponentially fast. Finally, using the monotonicity of the capacity we deduce the convergence along the full sequence.

PROOF OF THEOREM 1.1. Choose L as a function of n , such that $(\log n)^4 \leq 2^L \leq 2(\log n)^4$. In particular, one has $L \asymp \log \log n$. Proposition 2.3 shows that

$$(3.5) \quad \text{Cap}(\mathcal{R}_n) = \sum_{j=1}^{2^L} \text{Cap}(\mathcal{R}_n^{(L,j)}) - \xi_n + \varepsilon_n,$$

where $\mathbb{E}[\varepsilon_n] = \mathcal{O}(2^L \log n) = \mathcal{O}((\log n)^5)$, and

$$\xi_n = \sum_{i=1}^L \sum_{j=1}^{2^{i-1}} \chi_n(i, j).$$

Now (3.2) shows that for all $i \leq L$ and $j \leq 2^{i-1}$

$$\mathbb{E}[\chi_n(i, j)^2] \lesssim n^2 \cdot \frac{(\log \log n)^2}{2^{2i} (\log n)^4}.$$

Using Cauchy–Schwarz and the independence of the $(\chi_n(i, j))_j$ for any fixed i , one obtains

$$\begin{aligned} \text{Var}(\xi_n) &\leq L \cdot \sum_{i=1}^L \text{Var}\left(\sum_{j=1}^{2^{i-1}} \chi_n(i, j)\right) = L \cdot \sum_{i=1}^L 2^{i-1} \text{Var}(\chi_n(i, 1)) \\ &= \mathcal{O}\left(n^2 \cdot \frac{(\log \log n)^3}{(\log n)^4}\right). \end{aligned}$$

This together with Chebyshev's inequality give

$$(3.6) \quad \mathbb{P}\left(|\xi_n - \mathbb{E}[\xi_n]| > \varepsilon \frac{n}{\log n}\right) \lesssim \frac{(\log \log n)^3}{(\log n)^2},$$

where $\varepsilon > 0$. On the other hand, using the trivial bound $\text{Cap}(\mathcal{R}_n) \leq |\mathcal{R}_n| \leq n$, and Chebyshev's inequality again for a sum of independent terms, we get

$$(3.7) \quad \mathbb{P}\left(\sum_{j=1}^{2^L} |\text{Cap}(\mathcal{R}_n^{(L,j)}) - \mathbb{E}[\text{Cap}(\mathcal{R}_n^{(L,j)})]| > \varepsilon \frac{n}{\log n}\right) \lesssim \frac{1}{(\log n)^2}.$$

Now consider the subsequence $a_n = \exp(n^{3/4})$, and observe that it satisfies

$$\lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} = 1 \quad \text{and} \quad \sum_n \frac{(\log \log a_n)^3}{(\log a_n)^2} < \infty.$$

Therefore (3.1), (3.6) and Borel–Cantelli show that almost surely

$$\lim_{n \rightarrow \infty} \frac{\log a_n}{a_n} \cdot \xi_{a_n} = 0.$$

Similarly using the bound $\mathbb{E}[\varepsilon_n] = \mathcal{O}((\log n)^5)$, we deduce using Markov’s inequality that almost surely

$$\lim_{n \rightarrow \infty} \frac{\log a_n}{a_n} \cdot \varepsilon_{a_n} = 0.$$

Finally, using (1.4), (3.5), (3.7) and Borel–Cantelli again, we deduce that almost surely

$$\lim_{n \rightarrow \infty} \frac{\log a_n}{a_n} \cdot \text{Cap}(\mathcal{R}_{a_n}) = \frac{\pi^2}{8}.$$

To conclude, first observe that $\text{Cap}(\mathcal{R}_n)$ is nondecreasing, since for any $A \subset B$, one has $\text{Cap}(A) \leq \text{Cap}(B)$. Thus if for $n \geq 1$, we define k_n as the unique integer, such that $a_{k_n} \leq n < a_{k_n+1}$, we have

$$\text{Cap}(\mathcal{R}[0, a_{k_n}]) \leq \text{Cap}(\mathcal{R}[0, n]) \leq \text{Cap}(\mathcal{R}[0, a_{k_n+1}]).$$

Since a_{k_n+1}/a_{k_n} goes to 1, as $n \rightarrow \infty$, the sequence $(\log n/n) \cdot \text{Cap}(\mathcal{R}[0, n])$ converges. $\square$

REMARK 3.3. Note that in the last part of our proof, we followed the strategy of Dvoretzky and Erdős in their pioneering work on the range [13]. They first proved an almost sure limiting result along a subsequence growing subexponentially fast (using also Chebyshev’s inequality and good bounds on the variance), and then deduced the strong law of large numbers using a monotonicity argument. The idea that a decomposition like (1.6) could be useful in obtaining sharp variance bounds and a central limit theorem came much later, in Le Gall’s papers [28, 29].

4. Existence and definition of the limiting term. The goal of this section is to give a precise definition of the limiting term appearing in Theorem 1.2. We also prove a Carleman’s condition for the sum approximating it.

4.1. Carleman’s condition. We recall that Carleman’s condition holds for a nonnegative random variable X , if its sequence of moments $m_p := \mathbb{E}[X^p]$ satisfies:

$$\sum_{p \geq 1} (m_p)^{-\frac{1}{2p}} = \infty.$$

PROPOSITION 4.1. *Let*

$$X = \int_0^1 \int_0^1 \frac{1}{\|\beta_s - \tilde{\beta}_t\|^2} ds dt,$$

where β and $\tilde{\beta}$ are two independent standard 4-dimensional Brownian motions. There exist positive constants c and C , such that

$$c^p p^p \leq \mathbb{E}[X^p] \leq C^p \cdot p^{2p},$$

for all integers $p \geq 1$. In particular, the upper bound implies that Carleman's condition holds for X .

REMARK 4.2. Note that by reversibility of the Brownian path, and scaling invariance, one has

$$X \stackrel{(d)}{=} 2 \int_0^{1/2} \int_{1/2}^1 \frac{1}{\|\beta_s - \beta_t\|^2} ds dt.$$

PROOF. We first prove the upper bound. By using Jensen's inequality we get, for any integer p ,

$$\mathbb{E}[X^p] \lesssim \int_0^1 dt \mathbb{E} \left[\left(\int_0^1 \frac{ds}{\|\beta_s - \tilde{\beta}_t\|^2} \right)^p \right].$$

We define $\mathcal{C}_p := \{0 \leq s_1 \leq \dots \leq s_p \leq 1\}$, and by symmetry

$$(4.1) \quad \mathbb{E}[X^p] \lesssim p! \int_0^1 dt \int_{\mathcal{C}_p} \mathbb{E} \left[\frac{1}{\|\beta_{s_1} - \tilde{\beta}_t\|^2 \dots \|\beta_{s_p} - \tilde{\beta}_t\|^2} \right] ds_1 \dots ds_p.$$

By using standard properties of the Brownian motion, we write

$$\begin{aligned} & \int_{\mathcal{C}_p} \mathbb{E} \left[\prod_{i=1}^p \frac{1}{\|\beta_{s_i} - \tilde{\beta}_t\|^2} \right] \prod_{i=1}^p ds_i \\ (4.2) \quad &= \int_{\mathcal{C}_p} \mathbb{E} \left[\prod_{i=1}^{p-1} \frac{1}{\|\beta_{s_i} - \tilde{\beta}_t\|^2} \times \mathbb{E} \left(\frac{1}{\|\beta_{s_p} - \tilde{\beta}_t\|^2} \middle| (\beta_u)_{u \leq s_{p-1}}, \tilde{\beta}_t \right) \right] \prod_{i=1}^p ds_i \\ &= \int_{\mathcal{C}_{p-1}} \mathbb{E} \left[\prod_{i=1}^{p-1} \frac{1}{\|\beta_{s_i} - \tilde{\beta}_t\|^2} \int_0^{1-s_{p-1}} \mathbb{E} \left(\frac{1}{\|\beta_s - x_p\|^2} \middle| x_p \right) ds \right] \prod_{i=1}^{p-1} ds_i, \end{aligned}$$

with $x_p = \beta_{s_{p-1}} - \tilde{\beta}_t$. Now we need the following two lemmas.

LEMMA 4.3. *One has*

$$\int_0^1 \mathbb{E} \left[\frac{(a + |\log \|\beta_s - x\||)^k}{\|\beta_s - x\|^2} \right] ds \lesssim \sum_{\ell=0}^{k+1} (4k)^\ell \cdot (a + 1 + |\log \|x\||)^{k+1-\ell},$$

uniformly in $x \in \mathbb{R}^4 \setminus \{0\}$, $a \geq 0$ and $k \geq 0$.

LEMMA 4.4. For all $k \geq 1$ and $a \geq k$,

$$\int_0^1 \mathbb{E}[(a + |\log \|\tilde{\beta}_t\||)^k] dt \lesssim k \cdot a^k.$$

We prove these two lemmas in the [Appendix](#). Let us now conclude the proof of the proposition. First one can use Lemma 4.3 and by induction, we can bound the last integral in (4.2) by

$$C^p \cdot \sum_{\ell=0}^p \alpha_\ell \cdot (4p)^\ell \cdot \mathbb{E}[(p + |\log \|\tilde{\beta}_t\||)^{p-\ell}],$$

where C is a constant and

$$\alpha_\ell := \#\{1 \leq i_1 \leq \dots \leq i_\ell \leq p\} = \binom{p+\ell-1}{\ell}.$$

Then using Lemma 4.4, (4.1) and (4.2) we obtain the upper bound

$$\mathbb{E}[X^p] \lesssim C^p \sum_{\ell=0}^p \binom{p+\ell-1}{\ell} (4p)^\ell \cdot (p-\ell) \cdot p^{p-\ell} \lesssim \tilde{C}^p \cdot p^{2p},$$

where $\tilde{C}$ is a positive constant and this proves the upper bound.

Now we prove the lower bound. Define

$$\Gamma(p) := \inf_{\substack{x \in \mathcal{B}(0, 2/\sqrt{p}) \\ z, z' \in \mathcal{B}(0, 1/\sqrt{p})}} \int_0^{1/p} \int_0^{1/p} \mathbb{E}_{z, z'} \left[\frac{\mathbb{1}(\beta_{1/p}, \tilde{\beta}_{1/p} \in \mathcal{B}(0, 1/\sqrt{p}))}{\|x + \beta_s - \tilde{\beta}_t\|^2} \right] ds dt.$$

Note that by scaling $\Gamma(p) = \Gamma(1)/p$, and that

$$\Gamma(1) \geq \frac{1}{16} \mathbb{P}\left(\sup_{s \leq 1} \|\beta_s\| \leq 1\right)^2 := c_0 > 0.$$

Now define

$$\mathcal{D} = \left[0, \frac{1}{p}\right] \times \dots \times \left[\frac{p-1}{p}, 1\right] \quad \text{and} \quad A := \bigcap_{i=1}^p \left\{ \beta_{i/p}, \tilde{\beta}_{i/p} \in \mathcal{B}\left(0, \frac{1}{\sqrt{p}}\right) \right\}.$$

Then, for any permutation σ of the set $\{1, \dots, n\}$, one has by the Markov property and standard properties of Brownian motion,

$$\begin{aligned} & \int_{\mathcal{C}_p} \int_{\mathcal{C}_p} \mathbb{E} \left[\prod_{i=1}^p \frac{1}{\|\beta_{s_i} - \tilde{\beta}_{t_{\sigma(i)}}\|^2} \right] \prod_{i=1}^p ds_i dt_i \\ & \geq \int_{\mathcal{D}} \int_{\mathcal{D}} \mathbb{E} \left[\prod_{i=1}^p \frac{\mathbb{1}(A)}{\|\beta_{s_i} - \tilde{\beta}_{t_{\sigma(i)}}\|^2} \right] \prod_{i=1}^p ds_i dt_i \end{aligned}$$

$$\begin{aligned}
&= \int_{\mathcal{D}} \int_{\mathcal{D}} \mathbb{E} \left[\prod_{i=1}^p \frac{\mathbb{1}(A)}{\|x_i + (\beta_{s_i} - \beta_{i/p}) - (\tilde{\beta}_{t_{\sigma(i)}} - \tilde{\beta}_{\sigma(i)/p})\|^2} \right] \prod_{i=1}^p ds_i dt_i \\
&\geq \Gamma(p)^p \geq (c_0/p)^p,
\end{aligned}$$

with $x_i = \beta_{i/p} - \tilde{\beta}_{\sigma(i)/p}$, for $i = 1, \dots, p$. Now one has

$$\mathbb{E}[X^p] \geq (p!)^2 \inf_{\sigma} \int_{\mathcal{C}_p} \int_{\mathcal{C}_p} \prod ds_i \prod dt_i \mathbb{E} \left[\prod_{i=1}^p \frac{1}{\|\beta_{s_i} - \tilde{\beta}_{t_{\sigma(i)}}\|^2} \right],$$

and this gives the lower bound using the previous bound and Stirling's formula. $\square$

4.2. The limiting term. We have now all the ingredients to define properly the term $\gamma_G([0, 1]^2)$, appearing in Theorem 1.2. First, define the following subsquares of $[0, 1]^2$:

$$A_{i,j} = [(2j-2)2^{-i}, (2j-1)2^{-i}] \times [(2j-1)2^{-i}, (2j)2^{-i}],$$

for $i \geq 1$ and $j \leq 2^{i-1}$. Define also $\mathcal{C} = \{s, t \in [0, 1] : s \leq t\}$, the closure of the union of all these squares.

A straightforward computation shows that

$$\mathbb{E} \left[\int_0^1 \int_0^1 G(\beta_s, \beta_t) ds dt \right] = \infty.$$

However, if we consider disjoint intervals, then this expectation is finite as we already proved in Proposition 4.1, that is,

$$\mathbb{E} \left[\int_0^{1/2} \int_{1/2}^1 G(\beta_s, \beta_t) ds dt \right] < \infty,$$

and by scaling the same fact holds when integrating over any of the squares $A_{i,j}$. This observation is at the heart of the following proposition.

PROPOSITION 4.5. *Let β be a standard 4-dimensional Brownian motion. Then the limit $\gamma_G(\mathcal{C}) := \sum_{i=1}^{\infty} \sum_{j=1}^{2^{i-1}} \gamma_G(A_{i,j})$ exists in L^2 , where*

$$\gamma_G(A_{i,j}) := \int_{A_{i,j}} G(\beta_s, \beta_t) ds dt - \mathbb{E} \left[\int_{A_{i,j}} G(\beta_s, \beta_t) ds dt \right].$$

Moreover, there exists $\lambda \in \mathbb{R}$, such that

$$\mathbb{E}[e^{\lambda \cdot \gamma_G(\mathcal{C})}] = \infty.$$

REMARK 4.6. We take by definition $\gamma_G([0, 1]^2) := 2\gamma_G(\mathcal{C})$.

PROOF. For any $j \leq 2^{i-1}$, note that By Brownian scaling $\gamma_G(A_{i,j})$ has the same law as $2^{1-i}\gamma_G(A_{1,1})$. By Proposition 4.1 these random variables have finite second moment. Now by the triangle inequality for the L^2 -norm, we have for $m < n$

$$\left\| \sum_{i=m}^n \sum_{j=1}^{2^{i-1}} \gamma_G(A_{i,j}) \right\|_2 \leq \sum_{i=m}^n \left\| \sum_{j=1}^{2^{i-1}} \gamma_G(A_{i,j}) \right\|_2 = \sum_{i=m}^n \left(\sum_{j=1}^{2^{i-1}} \mathbb{E}[(\gamma_G(A_{i,j}))^2] \right)^{1/2},$$

where for the last inequality we used the independence of the terms $\gamma_G(A_{i,j})$, when the rectangles $A_{i,j}$ are all in the same subdivision. By scale invariance mentioned earlier, this last sum is equal to

$$\sum_{i=m}^n \sqrt{\frac{2^{i-1}}{2^{2(i-1)}} \cdot \text{Var}(\gamma_G(A_{1,1}))} = \sqrt{\text{Var}(\gamma_G(A_{1,1}))} \sum_{i=m}^n \frac{1}{2^{(i-1)/2}}.$$

This shows that the sequence $\sum_{i=1}^n \sum_{j=1}^{2^{i-1}} \gamma_G(A_{i,j})$ is Cauchy in L^2 , and hence it converges in L^2 . Now we prove that this limit random variable has some infinite exponential moment, using the same argument as in Le Gall [30]. First, note that by definition we can express $\gamma_G(\mathcal{C})$ almost surely as

$$\gamma_G(\mathcal{C}) = \frac{1}{2} \cdot \left(\gamma_1 + \gamma_2 + \frac{1}{2\pi^2} \cdot (X - \mathbb{E}[X]) \right),$$

with γ_1 and γ_2 two independent random variables with the same law as $\gamma_G(\mathcal{C})$, and X a random variable with the same law as the random variable from Proposition 4.1. Now the lower bound in the latter proposition shows that there exists $\lambda > 0$ such that $\mathbb{E}[\exp(\lambda X)] = \infty$. Since $X = 2\pi^2(2\gamma_G(\mathcal{C}) - \gamma_1 - \gamma_2) + \mathbb{E}[X]$, it implies by Cauchy–Schwarz that either $\mathbb{E}[\exp(8\pi^2\lambda\gamma_G(\mathcal{C}))]$ or $\mathbb{E}[\exp(-4\pi^2\lambda\gamma_G(\mathcal{C}))]$ is infinite. $\square$

5. Intersection and nonintersection probabilities. The goal of this section is to obtain an asymptotic expression for the probability of nonintersection of a two-sided walk with simple random walk, when one walk is conditioned to end up at a specific location. Our proofs will rely heavily on the following estimate of Lawler on the nonintersection probability of a two-sided walk with a simple random walk.

THEOREM 5.1 ([24], Corollary 4.2.5). *Let $\mathcal{R}^1, \mathcal{R}^2$ and $\mathcal{R}^3$ be the ranges of three independent random walks in $\mathbb{Z}^4$ starting at 0. Then*

$$(5.1) \quad \lim_{n \rightarrow \infty} \log n \cdot \mathbb{P}((\mathcal{R}^1[0, n] \cup \mathcal{R}^2[0, n]) \cap \mathcal{R}^3[1, \infty) = \emptyset, 0 \notin \mathcal{R}^1[1, n]) = \frac{\pi^2}{8}.$$

Recall the definition of $f_k(x)$ from (2.1) and also the shorthand notation $n_\alpha = n/(\log n)^\alpha$.

PROPOSITION 5.2. *Let $\alpha > 8$, $n - n_\alpha > k > n_\alpha$ and $x \in \mathbb{Z}^4$ with $\sqrt{n_{2\alpha}} \leq \|x\| \leq \sqrt{n}(\log n)^2$. Let $\mathcal{R}^1, \mathcal{R}^2$ and $\mathcal{R}^3$ be the ranges of three independent random walks starting from 0. Then*

$$\begin{aligned} \mathbb{P}((\mathcal{R}^1[0, k] \cup \mathcal{R}^2[0, n]) \cap \mathcal{R}^3[1, \infty) = \emptyset, S^1(k) = x, 0 \notin \mathcal{R}^1[1, k]) \\ = \frac{\pi^2}{8} \cdot \frac{1}{\log n} \cdot p_k(x) \cdot (1 + o(1)) + \mathcal{O}\left(\frac{\log \log n}{(\log n)^{3/2}} \cdot f_k(x/2)\right). \end{aligned}$$

Note that in the expression above we cannot always absorb the second term in the $o(1)$ term, since $p_k(x)$ is not always comparable to $f_k(x/2)$. However, this second term is going to be negligible when we take the sum over all time indices and all points in space.

The rest of this section is devoted to the proof of the above proposition.

The following lemma on the probability that two walks intersect when one walk is conditioned to end up at a specific location is a crucial ingredient in the proof of the proposition above and will also be used in later parts.

LEMMA 5.3. *There exists a positive constant C so that the following holds. Let $z \in \mathbb{Z}^4$ and let S^1 and S^2 be two independent simple random walks in $\mathbb{Z}^4$ starting from 0 and z , respectively. For $a, k, f \in \mathbb{N}$ and $b \in \mathbb{N} \cup \{\infty\}$ let $A(a, b, f, k) = \{\mathcal{R}^1[f, k] \cap \mathcal{R}^2[a, b] \neq \emptyset\}$. Then for all $x \in \mathbb{Z}^4$ with $\|x\| \leq k^{2/3}$ we have*

$$\begin{aligned} \mathbb{P}_{0,z}(A(a, b, f, k), S^1(k) = x) \\ \leq C f_k(x/2) \cdot \max(\mathbb{P}_{0,z}(A(a, b, f, k)), \mathbb{P}_{x,z}(A(a, b, 0, k))). \end{aligned}$$

We now state two lemmas and a claim whose proofs are deferred after the proof of Proposition 5.2.

LEMMA 5.4. *Let $\alpha > 8$. Let $n_\alpha < k < n - n_\alpha$, $i = k/(\log n)^{5\alpha}$ and $\sqrt{n_{2\alpha}} \leq \|x\| \leq \sqrt{n}(\log n)^2$. Let $\mathcal{R}^1, \mathcal{R}^2$ and $\mathcal{R}^3$ be the ranges of three independent random walks starting from 0. Then we have*

$$\begin{aligned} \mathbb{P}((\mathcal{R}^1[0, i] \cup \mathcal{R}^2[0, n]) \cap \mathcal{R}^3[1, \infty) = \emptyset, S^1(k) = x, 0 \notin \mathcal{R}^1[1, i]) \\ = \frac{\pi^2}{8} \cdot \frac{1}{\log n} \cdot p_k(x) \cdot (1 + o(1)). \end{aligned}$$

LEMMA 5.5. *Same assumptions as in Lemma 5.4. We have*

$$\begin{aligned} \mathbb{P}((\mathcal{R}^1[0, i] \cup \mathcal{R}^2[0, n]) \cap \mathcal{R}^3[1, \infty) = \emptyset, \mathcal{R}^1[i, k] \cap \mathcal{R}^3[1, \infty) \neq \emptyset, S^1(k) = x) \\ \lesssim f_k(x/2) \cdot \frac{\log \log n}{(\log n)^{3/2}}. \end{aligned}$$

CLAIM 5.6. *Let $\alpha > 0$, $k > n_\alpha$ and $\sqrt{n_{2\alpha}} \leq \|x\| \leq k^{3/5}$. Let $\text{hit}[a, b] = \{\exists \ell \in [a, b] : S^1(\ell) = 0\}$. Then we have*

$$\mathbb{P}(S^1(k) = x, \text{hit}[n_\alpha, k]) \lesssim f_k(x) \cdot \frac{(\log n)^{2\alpha}}{n}.$$

We now give the proof of Proposition 5.2 which is an easy consequence of the results above and then we will prove Lemmas 5.3, 5.4 and 5.5 and Claim 5.6.

PROOF OF PROPOSITION 5.2. Let $i = k/(\log n)^{5\alpha}$ as in Lemma 5.4. Then we can write

$$\begin{aligned} & \mathbb{P}((\mathcal{R}^1[0, k] \cup \mathcal{R}^2[0, n]) \cap \mathcal{R}^3[1, \infty) = \emptyset, S^1(k) = x, 0 \notin \mathcal{R}^1[1, k]) \\ &= \mathbb{P}((\mathcal{R}^1[0, i] \cup \mathcal{R}^2[0, n]) \cap \mathcal{R}^3[1, \infty) = \emptyset, S^1(k) = x, 0 \notin \mathcal{R}^1[1, i]) \\ &\quad - \mathbb{P}(\mathcal{R}^1[0, i] \cup \mathcal{R}^2[0, n]) \cap \mathcal{R}^3[1, \infty) = \emptyset, S^1(k) = x, 0 \notin \mathcal{R}^1[1, i], \\ &\quad \{\mathcal{R}^1[i, k] \cap \mathcal{R}^3[1, \infty) \neq \emptyset\} \cup \{0 \in \mathcal{R}^1[i, k]\}). \end{aligned}$$

For the first term, we use Lemma 5.4 to get the asymptotic expression of the statement. Regarding the second probability by the union bound it is upper bounded by

$$\begin{aligned} & \mathbb{P}((\mathcal{R}^1[0, i] \cup \mathcal{R}^2[0, n]) \cap \mathcal{R}^3[1, \infty) = \emptyset, S^1(k) = x, \mathcal{R}^1[i, k] \cap \mathcal{R}^3[1, \infty) \neq \emptyset) \\ &+ \mathbb{P}(S^1(k) = x, 0 \notin \mathcal{R}^1[1, i], 0 \in \mathcal{R}^1[i, k]). \end{aligned}$$

The first term can be bounded using Lemma 5.5 and the second one using Claim 5.6. $\square$

PROOF OF LEMMA 5.3. We assume that $p_k(x) > 0$, since otherwise the statement is trivial. Suppose first that $\|x\|^2 \leq k$. Then in this case $p_k(x) \asymp 1/k^2$, and hence for all $y, z \in \mathbb{Z}^4$ we have $p_{k/2}(y, z) \leq C/k^2 \asymp p_k(x)$. We now get

$$\begin{aligned} & \mathbb{P}_{0,z}(A(a, b, f, k), S^1(k) = x) \\ &= \sum_y \mathbb{P}_{0,z}(A(a, b, f, k), S^1(k) = x, S^1(k/2) = y) \\ &\leq \sum_y \mathbb{P}_{0,z}(\mathcal{R}^2[a, b] \cap \mathcal{R}^1[f, k/2] \neq \emptyset, S^1(k/2) = y, S^1(k) = x) \\ &\quad + \sum_y \mathbb{P}_{0,z}(\mathcal{R}^2[a, b] \cap \mathcal{R}^1[k/2, k] \neq \emptyset, S^1(k/2) = y, S^1(k) = x). \end{aligned}$$

Using time reversibility and the Markov property this last sum is equal to

$$\sum_y \mathbb{P}_{0,z}(A(a, b, f, k/2), S^1(k/2) = y) \cdot p_{k/2}(y, x)$$

$$\begin{aligned}
& + \sum_y \mathbb{P}_{x,z}(A(a, b, 0, k/2), S^1(k/2) = y) \cdot p_{k/2}(y, 0) \\
& \lesssim p_k(x) \cdot \sum_y \mathbb{P}_{0,z}(A(a, b, f, k/2), S^1(k/2) = y) \\
& \quad + p_k(x) \cdot \sum_y \mathbb{P}_{x,z}(A(a, b, 0, k/2), S^1(k/2) = y) \\
& \lesssim p_k(x) \cdot \mathbb{P}_{0,z}(A(a, b, f, k/2)) + p_k(x) \cdot \mathbb{P}_{x,z}(A(a, b, 0, k/2)) \\
& \lesssim p_k(x) \cdot \max(\mathbb{P}_{0,z}(A(a, b, f, k)), \mathbb{P}_{x,z}(A(a, b, 0, k))).
\end{aligned}$$

This completes the proof in the case when $\|x\|^2 \leq k$, since $f_k(x/2) \gtrsim 1/k^2$.

Suppose next that $\|x\|^2 \geq k$. We write $\mathcal{B}_x = \mathcal{B}(x, \|x\|/2)$ and $\mathcal{S}_x = \partial\mathcal{B}_x$. We define σ_x as the last visit time to $\mathcal{S}_x$ before time k and τ_x for the first hitting time of $\mathcal{S}_x$ (both for the walk S^1). For all $i \leq k$ and $w \in \mathbb{Z}^4$, we have

$$\begin{aligned}
(5.2) \quad & \mathbb{P}_w(S^1(i) = x, \tau_x \leq i) \\
& = \sum_{j \leq i} \sum_{y \in \mathcal{S}_x} \mathbb{P}_w(\tau_x = j, S^1(j) = y, S^1(i) = x) \\
& = \sum_{j \leq i} \sum_{y \in \mathcal{S}_x} \mathbb{P}_w(\tau_x = j, S^1(j) = y) \cdot p_{i-j}(y, x) \\
& \lesssim f_k(x/2) \cdot \mathbb{P}_w(\tau_x \leq i) \lesssim f_k(x/2),
\end{aligned}$$

where the first inequality follows from Claim 2.1. Now one can write

$$\begin{aligned}
(5.3) \quad & \mathbb{P}_{0,z}(A(a, b, f, k), S^1(k) = x) \\
& \leq \mathbb{P}_{0,z}(\mathcal{R}^2[a, b] \cap \mathcal{R}^1[f, \sigma_x] \neq \emptyset, S^1(k) = x) \\
& \quad + \mathbb{P}_{0,z}(\mathcal{R}^2[a, b] \cap \mathcal{R}^1[\sigma_x, k] \neq \emptyset, S^1(k) = x).
\end{aligned}$$

In order to bound the first term, let us define

$$\mathcal{I} := \inf\{i \geq f : S^1(i) \in \mathcal{R}^2[a, b]\}.$$

Note that for any i , the event $\{\mathcal{I} = i\}$ is $\sigma(\mathcal{R}^1[f, i]) \vee \sigma(\mathcal{R}^2[a, b])$ -measurable. Therefore, by the Markov property we obtain

$$\begin{aligned}
& \mathbb{P}_{0,z}(\mathcal{R}^2[a, b] \cap \mathcal{R}^1[f, \sigma_x] \neq \emptyset, S^1(k) = x) \\
& = \sum_w \sum_{f \leq i \leq k} \mathbb{P}_{0,z}(\mathcal{I} = i, \sigma_x \geq i, S^1(i) = w, S^1(k) = x) \\
& = \sum_{w, f \leq i \leq k} \mathbb{P}_{0,z}(\mathcal{I} = i, S^1(i) = w) \cdot \mathbb{P}_w(S^1(k-i) = x, \tau_x \leq k-i) \\
& \lesssim f_k(x/2) \cdot \sum_{w, f \leq i \leq k} \mathbb{P}_{0,z}(\mathcal{I} = i, S^1(i) = w)
\end{aligned}$$

$$\lesssim f_k(x/2) \cdot \mathbb{P}_{0,z}(A(a, b, f, k)),$$

where we used (5.2) for the first inequality. Now concerning the second term in (5.3), one can look at the path backwards in time, and observe that seen from x , σ_x is now the first hitting time of $\mathcal{S}_x$, namely τ_x . Therefore, using the strong Markov property,

$$\begin{aligned} & \mathbb{P}_{0,z}(\mathcal{R}^2[a, b] \cap \mathcal{R}^1[\sigma_x, k] \neq \emptyset, S^1(k) = x) \\ &= \sum_{y \in \mathcal{S}_x, i \leq k} \mathbb{P}_{0,z}(\mathcal{R}^2[a, b] \cap \mathcal{R}^1[i, k] \neq \emptyset, S^1(i) = y, \sigma_x = i, S^1(k) = x) \\ &= \sum_{y, i} \mathbb{P}_{x,z}(A(a, b, 0, k-i), S^1(k-i) = y, \tau_x = k-i) \cdot p_i(y) \\ &\lesssim f_k(x/2) \sum_{y, i} \mathbb{P}_{x,z}(A(a, b, 0, k), S^1(k-i) = y, \tau_x = k-i) \\ &\lesssim f_k(x/2) \cdot \mathbb{P}_{x,z}(A(a, b, 0, k)), \end{aligned}$$

where for the first inequality we used Claim 2.1 again. This now completes the proof. $\square$

PROOF OF LEMMA 5.4. This proof is very similar to [24], Proposition 4.3.2, but we include it here for the sake of completeness. Again we assume that $p_k(x) > 0$, otherwise the statement is trivial.

We define $D = \{\|S^1(i)\| \leq \sqrt{i}(\log n)^{\alpha+6}\}$. Then $\mathbb{P}(D^c) \leq \exp(-(\log n)^{\alpha+6})$. We set

$$A = \{(\mathcal{R}^1[0, i] \cup \mathcal{R}^2[0, n]) \cap \mathcal{R}^3[1, \infty) = \emptyset, 0 \notin \mathcal{R}^1[1, i]\}.$$

Then by Lawler's estimate, Theorem 5.1, we have that

$$\mathbb{P}(A) = \frac{\pi^2}{8} \cdot \frac{1}{\log n} \cdot (1 + o(1)).$$

We now obtain

$$\mathbb{P}(A, S^1(k) = x, D) \leq \mathbb{P}(A, S^1(k) = x) \leq \mathbb{P}(A, S^1(k) = x, D) + \mathbb{P}(D^c).$$

By the Markov property, we now get

$$\begin{aligned} \mathbb{P}(A \cap D, S^1(k) = x) &= \mathbb{P}(S^1(k) = x | A \cap D) \mathbb{P}(A \cap D) \\ &= p_k(x)(1 + o(1)) \mathbb{P}(A \cap D), \end{aligned}$$

where the last equality follows from (2.3) and Claim 2.2, since after conditioning on D , the time changes to $k-i$ and the walk starts from some z with $\|z\| \leq \sqrt{i}(\log n)^{\alpha+6}$. Note that for $\alpha > 8$ the assumptions of Claim 2.2 are satisfied. We also have

$$\mathbb{P}(A \cap D) = \mathbb{P}(A) - \mathbb{P}(A \cap D^c) = \mathbb{P}(A)(1 + o(1)),$$

since $\mathbb{P}(A \cap D^c) \leq \mathbb{P}(D^c) \leq \exp(-(\log n)^{\alpha+6})$ and $\mathbb{P}(A) \asymp 1/\log n$. So far we have showed that

$$\mathbb{P}(A \cap D, S^1(k) = x) = p_k(x)(1 + o(1))\mathbb{P}(A).$$

By the assumption on the values of x and k , we get that

$$\mathbb{P}(A)p_k(x) \gtrsim \exp(-c(\log n)^{\alpha+4}).$$

Therefore,

$$\mathbb{P}(A \cap D, S^1(k) = x) + \mathbb{P}(D^c) = p_k(x)\mathbb{P}(A)(1 + o(1))$$

and this completes the proof. $\square$

PROOF OF LEMMA 5.5. Define $\ell = i/(\log n)^{2\alpha+20}$. Then we can upper bound the probability of the statement as follows (with $\mathcal{R}_+$ for $\mathcal{R}[1, \infty)$)

$$\begin{aligned} (5.4) \quad & \mathbb{P}((\mathcal{R}^1[0, i] \cup \mathcal{R}^2[0, n]) \cap \mathcal{R}_+^3 = \emptyset, \mathcal{R}^1[i, k] \cap \mathcal{R}_+^3 \neq \emptyset, S^1(k) = x) \\ & \leq \mathbb{P}(\mathcal{R}^2[0, n] \cap \mathcal{R}^3[1, \ell] = \emptyset, \mathcal{R}^1[i, k] \cap \mathcal{R}_+^3 \neq \emptyset, S^1(k) = x) \\ & \leq \mathbb{P}(\mathcal{R}^2[0, n] \cap \mathcal{R}^3[1, \ell] = \emptyset, \mathcal{R}^1[i, k] \cap \mathcal{R}^3[1, \ell] \neq \emptyset, S^1(k) = x) \\ & \quad + \mathbb{P}(\mathcal{R}^2[0, n] \cap \mathcal{R}^3[1, \ell] = \emptyset, \mathcal{R}^1[i, k] \cap \mathcal{R}^3[\ell, \infty) \neq \emptyset, S^1(k) = x). \end{aligned}$$

From Lemma 5.3, we have

$$\begin{aligned} & \mathbb{P}(\mathcal{R}^1[i, k] \cap \mathcal{R}^3[1, \ell] \neq \emptyset, S^1(k) = x) \\ & \lesssim f_k(x/2) \cdot \max(\mathbb{P}_{x,0}(\mathcal{R}^1[0, k] \cap \mathcal{R}^3[1, \ell] \neq \emptyset), \\ & \quad \mathbb{P}(\mathcal{R}^1[i, k] \cap \mathcal{R}^3[1, \ell] \neq \emptyset)). \end{aligned}$$

We now define

$$D_3 = \left\{ \max_{j \leq \ell} \|S^3(j)\| \leq \frac{\sqrt{i}}{(\log n)^4} \right\}.$$

By the choice of ℓ and (2.2), we have $\mathbb{P}(D_3^c) \lesssim \exp(-(\log n)^{\alpha+6})$. We also let

$$D_1 = \left\{ \|S^1(i)\| \geq \frac{\sqrt{i}}{(\log n)^2} \right\}.$$

Then by (2.4) we get $\mathbb{P}(D_1^c) \lesssim \frac{1}{(\log n)^8}$, and hence we obtain that

$$\begin{aligned} & \mathbb{P}(\mathcal{R}^1[i, k] \cap \mathcal{R}^3[1, \ell] \neq \emptyset) \\ & \leq \mathbb{P}(D_3^c) + \mathbb{P}(D_1^c) + \mathbb{P}(\mathcal{R}^1[i, k] \cap \mathcal{R}^3[1, \ell] \neq \emptyset, D_1, D_3) \\ & \lesssim \frac{2}{(\log n)^8} + \max_{\|w\| \geq \sqrt{i}/(\log n)^2} \mathbb{P}_w(H_{B(0, \frac{\sqrt{i}}{(\log n)^4})} < \infty) \lesssim \frac{1}{(\log n)^4}, \end{aligned}$$

where for the last inequality we used (2.7). When S^1 starts from x , we then get

$$\begin{aligned} \mathbb{P}_{x,0}(\mathcal{R}^1[0, k] \cap \mathcal{R}^3[1, \ell] \neq \emptyset) &\lesssim \mathbb{P}(D_3^c) + \mathbb{P}_{x,0}(\mathcal{R}^1[0, k] \cap \mathcal{R}^3[1, \ell] \neq \emptyset, D_3) \\ &\leq \exp(-(\log n)^{\alpha+6}) + \mathbb{P}_x(H_{\mathcal{B}(0, \frac{\sqrt{i}}{(\log n)^4})} < \infty) \\ &\leq \exp(-(\log n)^{\alpha+6}) + \frac{1}{(\log n)^{8+3\alpha}}, \end{aligned}$$

where for the last inequality we used again (2.7) and the assumption on i and $\|x\|$.

So we overall get that

$$\mathbb{P}(\mathcal{R}^2[0, n] \cap \mathcal{R}^3[1, \ell] = \emptyset, \mathcal{R}^1[i, k] \cap \mathcal{R}^3[1, \ell] \neq \emptyset, S^1(k) = x) \lesssim \frac{f_k(\frac{x}{2})}{(\log n)^4}.$$

Returning to the probability appearing in the penultimate line of (5.4), we obtain

$$\begin{aligned} &\mathbb{P}(\mathcal{R}^2[0, n] \cap \mathcal{R}^3[1, \ell] = \emptyset, \mathcal{R}^1[i, k] \cap \mathcal{R}^3[\ell, \infty) \neq \emptyset, S^1(k) = x) \\ &\leq \mathbb{P}(\mathcal{R}^2[0, n] \cap \mathcal{R}^3[1, \ell] = \emptyset, \mathcal{R}^1[i, k] \cap \mathcal{R}^3[\ell, \infty) \neq \emptyset, S^1(k) = x, D_3) \\ (5.5) \quad &+ \mathbb{P}(D_3^c) \\ &\lesssim \mathbb{P}(\mathcal{R}^2[0, n] \cap \mathcal{R}^3[1, \ell] = \emptyset, \mathcal{R}^1[i, k] \cap \mathcal{R}^3[\ell, \infty) \neq \emptyset, S^1(k) = x, D_3) \\ &+ \frac{f_k(\frac{x}{2})}{(\log n)^4}, \end{aligned}$$

where for the last inequality we used that for x and k as in the statement of the lemma we have

$$p_k(x) \gtrsim \exp(-2(\log n)^{\alpha+4}) \quad \text{and} \quad \mathbb{P}(D_3^c) \leq \exp(-(\log n)^{\alpha+6}).$$

The first term appearing in the last line of (5.5) is upper bounded by

$$\begin{aligned} &\sum_{\|z\|^2 \leq \frac{i}{\log^8 n}} \mathbb{P}_{0,z}(\mathcal{R}^1[i, k] \cap \mathcal{R}_+^3 \neq \emptyset, S^1(k) = x) \\ (5.6) \quad &\times \mathbb{P}(\mathcal{R}^2[0, n] \cap \mathcal{R}^3[1, \ell] = \emptyset, S^3(\ell) = z). \end{aligned}$$

Using Lemma 5.3 again and [24], Theorem 4.3.3, we get that for all z in the range as above

$$\begin{aligned} &\mathbb{P}_{0,z}(\mathcal{R}^1[i, k] \cap \mathcal{R}_+^3 \neq \emptyset, S^1(k) = x) \\ &\lesssim f_k(x/2) \max(\mathbb{P}_{0,z}(\mathcal{R}^1[i, k] \cap \mathcal{R}_+^3 \neq \emptyset), \mathbb{P}_{x,z}(\mathcal{R}^1[0, k] \cap \mathcal{R}_+^3 \neq \emptyset)) \\ &\lesssim f_k(x/2) \cdot \frac{\log \log n}{\log n}. \end{aligned}$$

Therefore, the sum of (5.6) becomes upper bounded by

$$\mathbb{P}(\mathcal{R}^2[0, n] \cap \mathcal{R}^3[1, \ell] = \emptyset) \cdot f_k(x/2) \cdot \frac{\log \log n}{\log n} \asymp f_k(x/2) \cdot \frac{\log \log n}{(\log n)^{3/2}},$$

where for the equivalence we used [24], Theorem 4.4.1. Substituting this into (5.5) completes the proof. $\square$

PROOF OF CLAIM 5.6. In order to upper bound the probability of this event, we consider two cases, either $\|x\| \leq \sqrt{k}$ or $\|x\| > \sqrt{k}$. If $\|x\| \leq \sqrt{k}$, then using reversibility and the Markov property we obtain

$$\begin{aligned} \mathbb{P}(S^1(k) = x, \text{hit}[n_\alpha, k]) &= \mathbb{P}(S^1(k) = x, \text{hit}[n_\alpha, k/2]) + \mathbb{P}(S^1(k) = x, \text{hit}[k/2, k]) \\ &= \sum_z \mathbb{P}(S^1(k/2) = z, S^1(k) = x, \text{hit}[n_\alpha, k/2]) \\ &\quad + \sum_z \mathbb{P}(S^1(k/2) = z, S^1(k) = x, \text{hit}[k/2, k]) \\ &\lesssim p_k(x) \mathbb{P}(\text{hit}[n_\alpha, k/2]) + p_k(x) \mathbb{P}_x(H_0 < \infty) \\ &\lesssim p_k(x) \cdot \frac{(\log n)^{2\alpha}}{n}. \end{aligned}$$

We turn to the case $\|x\| > \sqrt{k}$. We now have using the Markov property

$$\begin{aligned} \mathbb{P}(S^1(k) = x, \text{hit}[n_\alpha, k]) &= \mathbb{P}(S^1(k) = x | \text{hit}[n_\alpha, k]) \mathbb{P}(\text{hit}[n_\alpha, k]) \\ &\leq \sup_{i < k} p_i(x) \cdot \mathbb{P}(\text{hit}[n_\alpha, k]) \lesssim f_k(x) \cdot \frac{(\log n)^{2\alpha}}{n}, \end{aligned}$$

where for the last inequality we used Claim 2.1. $\square$

6. Joint convergence in law of the cross terms. The main purpose of this section is to prove the joint convergence in law of the cross terms. First, recall the definition of the squares

$$A_{i,j} = [(2j-2)2^{-i}, (2j-1)2^{-i}] \times [(2j-1)2^{-i}, (2j)2^{-i}],$$

for $i \geq 1$ and $j \leq 2^{i-1}$. Recall also the definition of the cross terms from Proposition 2.3:

$$\chi_n(i, j) = \chi(\mathcal{R}_n^{(i, 2j-1)}, \mathcal{R}_n^{(i, 2j)}) + \chi(\mathcal{R}_n^{(i, 2j)}, \mathcal{R}_n^{(i, 2j-1)}),$$

with

$$(6.1) \quad \mathcal{R}_n^{(i,j)} = \mathcal{R}[(j-1)2^{-i}n, j2^{-i}n].$$

PROPOSITION 6.1. *Let $p \geq 1$ be a fixed integer. Then as $n \rightarrow \infty$,*

$$(6.2) \quad \left(\frac{2(\log n)^2}{\pi^4 \cdot n} \cdot \chi_n(i, j) \right)_{\substack{1 \leq i \leq p \\ 1 \leq j \leq 2^{i-1}}} \xrightarrow{(d)} \left(\int_{A_{i,j}} G(\beta_s, \beta_t) ds dt \right)_{\substack{1 \leq i \leq p \\ 1 \leq j \leq 2^{i-1}}}.$$

Moreover, the convergence also holds in expectation.

Our strategy for proving this proposition is first to localize in a certain sense all the $\chi_n(i, j)$. More precisely, we prove that for any given i and $j \leq 2^{i-1}$, the term $\chi_n(i, j)$ can be written as a sum of two elements, one being a localised version of this cross term [the so-called $\chi_{n,\alpha}(i, j)$, see below], and the other one having a negligible expectation. So to prove the joint convergence in law of the cross terms, we are led to prove only the joint convergence in law of the $\chi_{n,\alpha}(i, j)$. To prove this in turn, we show the convergence of the joint moments, which is indeed sufficient thanks to the results of Section 4.1 and Carleman's criterion (see Section 3.3.5 in [12]).

Now the $\chi_{n,\alpha}(i, j)$ have the great advantage that their joint moments reduce (after some tedious computation) to a product of nonintersection probabilities (whose asymptotics have been computed in the previous section) times a product of Green's function. Then a separate argument shows that this product of Green's functions converges to its continuous counterpart.

Before digging into the proof, we gather some basic preliminary estimates in the next subsection.

6.1. *Preliminaries.* We start with an elementary fact which directly follows from the local CLT (2.3) and (2.2): there is a constant $C > 0$, such that for all $k \geq 0$,

$$(6.3) \quad \mathbb{E}[G_d(S_k)] \leq C \cdot \frac{1}{k+1} \quad \text{and} \quad \mathbb{E}[G_d(S_k)^2] \leq C \cdot \frac{1}{k^2+1}.$$

Now for $\alpha > 0$, recall that $n_\alpha = n/(\log n)^\alpha$, and define the set

$$(6.4) \quad B_\alpha = \{(x, y) : \sqrt{n_{2\alpha}} \leq \|x\|, \|y\| \leq \sqrt{n}(\log n)^2, \|x - y\| \geq \sqrt{n_{2\alpha}}\}.$$

LEMMA 6.2. *Let S and $\tilde{S}$ be two independent random walks and let $\alpha > 0$. Then*

$$(6.5) \quad \sum_{k=0}^n \sum_{\ell=0}^{n_\alpha} \mathbb{E}[G_d(S_k - \tilde{S}_\ell)] \lesssim n_\alpha \cdot \log n,$$

$$(6.6) \quad \sum_{k=0}^n \sum_{\ell=n-n_\alpha}^n \mathbb{E}[G_d(S_k - \tilde{S}_\ell)] \lesssim n_\alpha$$

and

$$(6.7) \quad \sum_{k,\ell=0}^n \sum_{(x,y) \notin B_\alpha} \mathbb{P}(S_k = x, \tilde{S}_\ell = y) \cdot G_d(x, y) \lesssim n_{2\alpha} \cdot (\log n)^2.$$

PROOF. Note that $S_k - \tilde{S}_\ell$ is equal in law to $S_{k+\ell}$. Thus by using (6.3), we deduce that for any $k \geq 1$,

$$\sum_{\ell=0}^{n_\alpha} \mathbb{E}[G_d(S_{k+\ell})] \lesssim \sum_{\ell=0}^{n_\alpha} \frac{1}{k+\ell} \lesssim \frac{n_\alpha}{k}.$$

Summing over k proves (6.5). The proof of (6.6) is entirely similar.

We prove now (6.7). Using (2.4) yields

$$\mathbb{P}(\|S_k - \tilde{S}_\ell\|^2 \leq n_{2\alpha}) = \mathbb{P}(\|S_{k+\ell}\|^2 \leq n_{2\alpha}) \lesssim \frac{n_{2\alpha}^2}{1 + (k + \ell)^2}.$$

Similarly, one has

$$\mathbb{P}(\|S_k\|^2 \leq n_{2\alpha}) \lesssim \frac{n_{2\alpha}^2}{1 + k^2} \quad \text{and} \quad \mathbb{P}(\|\tilde{S}_\ell\|^2 \leq n_{2\alpha}) \lesssim \frac{n_{2\alpha}^2}{1 + \ell^2}.$$

Therefore, by using (6.3), (2.2), (2.4) and Cauchy–Schwarz, we get

$$\mathbb{E}[G_d(S_k, \tilde{S}_\ell) \mathbb{1}((S_k, \tilde{S}_\ell) \notin B_\alpha)] \lesssim \frac{n_{2\alpha}}{1 + k + \ell} \left(\frac{1}{1 + k} + \frac{1}{1 + \ell} \right).$$

The result follows by summing over k and ℓ . $\square$

LEMMA 6.3. *There exists a constant $C > 0$ so that for all $i \geq 1$ and for all $j \leq 2^{i-1}$, one has*

$$\frac{1}{n} \cdot \mathbb{E} \left[\sum_{k, \ell \leq n} \mathbb{1}(\|S_k - S_\ell\| \leq \varepsilon \sqrt{n}) \cdot G_d(S_k, S_\ell) \right] \leq C \cdot \varepsilon \log \frac{1}{\varepsilon}.$$

PROOF. By considering two independent random walks, we get

$$\begin{aligned} & \frac{1}{n} \cdot \mathbb{E} \left[\sum_{k, \ell \leq n} \mathbb{1}_{\|S_k - S_\ell\| \leq \varepsilon \sqrt{n}} \cdot G_d(S_k, S_\ell) \right] \\ (6.8) \quad & \lesssim \frac{1}{n} \sum_{k=1}^{2n} k \sum_{z: \|z\| \leq \varepsilon \sqrt{n}} p_k(z) G_d(z) \\ & \lesssim \frac{1}{n} \sum_{k=1}^{2n} \sum_{z: \|z\| \leq \varepsilon \sqrt{n}} k \times \frac{\exp(-\|z\|^2/(2k))}{k^2} G_d(z). \end{aligned}$$

Summing over z , we now obtain

$$\begin{aligned} \sum_{z: \|z\| \leq \varepsilon \sqrt{n}} \exp(-\|z\|^2/(2k)) G_d(z) & \lesssim \int_0^{\varepsilon \sqrt{n}} \frac{\exp(-r^2/(2k))}{r^2} r^3 dr \\ & = k(1 - \exp(-\varepsilon^2 n/(2k))). \end{aligned}$$

Summing over k , we get

$$\frac{1}{n} \sum_{k=1}^{2n} (1 - \exp(-\varepsilon^2 n/2k)) \leq \varepsilon + \frac{1}{n} \sum_{k=\varepsilon n}^{2n} \varepsilon \frac{n}{2k} \lesssim \varepsilon \log(1/\varepsilon)$$

and this concludes the proof. $\square$

LEMMA 6.4. *We have*

$$\sum_{x,y \in \mathbb{Z}^4} \sum_{k \leq n} \sum_{\ell \leq n} f_k(x/2) f_\ell(y/2) G_d(x, y) \lesssim n.$$

PROOF. The proof follows immediately by substituting the expression for f and using (2.6). $\square$

6.2. *Localisation of one cross term.* For any $n \geq 1, i \geq 1$ and $j \leq 2^{i-1}$, define

$$A_{i,j}^n := \{(2j-2)2^{-i}n, \dots, (2j-1)2^{-i}n\} \times \{(2j-1)2^{-i}n, \dots, j2^{-i+1}n\},$$

and for any $\alpha > 0$,

$$\begin{aligned} A_{i,j}^{n,\alpha} &:= \{(2j-2)2^{-i}n + n_\alpha, \dots, (2j-1)2^{-i}n - n_\alpha\} \\ &\quad \times \{(2j-1)2^{-i}n + n_\alpha, \dots, j2^{-i+1}n - n_\alpha\} \end{aligned}$$

at least for n large enough, to make sense of this definition, and with the usual convention to take integer parts when needed.

LEMMA 6.5. *Let i, j be positive integers with $j \leq 2^{i-1}$. For all $\alpha > 3$, we have*

$$\chi_n(i, j) = 2 \cdot \chi_{n,\alpha}(i, j) + \varepsilon_{n,\alpha}(i, j),$$

where $\mathbb{E}[\varepsilon_{n,\alpha}(i, j)] = o(n/(\log n)^2)$ and

$$\begin{aligned} \chi_{n,\alpha}(i, j) &= \sum_{\substack{(k,\ell) \in A_{i,j}^{n,\alpha} \\ (x,y) \in B_\alpha}} \mathbb{P}(\mathcal{R}[k - n_{4\alpha}, k + n_{4\alpha}] \cap (x + \mathcal{R}^1) = \emptyset, \\ &\quad S_k = x \notin \mathcal{R}[k - n_{4\alpha}, k] | S) \\ &\quad \times G_d(x, y) \cdot \mathbb{P}(\mathcal{R}[\ell - n_{4\alpha}, \ell + n_{4\alpha}] \cap (y + \mathcal{R}^2) = \emptyset, \\ &\quad S_\ell = y \notin \mathcal{R}[\ell - n_{4\alpha}, \ell] | S), \end{aligned}$$

with B_α as in (6.4) and $\mathcal{R}^1$ and $\mathcal{R}^2$ the ranges of two independent random walks starting from 0 and where for simplicity we used the convention $\mathcal{R}^1 = \mathcal{R}^1[1, \infty)$ and similarly for $\mathcal{R}^2$.

In order to prove the lemma above, we first approximate $\chi_n(i, j)$ by an expression without localisation which we call $\tilde{\chi}_{n,\alpha}(i, j)$, and which decorrelates the two parts of the range in some sense.

LEMMA 6.6. *With the same notation as in Lemma 6.5 and (6.1), we have*

$$\chi_n(i, j) = 2 \cdot \tilde{\chi}_{n,\alpha}(i, j) + \tilde{\varepsilon}_{n,\alpha}(i, j),$$

where $\mathbb{E}[\tilde{\varepsilon}_{n,\alpha}(i, j)] = o(n/(\log n)^2)$ and

$$\begin{aligned} \tilde{\chi}_{n,\alpha}(i, j) &= \sum_{\substack{(k,\ell) \in A_{i,j}^{n,\alpha} \\ (x,y) \in B_\alpha}} \mathbb{P}(\mathcal{R}_n^{(i,2j)} \cap (y + \mathcal{R}^2) = \emptyset, \\ &\quad S_\ell = y \notin \mathcal{R}[(2j-1)2^{-i}n, \ell] | S) \\ &\quad \times G_d(x, y) \cdot \mathbb{P}(\mathcal{R}_n^{(i,2j-1)} \cap (x + \mathcal{R}^1) = \emptyset, \\ &\quad S_k = x \notin \mathcal{R}[(2j-2)2^{-i}n, k] | S). \end{aligned}$$

PROOF OF LEMMA 6.6. Since i and j are going to be kept fixed while n will tend to infinity, we will not lose generality in doing the proof for $i = 0$ and $j = 1$. Also by moving the origin to $S(n)$, and looking at the range $\mathcal{R}[0, n]$ backwards, one is led to consider $\chi(\mathcal{R}_n, \tilde{\mathcal{R}}_n) + \chi(\tilde{\mathcal{R}}_n, \mathcal{R}_n)$, with $\mathcal{R}_n$ and $\tilde{\mathcal{R}}_n$ two independent ranges. So it suffices to treat the term $\chi(\mathcal{R}_n, \tilde{\mathcal{R}}_n)$.

By the independence of $\mathcal{R}^1$ and $\mathcal{R}^2$, we get

$$\begin{aligned} \chi(\mathcal{R}_n, \tilde{\mathcal{R}}_n) &= \sum_{k,\ell=0}^n \sum_{x,y} G_d(x, y) \mathbb{P}((\mathcal{R}_n \cup \tilde{\mathcal{R}}_n) \cap (x + \mathcal{R}^1) = \emptyset, \\ &\quad S_k = x \notin \mathcal{R}[0, k] | S, \tilde{S}) \\ &\quad \times \mathbb{P}(\tilde{\mathcal{R}}_n \cap (y + \mathcal{R}^2) = \emptyset, \\ &\quad \tilde{S}_\ell = y \notin \tilde{\mathcal{R}}[0, \ell] | \tilde{S}). \end{aligned}$$

Lemma 6.2 shows that we can restrict the sum over $n_\alpha \leq k, \ell \leq n - n_\alpha$ and $(x, y) \in B_\alpha$ at a cost of at most $n_\alpha \log n$ in expectation. The probability term appearing above is equal to

$$\begin{aligned} &\mathbb{P}(\mathcal{R}_n \cap (x + \mathcal{R}^1) = \emptyset, S_k = x \notin \mathcal{R}[0, k] | S) \\ &\quad \times \mathbb{P}(\tilde{\mathcal{R}}_n \cap (y + \mathcal{R}^2) = \emptyset, \tilde{S}_\ell = y \notin \tilde{\mathcal{R}}[0, \ell] | \tilde{S}) \\ &\quad - \mathbb{P}(\mathcal{R}_n \cap (x + \mathcal{R}^1) = \emptyset, \tilde{\mathcal{R}}_n \cap (y + \mathcal{R}^2) = \emptyset, \tilde{\mathcal{R}}_n \cap (x + \mathcal{R}^1) \neq \emptyset, \\ &\quad S_k = x \notin \mathcal{R}[0, k], \tilde{S}_\ell = y \notin \tilde{\mathcal{R}}[0, \ell] | S, \tilde{S}). \end{aligned}$$

The first term is equal to the probability term in the expression of $\tilde{\chi}_{n,\alpha}$. So we now turn to the second term. On the event $\{S_k = x\}$, by moving the origin to point x and reversing time we can write

$$\{\mathcal{R}_n \cap (x + \mathcal{R}^1) = \emptyset\} = \{(\mathcal{R}^3[0, k] \cup \mathcal{R}^4[0, n - k]) \cap \mathcal{R}^1 = \emptyset\},$$

where $\mathcal{R}^3$ and $\mathcal{R}^4$ are the ranges of two independent walks starting from 0. Applying the same to $\tilde{\mathcal{R}}$, we get that the event under consideration is contained in the intersection of the following events:

$$(6.9) \quad \begin{aligned} & \{(\mathcal{R}^3[0, k] \cup \mathcal{R}^4[0, n - k]) \cap \mathcal{R}^1 = \emptyset, S^3(k) = -x\}, \\ & \{(\tilde{\mathcal{R}}^3[0, \ell] \cup \tilde{\mathcal{R}}^4[0, n - \ell]) \cap \mathcal{R}^2 = \emptyset, \tilde{S}^3(\ell) = -y\}, \\ & \{(\tilde{\mathcal{R}}^3[0, \ell] \cup \tilde{\mathcal{R}}^4[0, n - \ell]) \cap (x - y + \mathcal{R}^1) \neq \emptyset, \tilde{S}^3(\ell) = -y\}. \end{aligned}$$

Setting $i = n_\beta$ with $\beta = 10\alpha + 4$, we next define

$$\begin{aligned} A_1 &= \{(\mathcal{R}^3[0, k] \cup \mathcal{R}^4[0, n - k]) \cap \mathcal{R}^1[0, i] = \emptyset, S^3(k) = -x\}, \\ A_2 &= \{\tilde{\mathcal{R}}^3[0, \ell] \cap \mathcal{R}^2 = \emptyset, \tilde{S}^3(\ell) = -y\}, \\ A_3 &= \{\tilde{\mathcal{R}}^4[0, n - \ell] \cap \mathcal{R}^2 = \emptyset\}, \\ A_4 &= \{\tilde{\mathcal{R}}^3[0, \ell] \cap (x - y + \mathcal{R}^1[0, i]) \neq \emptyset, \tilde{S}^3(\ell) = -y\}, \\ A_5 &= \{\tilde{\mathcal{R}}^3[0, \ell] \cap (x - y + \mathcal{R}^1[i, \infty)) \neq \emptyset, \tilde{S}^3(\ell) = -y\}, \\ A_6 &= \{\tilde{\mathcal{R}}^4[0, n - \ell] \cap (x - y + \mathcal{R}^1[0, i]) \neq \emptyset\}, \\ A_7 &= \{\tilde{\mathcal{R}}^4[0, n - \ell] \cap (x - y + \mathcal{R}^1[i, \infty)) \neq \emptyset\}. \end{aligned}$$

The first event in (6.9) is contained in A_1 ; the second one is contained in $A_2 \cap A_3$ and the third one is contained in the union of A_4, A_5, A_6 and A_7 . Therefore, we get that the probability of the intersection of the events appearing in (6.9) is upper bounded by

$$(6.10) \quad \mathbb{P}(A_1, A_2, A_6) + \mathbb{P}(A_1, A_2, A_7) + \mathbb{P}(A_1, A_3, A_4) + \mathbb{P}(A_1, A_3, A_5).$$

The first step now is to decorrelate the events where the range $\mathcal{R}^1$ appears. Lemma 5.3 gives

$$(6.11) \quad \begin{aligned} & \mathbb{P}(A_4) = \mathbb{P}(\tilde{\mathcal{R}}^3[0, \ell] \cap (x - y + \mathcal{R}^1[0, i]) \neq \emptyset, \tilde{S}^3(\ell) = -y) \\ & \lesssim f_\ell(y/2) \times \max(\mathbb{P}(\tilde{\mathcal{R}}^3[0, \ell] \cap (x - y + \mathcal{R}^1[0, i]) \neq \emptyset), \\ & \quad \mathbb{P}_{-y,0}(\tilde{\mathcal{R}}^3[0, \ell] \cap (x - y + \mathcal{R}^1[0, i]) \neq \emptyset)). \end{aligned}$$

Defining the event $D = \{\|S^1(r)\| \leq \sqrt{i}(\log n)^{\alpha+2}, \forall r \leq i\}$ we get from (2.2) for a positive constant c

$$(6.12) \quad \mathbb{P}(D^c) \lesssim \exp(-c(\log n)^{2\alpha+4}).$$

On the event D , in order for $\tilde{\mathcal{R}}^3[0, \ell]$ and $x - y + \mathcal{R}^1[0, i]$ to intersect, $\tilde{S}^3$ must hit a ball centred at $x - y$ of radius $\sqrt{i}(\log n)^{\alpha+2}$ or a ball centred at x of the same radius (depending on whether we start from 0 or $-y$). Since $\|x - y\| \geq \sqrt{n_{2\alpha}}$ and $\|x\| \geq \sqrt{n_{2\alpha}}$, using (2.7) and (6.12) we get

$$(6.13) \quad \begin{aligned} & \mathbb{P}(\tilde{\mathcal{R}}^3[0, \ell] \cap (x - y + \mathcal{R}^1[0, i]) \neq \emptyset) \\ & \leq \mathbb{P}_{-y, 0}(\tilde{\mathcal{R}}^3[0, \ell] \cap (x - y + \mathcal{R}^1[0, i]) \neq \emptyset) \\ & \lesssim \frac{1}{(\log n)^4}, \end{aligned}$$

where the last inequality follows from the choice of β (recall that $i = n_\beta$ with $\beta = 10\alpha + 4$).

Using (6.11), (6.13) and the independence between $\tilde{\mathcal{R}}^3$ and $\tilde{\mathcal{R}}^4$, we get

$$\mathbb{P}(A_1, A_3, A_4) \lesssim \frac{1}{(\log n)^4} \cdot p_k(x) f_\ell(y/2) \cdot \mathbb{P}(A_3) \leq \frac{1}{(\log n)^4} \cdot p_k(x) f_\ell(y/2).$$

Similarly, we get the same upper bound for $\mathbb{P}(A_1, A_2, A_6)$. It remains to bound the probabilities $\mathbb{P}(A_1, A_2, A_7)$ and $\mathbb{P}(A_1, A_3, A_5)$. By the independence between the walks again, we get

$$\mathbb{P}(A_1, A_2, A_7) = \mathbb{P}(A_2)\mathbb{P}(A_1, A_7) \quad \text{and} \quad \mathbb{P}(A_1, A_3, A_5) = \mathbb{P}(A_3)\mathbb{P}(A_1, A_5).$$

For the probability of the event A_2 by exactly the same proof as in Lemma 5.4, we have for a suitable $\gamma > 0$,

$$\mathbb{P}(A_2) \leq \mathbb{P}\left(\tilde{\mathcal{R}}^3\left[0, \frac{\ell}{(\log n)^\gamma}\right] \cap \mathcal{R}^2 = \emptyset, \tilde{S}^3(\ell) = -y\right) \lesssim p_\ell(y) \cdot \frac{1}{\sqrt{\log n}},$$

where in the last inequality we also used [24], Theorem 4.4.1. By [24], Theorem 4.4.1, again we get

$$\mathbb{P}(A_3) \lesssim \frac{1}{\sqrt{\log n}}.$$

We now upper bound the probability $\mathbb{P}(A_1, A_5)$. The probability $\mathbb{P}(A_1, A_7)$ can be bounded using similar ideas. Recall the definition of the event D above. Then from (6.12) and the independence between the three walks, we get

$$\mathbb{P}(D^c, S^3(k) = -x, \tilde{S}^3(\ell) = -y) \leq \frac{1}{(\log n)^4} p_k(x) p_\ell(y).$$

Conditioning on $S^1(i)$, the events A_1 and A_5 become independent, and hence we obtain

$$(6.14) \quad \begin{aligned} & \mathbb{P}(A_1, A_5, D) \\ & \leq \sum_{\|z\| \leq \sqrt{i}(\log n)^{\alpha+2}} \mathbb{P}(A_1 | S^1(i) = z) \mathbb{P}(S^1(i) = z) \mathbb{P}(A_5 | S^1(i) = z). \end{aligned}$$

From Lemma 5.3 again and [24], Theorem 4.3.3, we obtain for all z with $\|z\| \leq \sqrt{i}(\log n)^{\alpha+2}$

$$\mathbb{P}(A_5 | S^1(i) = z) \lesssim \frac{\log \log n}{\log n} \cdot f_\ell(y/2).$$

Plugging this into (6.14) gives

$$\begin{aligned} \mathbb{P}(A_1, A_5, D) &\lesssim f_\ell(y/2) \frac{\log \log n}{\log n} \mathbb{P}(A_1) \lesssim f_\ell(y/2) p_k(x) \frac{\log \log n}{(\log n)^2} \\ &\quad + \mathcal{O}\left(\frac{(\log \log n)^2}{(\log n)^{5/2}} \cdot f_k(x/2) \cdot f_\ell(y/2)\right), \end{aligned}$$

where for the last inequality we used Proposition 5.2. Therefore, we conclude that the sum of probabilities appearing in (6.10) is upper bounded by

$$\begin{aligned} &\frac{1}{(\log n)^{7/3}} \cdot (p_k(x) p_\ell(y) + p_k(x) f_\ell(y/2) + f_k(x/2) f_\ell(y/2)) \\ &\lesssim \frac{1}{(\log n)^{7/3}} f_k(x/2) f_\ell(y/2). \end{aligned}$$

Taking the sum over all k, ℓ and x, y and applying Lemma 6.4 completes the proof. $\square$

PROOF OF LEMMA 6.5. Using Lemma 6.6, it suffices to prove that for all $i, j \in \mathbb{N}$ we have

$$\tilde{\chi}_{n,\alpha}(i, j) = \chi_{n,\alpha}(i, j) + \varepsilon_{n,\alpha}(i, j),$$

where $\mathbb{E}[\varepsilon_{n,\alpha}(i, j)] = o(n/(\log n)^2)$. As in the proof of the previous lemma, we only prove the result for $i = 0$ and $j = 1$, and by using reversibility of the walk, we are led to consider two independent ranges $\mathcal{R}_n$ and $\tilde{\mathcal{R}}_n$ between times 0 and n .

We now define

$$H = \{\mathcal{R}_n \cap (x + \mathcal{R}^1) = \emptyset, S_k = x \notin \mathcal{R}[0, k]\}.$$

Then we have that $H = H_1 \cap H_2$, where

$$H_1 = \{\mathcal{R}[k - n_{4\alpha}, k + n_{4\alpha}] \cap (x + \mathcal{R}^1) = \emptyset, S_k = x \notin \mathcal{R}[k - n_{4\alpha}, k]\},$$

$$H_2 = \{(\mathcal{R}[0, k - n_{4\alpha}] \cup \mathcal{R}[k + n_{4\alpha}, n]) \cap (x + \mathcal{R}^1) = \emptyset, x \notin \mathcal{R}[0, k - n_{4\alpha}]\}.$$

Since $\mathbb{P}(H) = \mathbb{P}(H_1) - \mathbb{P}(H_1 \cap H_2^c)$, using Lemma 6.4 it suffices to prove that for all x and k satisfying $\sqrt{n_{2\alpha}} \leq \|x\| \leq \sqrt{n}(\log n)^2$ and $n_\alpha \leq k \leq n - n_\alpha$, we have

$$(6.15) \quad \mathbb{P}(H_1 \cap H_2^c) \lesssim f_k(x/2) \cdot \frac{\log \log n}{(\log n)^{3/2}}.$$

So we now turn to prove (6.15). We first note that $H_1 \cap H_2^c \subseteq F_1 \cup F_2 \cup F_3$, where

$$\begin{aligned} F_1 &= \{\mathcal{R}[0, k - n_{4\alpha}] \cap (x + \mathcal{R}^1) \neq \emptyset, \mathcal{R}[k, k + n_{4\alpha}] \cap (x + \mathcal{R}^1) = \emptyset, S_k = x\}, \\ F_2 &= \{\mathcal{R}[k + n_{4\alpha}, n] \cap (x + \mathcal{R}^1) \neq \emptyset, \mathcal{R}[k - n_{4\alpha}, k] \cap (x + \mathcal{R}^1) = \emptyset, S_k = x\}, \\ F_3 &= \{S_k = x \in \mathcal{R}[0, k - n_{4\alpha}]\}. \end{aligned}$$

We start by proving the upper bound of (6.15) for $\mathbb{P}(F_1)$. The probability $\mathbb{P}(F_2)$ can be treated in exactly the same way.

First, we decorrelate the two events appearing in F_1 , by conditioning on $S_k = x$ and also by considering $\mathcal{R}^1$ in separate time intervals just like we did in the proof of Lemma 6.6. Let $i = n_{10\alpha} + 4$. Then subtracting x and reversing time we obtain $F_1 \subseteq F_1(1) \cup F_1(2)$, where

$$\begin{aligned} F_1(1) &= \{\mathcal{R}^3[n_{4\alpha}, k] \cap \mathcal{R}^1[1, i] \neq \emptyset, S^3(k) = -x\}, \\ F_1(2) &= \{\mathcal{R}^3[n_{4\alpha}, k] \cap \mathcal{R}^1[i, \infty) \neq \emptyset, \mathcal{R}^4[0, n_{4\alpha}] \cap \mathcal{R}^1[1, i] = \emptyset, S^3(k) = -x\}, \end{aligned}$$

and $\mathcal{R}^3$ and $\mathcal{R}^4$ are two independent ranges. From Lemma 5.3, we get

$$\begin{aligned} \mathbb{P}(F_1(1)) &\lesssim f_k(x/2) \cdot \max(\mathbb{P}_{-x,0}(\mathcal{R}^3[0, k] \cap \mathcal{R}^1[1, i] \neq \emptyset), \\ &\quad \mathbb{P}(\mathcal{R}^3[n_{4\alpha}, k] \cap \mathcal{R}^1[1, i] \neq \emptyset)). \end{aligned}$$

Just like in the proof of Lemma 6.6 we define the event $D = \{\|S^1(r)\| \leq \sqrt{i}(\log n)^2, \forall r \leq i\}$. Then on the event D in order for $\mathcal{R}^3$ and $\mathcal{R}^1[1, i]$ to intersect, the range $\mathcal{R}^3$ must hit the ball centered at 0 of radius $\sqrt{i}(\log n)^2$. By the choice of x and (2.4), this now gives us

$$\max(\mathbb{P}_{-x,0}(\mathcal{R}^3[0, k] \cap \mathcal{R}^1[1, i] \neq \emptyset), \mathbb{P}(\mathcal{R}^3[n_{4\alpha}, k] \cap \mathcal{R}^1[1, i] \neq \emptyset)) \lesssim \frac{1}{(\log n)^4},$$

and hence

$$\mathbb{P}(F_1(1)) \lesssim f_k(x/2) \cdot \frac{1}{(\log n)^4}.$$

We now turn to bound $\mathbb{P}(F_1(2))$. Clearly $\mathbb{P}(F_1(2), D^c) \leq p_k(x)/(\log n)^4$. Now on the event D conditioning on $S^1(i)$, we get

$$\begin{aligned} \mathbb{P}(F_1(2), D) &\leq \sum_{\|z\| \leq \sqrt{i}(\log n)^2} \mathbb{P}(\mathcal{R}^4[0, n_{4\alpha}] \cap \mathcal{R}^1[1, i] = \emptyset, S^1(i) = z) \\ &\quad \times \mathbb{P}(\mathcal{R}^3[n_{4\alpha}, k] \cap \mathcal{R}^1[i, \infty) \neq \emptyset, S^3(k) = -x | S^1(i) = z). \end{aligned}$$

From Lemma 5.3 again, we obtain

$$\begin{aligned} &\mathbb{P}(\mathcal{R}^3[n_{4\alpha}, k] \cap \mathcal{R}^1[i, \infty) \neq \emptyset, S^3(k) = -x | S^1(i) = z) \\ &\lesssim f_k(x/2) \cdot \max(\mathbb{P}_{z,0}(\mathcal{R}^3[n_{4\alpha}, k] \cap \mathcal{R}^1[0, \infty) \neq \emptyset), \end{aligned}$$

$$\begin{aligned} & \mathbb{P}_{z,-x}(\mathcal{R}^3[0, k] \cap \mathcal{R}^1[0, \infty) \neq \emptyset) \\ & \lesssim f_k(x/2) \cdot \frac{\log \log n}{\log n}, \end{aligned}$$

where the last inequality follows from [24], Theorem 4.3.3.

Plugging this above, we finally deduce

$$\mathbb{P}(F_1(2), D) \leq \frac{\log \log n}{\log n} \cdot f_k(x/2) \cdot \frac{1}{\sqrt{\log n}},$$

where the last estimate follows from [24], Theorem 4.4.1.

To complete the proof, we only need to upper bound $\mathbb{P}(F_3)$. By reversing time again, we obtain

$$\mathbb{P}(F_3) \leq \mathbb{P}(S_k = -x, \exists \ell \in [n_{4\alpha}, k] : S_\ell = 0) \lesssim f_k(x/2) \cdot \frac{(\log n)^{2\alpha}}{n},$$

where the last inequality follows from Claim 5.6. Therefore, putting all these bounds together proves (6.15) and this now completes the proof. $\square$

6.3. Moments of χ . In this subsection, we prove the following result.

LEMMA 6.7. *Let $r \in \mathbb{N}$ and let $i_1, j_1, \dots, i_r, j_r$ be integers such that $j_m \leq 2^{i_m-1}$ for all $m \leq r$ (and possibly with repetition). Then for all $\alpha > 12$ as $n \rightarrow \infty$ we have*

$$(6.16) \quad \left(\frac{8 \log n}{\pi^2} \right)^{2r} \cdot \mathbb{E} \left[\prod_{m=1}^r \chi_{n,\alpha}(i_m, j_m) \right] \sim \mathbb{E} \left[\prod_{m=1}^r \left(\sum_{(k,\ell) \in A_{i_m,j_m}^n} G_d(S_k, S_\ell) \right) \right].$$

Before proving the lemma above we state a result on the convergence of discrete quantities to their continuous counterparts. We defer the proof after we prove Lemma 6.7. For all $\beta > 0$, we define the sets D_β and E_β to be the set of time indices and points in space at distance n_β and $\sqrt{n_{2\beta}}$ apart respectively. More precisely, we define

$$(6.17) \quad \begin{aligned} D_\beta &= \{(k_1, \ell_1, \dots, k_r, \ell_r) \in \mathbb{N}^{2r} : (k_m, \ell_m) \in A_{i_m, j_m}^{n, \beta}, \forall m \leq r \text{ and} \\ & |k_m - k_{m'}|, |k_m - \ell_{m'}|, |\ell_m - \ell_{m'}| \geq n_\beta, \forall m \neq m' \leq r\} \end{aligned}$$

and also, with B_β as in (6.4),

$$(6.18) \quad \begin{aligned} E_\beta &= \{(x_1, y_1, \dots, x_r, y_r) \in (\mathbb{Z}^d)^{2r} : (x_m, y_m) \in B_\beta, \forall m \leq r, \text{ and} \\ & \|x_m - x_{m'}\|, \|x_m - y_{m'}\|, \|y_m - y_{m'}\| \geq \sqrt{n_{2\beta}}, \forall m \neq m' \leq r\}. \end{aligned}$$

LEMMA 6.8. *Let $r \in \mathbb{N}$ and let $i_1, j_1, \dots, i_r, j_r$ be integers satisfying $j_m \leq 2^{i_m-1}$ for all $m \leq r$ (and possibly with repetition). Then as $n \rightarrow \infty$ we have*

$$\frac{1}{n^r} \cdot \prod_{m=1}^r \left(\sum_{(k, \ell) \in A_{i_m, j_m}^n} G_d(S_k, S_\ell) \right) \xrightarrow{(d)} \prod_{m=1}^r \int_{A_{i_m, j_m}} 16G(\beta_s, \beta_t) ds dt.$$

Moreover, for all $\beta > 2$ we have

$$\frac{1}{n^r} \cdot \sum_{D_\beta} \mathbb{1}_{(S_{k_m}, S_{\ell_m})_{m \leq r} \in E_\beta} \prod_{m=1}^r G_d(S_{k_m}, S_{\ell_m}) \xrightarrow{(d)} \prod_{m=1}^r \int_{A_{i_m, j_m}} 16G(\beta_s, \beta_t) ds dt,$$

where in the sum above we take $(k_1, \ell_1, \dots, k_r, \ell_r) \in D_\beta$. Finally, in both cases we also have convergence in expectation.

LEMMA 6.9. *There exists a positive constant C so that for all $r \geq 1$ and all $i_1, j_1, \dots, i_r, j_r$ (satisfying $j_m \leq 2^{i_m-1}$ for all $m \leq r$ and possibly with repetition), one has*

$$\mathbb{E} \left[\prod_{m=1}^r \sum_{(k, \ell) \in A_{i_m, j_m}^n} G_d(S_k, S_\ell) \right] \leq Cn^r.$$

REMARK 6.10. Since we allow repetition of the indices, Lemma 6.9 shows that the random variables

$$\frac{1}{n^r} \cdot \prod_{m=1}^r \sum_{(k, \ell) \in A_{i_m, j_m}^n} G_d(S_k, S_\ell)$$

are bounded in L^p for all $p \in \mathbb{N}$.

PROOF OF LEMMA 6.9. The proof of the lemma follows directly by the Cauchy–Schwarz inequality together with [4], Lemma 3.2. $\square$

PROOF OF LEMMA 6.7. For shorthand notation, we write for $k \geq 0$, $x \in \mathbb{Z}^d$, $\alpha > 0$, and $\Lambda \subset \mathbb{Z}^d$,

$$\mathcal{A}_\alpha(k, x, \Lambda) = \{\mathcal{R}[k - n_{4\alpha}, k + n_{4\alpha}] \cap (x + \Lambda) = \emptyset, S_k = x \notin \mathcal{R}[k - n_{4\alpha}, k]\}.$$

Now for multi-indices $i = (i_1, \dots, i_r)$ and $j = (j_1, \dots, j_r)$, write as above $A_{i, j}^{n, \alpha} = A_{i_1, j_1}^{n, \alpha} \times \dots \times A_{i_r, j_r}^{n, \alpha}$. Let $(\mathcal{R}^m)_m$ and $(\tilde{\mathcal{R}}^m)_m$ be the ranges of independent walks

starting from 0. Then we have

$$\begin{aligned}
 & \mathbb{E} \left[\prod_{m=1}^r \chi_{n,\alpha}(i_m, j_m) \right] \\
 (6.19) \quad &= \sum_{A_{i,j}^{n,\alpha}, B_\alpha} \mathbb{P} \left(\bigcap_{m=1}^r (\mathcal{A}_\alpha(k_m, x_m, \mathcal{R}^m) \cap \mathcal{A}_\alpha(\ell_m, y_m, \tilde{\mathcal{R}}^m)) \right) \\
 & \quad \times \prod_{m=1}^r G_d(x_m, y_m),
 \end{aligned}$$

where in the sum above we take $(k_1, \ell_1, \dots, k_r, \ell_r) \in A_{i,j}^{n,\alpha}$ and $(x_m, y_m) \in B_\alpha$ for all $m \leq r$.

First of all, it is obvious that

$$\begin{aligned}
 & \mathbb{E} \left[\prod_{m=1}^r \chi_{n,\alpha}(i_m, j_m) \right] \\
 (6.20) \quad & \geq \sum_{D_\alpha, E_\alpha} \mathbb{P} \left(\bigcap_{m=1}^r (\mathcal{A}_\alpha(k_m, x_m, \mathcal{R}^m) \cap \mathcal{A}_\alpha(\ell_m, y_m, \tilde{\mathcal{R}}^m)) \right) \\
 & \quad \times \prod_{m=1}^r G_d(x_m, y_m).
 \end{aligned}$$

Next, we establish that for a suitable $\beta > \alpha$ we have

$$\begin{aligned}
 & \mathbb{E} \left[\prod_{m=1}^r \chi_{n,\alpha}(i_m, j_m) \right] \\
 (6.21) \quad & \leq \sum_{D_\beta, E_\beta} \mathbb{P} \left(\bigcap_{m=1}^r (\mathcal{A}_\beta(k_m, x_m, \mathcal{R}^m) \cap \mathcal{A}_\beta(\ell_m, y_m, \tilde{\mathcal{R}}^m)) \right) \\
 & \quad \times \prod_{m=1}^r G_d(x_m, y_m) + o(n^r / (\log n)^{2r}).
 \end{aligned}$$

Indeed, we decompose (6.19) into two parts, one over the set $D_\beta \times E_\beta$ and one over the complementary set. Since $\beta > \alpha$, we notice that $\mathcal{A}_\alpha(k, x, \Lambda) \subseteq \mathcal{A}_\beta(k, x, \Lambda)$ for all k, x, Λ . It remains to show that the sum over $(D_\beta \times E_\beta)^c$ is $o(n^r / (\log n)^{2r})$. Forgetting about the intersection events, we can upper bound this sum by

$$\sum_{A_{i,j}^{n,\alpha}, D_\beta^c, B_\alpha} \mathbb{P} \left(\bigcap_{m=1}^r \{S_{k_m} = x_m, S_{\ell_m} = y_m\} \right) \cdot \prod_{m=1}^r G_d(x_m, y_m)$$

$$+ \sum_{A_{i,j}^{n,\alpha}, D_\beta, B_\alpha, E_\beta^c} \mathbb{P} \left(\bigcap_{m=1}^r \{S_{k_m} = x_m, S_{\ell_m} = y_m\} \right) \cdot \prod_{m=1}^r G_d(x_m, y_m).$$

We start by bounding the first sum appearing above. Using that $G_d(x, y) \lesssim 1/n_{2\alpha}$ for all $(x, y) \in B_\alpha$ gives

$$(6.22) \quad \sum_{A_{i,j}^{n,\alpha}, D_\beta^c, B_\alpha} \mathbb{P} \left(\bigcap_{m=1}^r \{S_{k_m} = x_m, S_{\ell_m} = y_m\} \right) \cdot \prod_{m=1}^r G_d(x_m, y_m) \\ \lesssim \frac{n^{2r-1}}{(n_{2\alpha})^r} \cdot n_\beta = \frac{n^r}{(\log n)^{\beta-2\alpha r}},$$

where we used that if $\beta > \alpha$, then $A_{i_m, j_m}^{n,\alpha} \subseteq A_{i_m, j_m}^{n,\beta}$. We now turn to the second sum. Using again the bound on $G_d(x, y)$ for $(x, y) \in B_\alpha$ as above and that on the set D_β all the indices are within distance at least n_β apart, we get

$$(6.23) \quad \sum_{A_{i,j}^{n,\alpha}, D_\beta, B_\alpha, E_\beta^c} \mathbb{P} \left(\bigcap_{m=1}^r \{S_{k_m} = x_m, S_{\ell_m} = y_m\} \right) \cdot \prod_{m=1}^r G_d(x_m, y_m) \\ \leq \frac{n^{2r}}{(n_{2\alpha})^r} \cdot \sup_{t \geq n_\beta} \mathbb{P}(\|S_t\| \leq \sqrt{n_{2\beta}}).$$

Note that we used that n^{2r} is an upper bound on the cardinality of $A_{i,j}^{n,\alpha}$.

Using now (2.4), we can upper bound the second sum by

$$\sum_{A_{i,j}^{n,\alpha}, D_\beta, B_\alpha, E_\beta^c} \mathbb{P} \left(\bigcap_{m=1}^r \{S_{k_m} = x_m, S_{\ell_m} = y_m\} \right) \cdot \prod_{m=1}^r G_d(x_m, y_m) \\ \lesssim \frac{n^r}{(\log n)^{2\beta-2\alpha r}}.$$

Thus taking $2\beta > 2\alpha r + 2r + 1$ proves (6.21).

We next show that for all β sufficiently large we have the following: for any $0 = k_0 < k_1 < \dots < k_r$ and any $0 = x_0, x_1, \dots, x_r$ satisfying $|k_{i+1} - k_i| \geq n_\beta$, $\|x_{i+1} - x_i\| \geq \sqrt{n_{2\beta}}$ and $\|x_i\| \leq \sqrt{n}(\log n)^2$ for all $i < r$ we have

$$(6.24) \quad \mathbb{P} \left(\bigcap_{m=1}^r \mathcal{A}_\beta(k_m, x_m, \mathcal{R}^m) \right) \\ = \left(\frac{\pi^2}{8} \cdot \frac{1}{\log n} \right)^r \cdot \prod_{m=1}^r p_{k_m - k_{m-1}}(x_{m-1}, x_m) \cdot (1 + o(1)) \\ + \mathcal{O} \left(\frac{\log \log n}{(\log n)^{r+\frac{1}{2}}} \cdot \prod_{m=1}^r f_{k_m - k_{m-1}}((x_{m-1} - x_m)/2) \right).$$

We show (6.24) by induction on r . The case $r = 1$ follows from Proposition 5.2. We now assume $r > 1$. We write $j = k_{r-1} + n_{4\beta}$ and define for $x \in \mathbb{Z}^d$

$$B(x) = \{z : \|z - x\| \leq \sqrt{n_{4\beta}} \cdot (\log n)^{3\beta/4}\} \quad \text{and} \quad D = \{S_j \in B(S_{k_{r-1}})\}.$$

We now obtain

$$(6.25) \quad \mathbb{P}\left(\bigcap_{m=1}^r \mathcal{A}_\beta(k_m, x_m, \mathcal{R}^m)\right) = \mathbb{P}\left(\bigcap_{m=1}^r \mathcal{A}_\beta(k_m, x_m, \mathcal{R}^m), D\right) + \mathbb{P}\left(\bigcap_{m=1}^r \mathcal{A}_\beta(k_m, x_m, \mathcal{R}^m), D^c\right).$$

For $z \in B(x_{r-1})$ by the Markov property, we have

$$\begin{aligned} \mathbb{P}\left(\bigcap_{m=1}^r \mathcal{A}_\beta(k_r, x_r, \mathcal{R}^r), S_j = z\right) &= \mathbb{P}(\mathcal{A}_\beta(k_r, x_r, \mathcal{R}^r) | S_j = z) \\ &\quad \times \mathbb{P}\left(\bigcap_{m=1}^{r-1} \mathcal{A}_\beta(k_m, x_m, \mathcal{R}^m), S_j = z\right). \end{aligned}$$

Taking β satisfying $\beta > 8$ and applying Proposition 5.2 to the first term appearing on the right hand side above, we obtain as $n \rightarrow \infty$

$$(6.26) \quad \begin{aligned} \mathbb{P}(\mathcal{A}_\beta(k_r, x_r, \mathcal{R}^r) | S_j = z) &= \frac{\pi^2}{8} \cdot \frac{1}{\log n} \cdot p_{k_r-j}(z, x_r) \cdot (1 + o(1)) \\ &\quad + \mathcal{O}\left(\frac{\log \log n}{(\log n)^{3/2}} \cdot f_{k_r-j}((z - x_r)/2)\right). \end{aligned}$$

Then by Claim 2.2 and (2.3) since $\beta > 8$ we get that

$$p_{k_r-j}(z, x_r) = p_{k_r-k_{r-1}}(x_{r-1}, x_r) \cdot (1 + o(1))$$

and similarly for f . Substituting this into (6.26) gives

$$\begin{aligned} \mathbb{P}(\mathcal{A}_\beta(k_r, x_r, \mathcal{R}^r) | S_j = z) &= \frac{\pi^2}{8} \cdot \frac{1}{\log n} \cdot p_{k_r-k_{r-1}}(x_{r-1}, x_r) \cdot (1 + o(1)) \\ &\quad + \mathcal{O}\left(\frac{\log \log n}{(\log n)^{3/2}} \cdot f_{k_r-k_{r-1}}((x_r - x_{r-1})/2)\right). \end{aligned}$$

Hence overall we obtain

$$\begin{aligned}
 & \mathbb{P}\left(\bigcap_{m=1}^r \mathcal{A}_\beta(k_m, x_m, \mathcal{R}^m), D\right) \\
 &= \sum_{z \in B(x_{p-1})} \mathbb{P}\left(\bigcap_{m=1}^{r-1} \mathcal{A}_\beta(k_m, x_m, \mathcal{R}^m), \mathcal{A}_\beta(k_r, x_r, \mathcal{R}^r), S_j = z\right) \\
 &= \frac{\pi^2}{8} \cdot \frac{1}{\log n} \cdot p_{k_r - k_{r-1}}(x_{r-1}, x_r) \cdot (1 + o(1)) \\
 (6.27) \quad & \times \mathbb{P}\left(\bigcap_{m=1}^{r-1} \mathcal{A}_\beta(k_m, x_m, \mathcal{R}^m), D\right) \\
 &+ \mathcal{O}\left(\mathbb{P}\left(\bigcap_{m=1}^{r-1} \mathcal{A}_\beta(k_m, x_m, \mathcal{R}^m), D\right)\right. \\
 &\quad \left. \times \frac{\log \log n}{(\log n)^{3/2}} \cdot f_{k_r - k_{r-1}}((x_r - x_{r-1})/2)\right).
 \end{aligned}$$

By the Markov property and (2.2), we have

$$\mathbb{P}\left(\bigcap_{m=1}^{r-1} \mathcal{A}_\beta(k_m, x_m, \mathcal{R}^m), D^c\right) \leq \prod_{m=1}^{r-1} p_{k_m - k_{m-1}}(x_{m-1}, x_m) \cdot \exp(-(\log n)^{3\beta/2}).$$

Now note that $p_{k_r - k_{r-1}}(x_{r-1}, x_r) \gtrsim e^{-2(\log n)^{\beta+4}}$ whenever $p_{k_r - k_{r-1}}(x_{r-1}, x_r)$ is positive, since we have taken $k_r - k_{r-1} \geq n^\beta$ and $\|x_r - x_{r-1}\| \leq 2\sqrt{n}(\log n)^2$. Taking β such that $3\beta/2 > \beta + 6$, we have

$$\begin{aligned}
 \mathbb{P}\left(\bigcap_{m=1}^r \mathcal{A}_\beta(k_m, x_m, \mathcal{R}^m), D^c\right) &\leq \prod_{m=1}^{r-1} p_{k_m - k_{m-1}}(x_{m-1}, x_m) \cdot \exp(-(\log n)^{\beta+6}) \\
 &\leq \prod_{m=1}^r p_{k_m - k_{m-1}}(x_{m-1}, x_m) \cdot \exp(-(\log n)^2).
 \end{aligned}$$

Therefore, the two inequalities above together with (6.25) and (6.27) give

$$\begin{aligned}
 & \mathbb{P}\left(\bigcap_{m=1}^r \mathcal{A}_\beta(k_m, x_m, \mathcal{R}^m)\right) \\
 &= \frac{\pi^2}{8 \log n} p_{k_r - k_{r-1}}(x_{r-1}, x_r) \cdot \mathbb{P}\left(\bigcap_{m=1}^{r-1} \mathcal{A}_\beta(k_m, x_m, \mathcal{R}^m)\right) (1 + o(1)) \\
 &\quad + \mathcal{O}\left(\prod_{m=1}^r p_{k_m - k_{m-1}}(x_{m-1}, x_m) \exp(-(\log n)^2)\right)
 \end{aligned}$$

$$\begin{aligned}
& + \mathcal{O}\left(\mathbb{P}\left(\bigcap_{m=1}^{r-1} \mathcal{A}_\beta(k_m, x_m, \mathcal{R}^m)\right) \frac{\log \log n}{(\log n)^{3/2}} f_{k_r-k_{r-1}}((x_r - x_{r-1})/2)\right) \\
& + \mathcal{O}\left(f_{k_r-k_{r-1}}((x_r - x_{r-1})/2)\right. \\
& \quad \times \left.\prod_{m=1}^{r-1} p_{k_m-k_{m-1}}(x_{m-1}, x_m) \exp(-(\log n)^{3\beta/2})\right).
\end{aligned}$$

Note that for (x_i) and (k_i) as above, we have from (2.3) that

$$p_{k_m-k_{m-1}}(x_{m-1}, x_m) \lesssim f_{k_m-k_{m-1}}((x_m - x_{m-1})/2).$$

So we can replace the product appearing in the last $\mathcal{O}$ term above by $\prod_{m=1}^{r-1} f_{k_m-k_{m-1}}((x_m - x_{m-1})/2)$. We use the previous equation to complete the inductive proof of (6.24).

We now explain how to conclude the proof of the lemma from (6.24).

Since (6.24) holds for all $\beta > 12$ and since $\alpha > 12$, we can plug (6.24) into (6.20) to get

$$\begin{aligned}
& \mathbb{E}\left[\prod_{m=1}^r \chi_{n,\alpha}(i_m, j_m)\right] \\
& \geq \left(\frac{\pi^2}{8} \cdot \frac{1}{\log n}\right)^{2r} \mathbb{E}\left[\sum_{D_\alpha} \mathbb{1}_{(S_{k_m}, S_{\ell_m})_{m \leq r} \in E_\alpha} \prod_{m=1}^r G_d(S_{k_m}, S_{\ell_m})\right] (1 + o(1)) \\
& \quad + \frac{\log \log n}{(\log n)^{2r+1/2}} \mathcal{O}\left(\sum_{D_\alpha, E_\alpha} \prod_{m=1}^{2r} f_{s_i-s_{i-1}}((z_i - z_{i-1})/2) \prod_{m=1}^r G_d(x_m, y_m)\right),
\end{aligned}$$

where (s_i) is the increasing rearrangement of the indices (k_m) and (ℓ_m) and (z_i) are the corresponding rearrangement of (x_m) and (y_m) . Similarly, plugging (6.24) into (6.21) we obtain

$$\begin{aligned}
& \mathbb{E}\left[\prod_{m=1}^r \chi_{n,\alpha}(i_m, j_m)\right] \\
& \leq o(n^r / (\log n)^{2r}) \\
& \quad + \left(\frac{\pi^2}{8} \cdot \frac{1}{\log n}\right)^{2r} \mathbb{E}\left[\sum_{D_\beta} \mathbb{1}_{(S_{k_m}, S_{\ell_m})_{m \leq r} \in E_\beta} \prod_{m=1}^r G_d(S_{k_m}, S_{\ell_m})\right] (1 + o(1)) \\
& \quad + \frac{\log \log n}{(\log n)^{2r+1/2}} \mathcal{O}\left(\sum_{D_\beta, E_\beta} \prod_{m=1}^{2r} f_{s_i-s_{i-1}}((z_i - z_{i-1})/2) \prod_{m=1}^r G_d(x_m, y_m)\right).
\end{aligned}$$

By Lemma 6.8, since both sequences appearing below converge in expectation to the same limit, we get that as $n \rightarrow \infty$,

$$\mathbb{E} \left[\sum_{D_\beta} \mathbb{1}_{(S_{k_m}, S_{\ell_m})_{m \leq r} \in E_\beta} \prod_{m=1}^r G_d(S_{k_m}, S_{\ell_m}) \right] \sim \mathbb{E} \left[\prod_{m=1}^r \left(\sum_{(k, \ell) \in A_{im, jm}^n} G_d(S_k, S_\ell) \right) \right].$$

Using that $f_k(x/2) \asymp f_{4k}(x) \asymp p_{4k}(x) + p_{4k+1}(x)$ and Lemma 6.9, we get that the $\mathcal{O}$ terms appearing in the two inequalities above are $\mathcal{O}(n^r)$ and this now completes the proof of the lemma. $\square$

6.4. Proof of Proposition 6.1. In this section, we give the proof of Proposition 6.1. We start by proving Lemma 6.8.

PROOF OF LEMMA 6.8. We start by proving the first statement of the lemma. Let $\varepsilon > 0$ and φ_ε be a continuous function satisfying

$$\mathbb{1}(\|x\| \geq \varepsilon) \leq \varphi_\varepsilon(x) \leq \mathbb{1}(\|x\| \geq \varepsilon/2).$$

Recall from (2.6) that

$$G_d(x) = 4G(x) + \mathcal{O}\left(\frac{1}{1 + \|x\|^4}\right).$$

To simplify notation, we now write

$$\begin{aligned} X(m) &= \sum_{(k, \ell) \in A_{im, jm}^n} G_d(S_k, S_\ell), \quad X_\varepsilon(m) \\ &= \sum_{(k, \ell) \in A_{im, jm}^n} G_d(S_k, S_\ell) \varphi_\varepsilon\left(\frac{S_k - S_\ell}{2\sqrt{n}}\right) \quad \text{and} \\ \tilde{X}_\varepsilon(m) &= \sum_{(k, \ell) \in A_{im, jm}^n} 4G(S_k, S_\ell) \varphi_\varepsilon\left(\frac{S_k - S_\ell}{2\sqrt{n}}\right). \end{aligned}$$

Using Lemma 6.9, it is straightforward to check that

$$(6.28) \quad \frac{1}{n^r} \cdot \mathbb{E} \left[\prod_{m=1}^r X_\varepsilon(m) - \prod_{m=1}^r \tilde{X}_\varepsilon(m) \right] = \mathcal{O}(1/n).$$

Note that the function $G(x)\varphi_\varepsilon(x)$ is continuous and bounded, and that by Donsker's invariance principle $(S_{[nt]}/2\sqrt{n}, t \geq 0)$ converges in law to a standard Brownian motion. Hence we obtain as $n \rightarrow \infty$,

$$(6.29) \quad \frac{1}{n^r} \cdot \prod_{m=1}^r \tilde{X}_\varepsilon(m) \xrightarrow{(d)} \prod_{m=1}^r \int_{A_{im, jm}} 16G(\beta_s, \beta_t) \varphi_\varepsilon(\beta_s - \beta_t) ds dt.$$

For all $m \leq r$, we have

$$X(m) - X_\varepsilon(m) \leq \sum_{(k,\ell) \in A_{i_m, j_m}^n} G_d(S_k, S_\ell) \mathbb{1}(\|S_k - S_\ell\| \leq 2\varepsilon\sqrt{n}),$$

and hence using Lemma 6.3 we get

$$(6.30) \quad \mathbb{E}[X(m) - X_\varepsilon(m)] \lesssim n \cdot \varepsilon \log\left(\frac{1}{\varepsilon}\right).$$

For each $\varepsilon > 0$, we now define

$$R_n(\varepsilon) = \frac{1}{n^r} \cdot \prod_{m=1}^r X(m) - \frac{1}{n^r} \cdot \prod_{m=1}^r X_\varepsilon(m).$$

In view of (6.28) and (6.29), in order to complete the proof of the first statement, it suffices to prove that $\mathbb{E}[R_n(\varepsilon)] \rightarrow 0$ as $\varepsilon \rightarrow 0$ uniformly in n .

With the definitions above, we can upper bound $R_n(\varepsilon)$ by

$$R_n(\varepsilon) \leq \frac{1}{n^r} \cdot \sum_{w=1}^r (X(w) - X_\varepsilon(w)) \cdot \prod_{\substack{m=1 \\ m \neq w}}^r X(m).$$

We now set $Z(w)$ to be equal to the product appearing above. Using the obvious upper bound $X(m) - X_\varepsilon(m) \leq (X(m) - X_\varepsilon(m))^{1/2} \cdot (X(m))^{1/2}$ and Hölder's inequality applied twice gives the following bound for $\mathbb{E}[R_n(\varepsilon)]$:

$$\frac{1}{n^r} \cdot \sum_{w=1}^r (\mathbb{E}[X(w) - X_\varepsilon(w)])^{1/2} \cdot (\mathbb{E}[(X(m))^3])^{1/6} \cdot (\mathbb{E}[(Z(w))^3])^{1/3}.$$

Lemma 6.9 and (6.30) now give that

$$(6.31) \quad \mathbb{E}[R_n(\varepsilon)] \leq C_1 \cdot \sqrt{\varepsilon \log\left(\frac{1}{\varepsilon}\right)} \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0$$

and this proves the first convergence.

For the second statement, we note that using Cauchy–Schwarz and similar arguments as in Lemma 6.2 one can remove the sets D_β and E_β and then apply the first part of the lemma.

Finally, we get the convergence in expectation as a consequence of weak convergence together with uniform integrability which follows directly from Lemma 6.9. $\square$

PROOF OF PROPOSITION 6.1. By Cramer–Wold in order to deduce the weak convergence, it suffices to prove that all linear combinations of variables on the left converge weakly to the corresponding linear combinations of variables on the right. Lemma 6.5 shows that one can replace the terms $\chi_n(i, j)$ by their localised

versions, $\chi_{n,\alpha}(i, j)$. Lemmas 6.7 and 6.8 show that the moments of all linear combinations of the $\chi_{n,\alpha}(i, j)$ do actually converge to the corresponding moments. We only need to ensure that the moments of the limiting object uniquely characterise the distribution. This now follows from Proposition 4.1 using Carleman's criterion (see Section 3.3.5 in [12]). Indeed, for any variable X on the right of (6.2), Proposition 4.1 gives that

$$\mathbb{E}[X^p] \leq C^p p^{2p}.$$

Therefore, if X and Y are two of the variables on the right, then by the triangle inequality for the L^p -norm we get

$$\mathbb{E}[(X + Y)^p]^{1/p} \leq \mathbb{E}[X^p]^{1/p} + \mathbb{E}[Y^p]^{1/p} \leq 2C \cdot p^2.$$

Therefore, Carleman's condition also holds for the sum $X + Y$, hence its distribution is uniquely characterised by its moments. $\square$

7. Central limit theorem. In this section, we finally give the proof of Theorem 1.2.

We start by proving an upper bound on the variance of $\text{Cap}(\mathcal{R}_n)$ using the same technique as Le Gall did for the range in dimension 2.

LEMMA 7.1. *We have $\mathbb{E}[(\text{Cap}(\mathcal{R}_n) - \mathbb{E}[\text{Cap}(\mathcal{R}_n)])^4] \lesssim n^4/(\log n)^8$.*

PROOF. The proof follows in the same way as [28], Lemma 6.2, and [4], Lemma 3.5. We write $X_n = \text{Cap}(\mathcal{R}_n)$ and $\bar{X} = X - \mathbb{E}[X]$. We now write $\ell = \lfloor n/2 \rfloor$ and $m = n - \ell$. Then from Proposition 2.3, we get

$$|\bar{X}_n - \bar{X}_\ell^{(1)} - \bar{X}_m^{(2)}| = |\overline{\chi_n(1, 1)} + \bar{\varepsilon}_n|.$$

Lemmas 6.7 and 6.9 give that

$$\mathbb{E}[\chi_n(1, 1)] \lesssim \frac{n}{(\log n)^2} \quad \text{and} \quad \mathbb{E}[(\overline{\chi_n(1, 1)})^2] \lesssim \frac{n^2}{(\log n)^4}.$$

Also from Proposition 2.3, we have that $\mathbb{E}[\varepsilon_n] = \mathcal{O}(\log n)$ and $\mathbb{E}[\varepsilon_n^2] = \mathcal{O}((\log n)^2)$. The rest of the proof follows in exactly the same way as in [4], Lemma 4.2. $\square$

PROOF OF THEOREM 1.2. For any fixed $p \geq 1$, Proposition 2.3 shows that

$$\text{Cap}(\mathcal{R}_n) = \sum_{j=1}^{2^p} \text{Cap}(\mathcal{R}_n^{(p,j)}) - \sum_{i=1}^p \sum_{j=1}^{2^{i-1}} \chi_n(i, j) + \varepsilon_n.$$

We write $\overline{X} = X - \mathbb{E}[X]$. Subtracting the expectation in the equation above, we obtain

$$\overline{\text{Cap}(\mathcal{R}_n)} = \sum_{j=1}^{2^p} \overline{\text{Cap}(\mathcal{R}_n^{(p,j)})} - \sum_{i=1}^p \sum_{j=1}^{2^{i-1}} \overline{\chi_n(i, j)} + \overline{\varepsilon_n}.$$

Lemma 7.1 and the independence of the ranges $\mathcal{R}_n^{(p,j)}$ immediately give that

$$\mathbb{E} \left[\left(\frac{(\log n)^2}{n} \sum_{j=1}^{2^p} \overline{\text{Cap}(\mathcal{R}_n^{(p,j)})} \right)^2 \right] \lesssim 2^{-p}.$$

Since $\mathbb{E}[\varepsilon_n] = o(n/(\log n)^2)$ from Proposition 2.3, we get that

$$\frac{(\log n)^2}{n} \cdot \overline{\varepsilon_n} \xrightarrow{\mathbb{P}} 0 \quad \text{as } n \rightarrow \infty.$$

Moreover, using Proposition 6.1 we get that for a fixed p , as n goes to infinity

$$\frac{2(\log n)^2}{\pi^4 \cdot n} \cdot \sum_{i=1}^p \sum_{j=1}^{2^{i-1}} \overline{\chi_n(i, j)} \xrightarrow{(d)} \sum_{i=1}^p \sum_{j=1}^{2^{i-1}} \gamma_G(A_{i,j}).$$

From Proposition 4.5, we also have in L^2 -norm:

$$\lim_{p \rightarrow \infty} \sum_{i=1}^p \sum_{j=1}^{2^{i-1}} \gamma_G(A_{i,j}) = \gamma_G(\mathcal{C}) = \frac{1}{2} \gamma_G([0, 1]^2).$$

By first taking p large enough, and then letting $n \rightarrow \infty$ completes the proof. $\square$

PROOF OF COROLLARY 1.5. Lemma 7.1 shows that $(\log n)^4/n^2(\text{Cap}(\mathcal{R}_n) - \mathbb{E}[\text{Cap}(\mathcal{R}_n)])^2$ is uniformly integrable. Hence this together with the convergence in distribution from Theorem 1.2 proves the corollary. $\square$

APPENDIX

In this section we prove the two lemma that were used in the proof of Proposition 4.1.

PROOF OF LEMMA 4.3. First, using the explicit density of the Brownian motion we get

$$\begin{aligned} \int_0^1 \mathbb{E} \left[\frac{(a + |\log \|\beta_s - x\||)^k}{\|\beta_s - x\|^2} \right] ds &\lesssim \int_0^1 \frac{ds}{s^2} \int_{\mathbb{R}^4} \frac{(a + |\log \|u\||)^k}{\|u\|^2} e^{-\frac{\|u+x\|^2}{2s}} du \\ &\lesssim \int_{\mathbb{R}^4} \frac{(a + |\log \|u\||)^k}{\|u\|^2 \cdot \|u+x\|^2} e^{-\frac{\|u+x\|^2}{2}} du, \end{aligned}$$

using Fubini at the second line. Now define

$$F(u, x) := \frac{(a + |\log \|u\||)^k}{\|u\|^2 \cdot \|u + x\|^2} \cdot e^{-\frac{\|u+x\|^2}{2}}.$$

We bound F as follows:

$$F(u, x) \lesssim \begin{cases} \frac{(a + |\log \|u\||)^k}{\|u\|^4} \cdot e^{-\frac{\|u\|^2}{8}} & \text{if } \|u\| \geq 2\|x\|, \\ \frac{(a + |\log \|u\||)^k}{\|u\|^2 \cdot \|x\|^2} \cdot e^{-\frac{\|x\|^2}{8}} & \text{if } \|u + x\| \geq \frac{\|x\|}{2} \text{ and} \\ & \|u\| \leq 2\|x\|, \\ \frac{(a + 1 + |\log \|x\||)^k}{\|x\|^2 \cdot \|u + x\|^2} \cdot e^{-\frac{\|u+x\|^2}{2}} & \text{if } \|u + x\| \leq \frac{\|x\|}{2}. \end{cases}$$

Then we obtain with appropriate changes of variables

$$\int_{\mathbb{R}^4} F(u, x) du \lesssim I_1(x) + I_2(x) + I_3(x),$$

with

$$\begin{aligned} I_1(x) &= \int_{2\|x\|}^{\infty} \frac{(a + |\log r|)^k}{r} \cdot e^{-\frac{r^2}{8}} dr, \\ I_2(x) &= \|x\|^{-2} \cdot e^{-\frac{\|x\|^2}{8}} \int_0^{2\|x\|} r(a + |\log r|)^k dr, \\ I_3(x) &= \frac{(a + 1 + |\log \|x\||)^k}{\|x\|^2} \int_0^{\|x\|} r e^{-r^2/2} dr. \end{aligned}$$

Note that $I_3(x) \lesssim (a + 1 + |\log \|x\||)^k$. Moreover, $I_1(x)$ and $I_2(x)$ will be bounded using the two following claims.

CLAIM A.1. For all $a \geq 0$, $b > 0$ and $k \geq 1$, we have

$$\int_0^b (a + |\log r|)^k r dr \lesssim b^2 \cdot \sum_{\ell=0}^k (a + |\log b|)^{k-\ell} \cdot k^\ell.$$

PROOF. Using the change of variable $r = \exp(-u)$, we obtain

$$f(k) := \int_0^b (a + |\log r|)^k r dr = \int_{-\log b}^{\infty} (a + |u|)^k e^{-2u} du.$$

Assume first that $b < 1$, so that $-\log b$ is nonnegative. Then an integration by parts gives

$$f(k) = \frac{b^2}{2} (a + |\log b|)^k + \frac{k}{2} f(k-1),$$

leading to the desired result by induction. Now if $b \geq 1$, one has

$$f(k) = \int_0^{\log b} (a+u)^k e^{2u} du + \int_0^\infty (a+u)^k e^{-2u} du,$$

and an integration by parts gives similarly

$$f(k) \lesssim b^2(a + \log b)^k + kf(k-1),$$

and the claim follows by induction. $\square$

CLAIM A.2. *For all $a \geq 0$, $b > 0$, and $k \geq 0$ we have*

$$\begin{aligned} & \int_b^\infty \frac{(a + |\log r|)^k}{r} e^{-r^2/8} dr \\ & \lesssim \frac{(a + |\log b|)^{k+1}}{k+1} + e^{-b^2/8} \cdot \sum_{\ell=0}^k (4k)^\ell (a + |\log b|)^{k-\ell}. \end{aligned}$$

PROOF. Assume first that $b \geq 1$, and note that

$$\int_b^\infty \frac{(a + |\log r|)^k}{r} e^{-r^2/8} dr \leq g(k, b) := \int_b^\infty (a + |\log r|)^k r e^{-r^2/8} dr.$$

Moreover, an integration by parts yields

$$g(k, b) \leq 4(a + \log b)^k e^{-b^2/8} + 4kg(k-1, b),$$

which gives the result by induction. Now if $b < 1$, we have

$$\begin{aligned} \int_b^\infty \frac{(a + |\log r|)^k}{r} e^{-r^2/8} dr & \leq \int_b^1 \frac{(a - \log r)^k}{r} dr + g(k, 1) \\ & = \frac{(a + |\log b|)^{k+1}}{k+1} + g(k, 1), \end{aligned}$$

and using the previous estimate for $g(k, 1)$, this concludes the proof of the claim. $\square$

Now we can just apply these two claims with $b = 2\|x\|$ and use that $|\log 2\|x\|| \leq 1 + |\log \|x\||$. This gives the desired upper bounds for $I_1(x)$ and $I_2(x)$ and concludes the proof of Lemma 4.3. $\square$

PROOF OF LEMMA 4.4. We have

$$\begin{aligned} & \int_0^1 \mathbb{E}[(a + |\log \|\tilde{\beta}_t\||)^k] dt \\ & \lesssim \int_0^1 \frac{dt}{t^2} \int_{\mathbb{R}^4} (a + |\log \|u\||)^k e^{-\|u\|^2/(2t)} du \end{aligned}$$

$$\begin{aligned} &\lesssim \int_{\mathbb{R}^4} \frac{(a + |\log \|u\||)^k}{\|u\|^2} e^{-\|u\|^2/2} du \lesssim \int_0^\infty (a + |\log r|)^k r e^{-r^2/2} dr \\ &\lesssim \int_0^1 (a + |\log r|)^k r dr + \int_1^\infty (a + |\log r|)^k r e^{-r^2/2} dr. \end{aligned}$$

Now using the same argument as in the proof of Claim A.2 for the second integral and Claim A.1 with $b = 1$ for the first one, we obtain the lemma. $\square$

Acknowledgements. We thank Yinshan Chang for kindly sharing with us his manuscript, which motivated us to look deeper at fluctuations in dimension four. We thank the referee for the thorough review of the paper.

REFERENCES

- [1] AIZENMAN, M. (1985). The intersection of Brownian paths as a case study of a renormalization group method for quantum field theory. *Comm. Math. Phys.* **97** 91–110. [MR0782960](#)
- [2] ALBEVERIO, S. and ZHOU, X. Y. (1996). Intersections of random walks and Wiener sausages in four dimensions. *Acta Appl. Math.* **45** 195–237. [MR1414282](#)
- [3] ASSELAH, A. and SCHAPIRA, B. (2017). Moderate deviations for the range of a transient random walk: Path concentration. *Ann. Sci. Éc. Norm. Supér. (4)* **50** 755–786. [MR3665554](#)
- [4] ASSELAH, A., SCHAPIRA, B. and SOUSI, P. (2018). Capacity of the range of random walk on $\mathbb{Z}^d$. *Trans. Amer. Math. Soc.* **370** 7627–7645. [MR3852443](#)
- [5] ASSELAH, A., SCHAPIRA, B. and SOUSI, P. (2018). Strong law of large numbers for the capacity of the Wiener sausage in dimension four. *Probab. Theory Related Fields* **173** 813–858. [MR3936147](#)
- [6] BRYDGES, D. C. and SPENCER, T. (1985). Self-avoiding random walk and the renormalisation group. In *Applications of Field Theory to Statistical Mechanics (Sitges, 1984)*. *Lecture Notes in Physics* **216** 189–198. Springer, Berlin. [MR0781962](#)
- [7] BURDZY, K. and LAWLER, G. F. (1990). Nonintersection exponents for Brownian paths. I. Existence and an invariance principle. *Probab. Theory Related Fields* **84** 393–410. [MR1035664](#)
- [8] CHANG, Y. (2017). Two observations on the capacity of the range of simple random walks on $\mathbb{Z}^3$ and $\mathbb{Z}^4$. *Electron. Commun. Probab.* **22** Paper No. 25, 9. [MR3652038](#)
- [9] CHANG, Y. and SAPOZHNIKOV, A. (2016). Phase transition in loop percolation. *Probab. Theory Related Fields* **164** 979–1025. [MR3477785](#)
- [10] DUPLANTIER, B. (1998). Random walks and quantum gravity in two dimensions. *Phys. Rev. Lett.* **81** 5489–5492. [MR1666816](#)
- [11] DUPLANTIER, B. and KWON, K.-H. (1988). Conformal invariance and intersections of random walks. *Phys. Rev. Lett.* **61** 2514–2517.
- [12] DURRETT, R. (2010). *Probability: Theory and Examples*, 4th ed. *Cambridge Series in Statistical and Probabilistic Mathematics* **31**. Cambridge Univ. Press, Cambridge. [MR2722836](#)
- [13] DVORETZKY, A. and ERDŐS, P. (1951). Some problems on random walk in space. In *Proceedings of the Second Berkeley Symposium on Mathematical Statistics and Probability*, 1950 353–367. Univ. California Press, Berkeley, CA. [MR0047272](#)
- [14] DVORETZKY, A., ERDŐS, P. and KAKUTANI, S. (1950). Double points of paths of Brownian motion in n -space. *Acta Sci. Math. (Szeged)* **12** 75–81. [MR0034972](#)
- [15] ERHARD, D. and POISAT, J. (2016). Asymptotics of the critical time in Wiener sausage percolation with a small radius. *ALEA Lat. Am. J. Probab. Math. Stat.* **13** 417–445. [MR3519252](#)

- [16] FELDER, G. and FRÖHLICH, J. (1985). Intersection properties of simple random walks: A renormalization group approach. *Comm. Math. Phys.* **97** 111–124. [MR0782961](#)
- [17] FERNÁNDEZ, R., FRÖHLICH, J. and SOKAL, A. D. (1992). *Random Walks, Critical Phenomena, and Triviality in Quantum Field Theory*. Springer, Berlin. [MR1219313](#)
- [18] JAIN, N. and OREY, S. (1968). On the range of random walk. *Israel J. Math.* **6** 373–380. [MR0243623](#)
- [19] JAIN, N. C. and PRUITT, W. E. (1971). The range of transient random walk. *J. Anal. Math.* **24** 369–393. [MR0283890](#)
- [20] KHOSHNEVISAN, D. (2003). Intersections of Brownian motions. *Expo. Math.* **21** 97–114. [MR1978059](#)
- [21] LAWLER, G. F. (1980). A self-avoiding random walk. *Duke Math. J.* **47** 655–693. [MR0587173](#)
- [22] LAWLER, G. F. (1982). The probability of intersection of independent random walks in four dimensions. *Comm. Math. Phys.* **86** 539–554. [MR0679202](#)
- [23] LAWLER, G. F. (1985). Intersections of random walks in four dimensions. II. *Comm. Math. Phys.* **97** 583–594. [MR0787120](#)
- [24] LAWLER, G. F. (1991). *Intersections of Random Walks*. Birkhäuser, Boston, MA. [MR1117680](#)
- [25] LAWLER, G. F. and LIMIC, V. (2010). *Random Walk: A Modern Introduction. Cambridge Studies in Advanced Mathematics* **123**. Cambridge Univ. Press, Cambridge. [MR2677157](#)
- [26] LAWLER, G. F., SCHRAMM, O. and WERNER, W. (2001). Values of Brownian intersection exponents. II. Plane exponents. *Acta Math.* **187** 275–308. [MR1879851](#)
- [27] LE GALL, J.-F. (1985). Sur le temps local d’intersection du mouvement brownien plan et la méthode de renormalisation de Varadhan. In *Séminaire de Probabilités, XIX, 1983/84. Lecture Notes in Math.* **1123** 314–331. Springer, Berlin. [MR0889492](#)
- [28] LE GALL, J.-F. (1986). Propriétés d’intersection des marches aléatoires. I. Convergence vers le temps local d’intersection. *Comm. Math. Phys.* **104** 471–507. [MR0840748](#)
- [29] LE GALL, J.-F. (1988). Fluctuation results for the Wiener sausage. *Ann. Probab.* **16** 991–1018. [MR0942751](#)
- [30] LE GALL, J.-F. (1994). Exponential moments for the renormalized self-intersection local time of planar Brownian motion. In *Séminaire de Probabilités, XXVIII. Lecture Notes in Math.* **1583** 172–180. Springer, Berlin. [MR1329112](#)
- [31] LE GALL, J.-F. and ROSEN, J. (1991). The range of stable random walks. *Ann. Probab.* **19** 650–705. [MR1106281](#)
- [32] MADRAS, N. and SLADE, G. (2013). *The Self-Avoiding Walk*. Birkhäuser/Springer, New York. [MR2986656](#)
- [33] PEMANTLE, R., PERES, Y. and SHAPIRO, J. W. (1996). The trace of spatial Brownian motion is capacity-equivalent to the unit square. *Probab. Theory Related Fields* **106** 379–399. [MR1418846](#)
- [34] RÁTH, B. and SAPOZHNIKOV, A. (2012). Connectivity properties of random interlacement and intersection of random walks. *ALEA Lat. Am. J. Probab. Math. Stat.* **9** 67–83. [MR2889752](#)
- [35] SYMANZIK, K. (1969). Euclidean quantum field theory. In *Local Quantum Theory* (R. Jost, ed.) 152–226. Academic Press, New York.
- [36] SZNITMAN, A.-S. (2010). Vacant set of random interacements and percolation. *Ann. of Math.* (2) **171** 2039–2087. [MR2680403](#)
- [37] VAN DEN BERG, M., BOLTHAUSEN, E. and DEN HOLLANDER, F. (2004). On the volume of the intersection of two Wiener sausages. *Ann. of Math.* (2) **159** 741–782. [MR2081439](#)
- [38] VAN DEN BERG, M., BOLTHAUSEN, E. and DEN HOLLANDER, F. (2018). Torsional rigidity for regions with a Brownian boundary. *Potential Anal.* **48** 375–403. [MR3779094](#)

A. ASSELAH
LAMA (UMR 8050),
UPEC, UPEMLV
CNRS
UNIVERSITÉ PARIS-EST
F-94010, CRÉTEIL
FRANCE
E-MAIL: amine.asselah@u-pec.fr

B. SCHAPIRA
I2M, UMR 7373
AIX-MARSEILLE UNIVERSITÉ
13453 MARSEILLE
FRANCE
E-MAIL: bruno.schapira@univ-amu.fr

P. SOUSI
DEPARTMENT OF PURE MATHEMATICS
CAMBRIDGE UNIVERSITY
CAMBRIDGE CB30WB
UNITED KINGDOM
E-MAIL: p.sousi@statslab.cam.ac.uk