



Contents lists available at ScienceDirect

C. R. Acad. Sci. Paris, Ser. I

www.sciencedirect.com



Probability Theory

A balanced excited random walk

Une marche excitée équilibrée

Itai Benjamini^a, Gady Kozma^a, Bruno Schapira^b

^a The Weizmann Institute of Science, Rehovot POB 76100, Israel

^b Département de Mathématiques, bâtiment 425, Université Paris-Sud 11, 91405 Orsay cedex, France

ARTICLE INFO

Article history:

Received 7 September 2010

Accepted after revision 21 February 2011

Available online 10 March 2011

Presented by Marc Yor

ABSTRACT

The following random process on $\mathbb{Z}^4$ is studied. At first visit to a site, the two first coordinates perform a (2-dimensional) simple random walk step. At further visits, it is the last two coordinates which perform a simple random walk step. We prove that this process is almost surely transient. The lower dimensional versions are discussed and various generalizations and related questions are proposed.

© 2011 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

RÉSUMÉ

Nous étudions le processus suivant sur $\mathbb{Z}^4$. À la première visite en un site, les deux premières coordonnées effectuent un saut d'une marche simple (2-dimensionnelle). Aux visites suivantes en ce site, ce sont les deux dernières coordonnées qui effectuent un saut de marche simple. Nous montrons que ce processus est presque sûrement transitoire. Nous discutons également des dimensions inférieures et divers généralisations et questions connexes sont proposées.

© 2011 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

1. Introduction

The excited random walk as defined by Benjamini and Wilson [2] has a bias in some fixed direction, a feature which is highly useful in its analysis. See e.g. [10] and references within. Attempts to relax the dependence of the proof structure on monotonicity resulted in a number of works where the walker has competing drifts. See [1,6,4]. One motivation was to get closer to standard models of reinforced random walks on $\mathbb{Z}^d$, which are symmetric in nature. We think for instance on the question of recurrence vs. transience of 1-reinforced random walks, which is still widely open (see the surveys [11] and [12]). With the same goal in mind, we started exploring excited-like models where the walker is in addition also a martingale or a bounded perturbation of one, and posed some questions in 2007 [7], which, it seems, are all still open. Progress on this kind of models was achieved in [5], but there the laws were not nearest-neighbors. Here we describe and solve one such model of a nearest-neighbor walk in 4 dimensions.

We describe a general form of the model in any dimension $d \geq 2$, but we will actually only deal with 4 dimensions here. So one has first to choose arbitrarily two integers $d_1 \geq 1$ and $d_2 \geq 1$ such that $d = d_1 + d_2$. Then we define the process $(S_n, n \geq 0)$ on $\mathbb{Z}^d$ as a mixture of two simple random walks in the following sense. Set $S_n = (X_n, Y_n)$, where $X_n \in \mathbb{Z}^{d_1}$ is the set of the first d_1 coordinates of S_n and $Y_n \in \mathbb{Z}^{d_2}$ is the set of the last d_2 coordinates. Now the rule is the following. First $S_0 = 0$. Next if S visits a site for the first time then only the X component performs a simple random walk step, that is:

E-mail addresses: itai.benjamini@weizmann.ac.il (I. Benjamini), gady.kozma@weizmann.ac.il (G. Kozma), bruno.schapira@math.u-psud.fr (B. Schapira).

$$\mathbb{P}[S_{n+1} - S_n = (0, \dots, 0, \pm 1, 0, \dots, 0) \mid \mathcal{F}_n] = \begin{cases} 1/(2d_1) & S_n \neq S_i \forall i < n, \\ 0 & \text{otherwise,} \end{cases}$$

where the ± 1 can be at any of the first d_1 coordinates. Otherwise, only Y performs a simple random walk step:

$$\mathbb{P}[S_{n+1} - S_n = (0, \dots, 0, \pm 1, 0, \dots, 0) \mid \mathcal{F}_n] = \begin{cases} 1/(2d_2) & \exists i < n \text{ } S_i = S_n, \\ 0 & \text{otherwise,} \end{cases}$$

where this time the ± 1 can be at any of the last d_2 coordinates. We call this process S the $M(d_1, d_2)$ -random walk.

Here we say that a process is transient if almost surely any site is visited only finitely many times. It is said to be recurrent if almost surely it visits all sites infinitely often. We will prove the following:

Theorem 1. *The $M(2, 2)$ -random walk is transient.*

The proof of Theorem 1 is elementary, and uses only basic estimates on the standard 2-dimensional simple random walk. It relies on finding good upper bounds for the probability of return to the origin and then uses the Borel–Cantelli Lemma (what makes however the proof nontrivial is that the two components X and Y are not independent).

Note that the canonical projections of S on $\mathbb{Z}^{d_1}$ and $\mathbb{Z}^{d_2}$ are usual (time changed) simple random walks. So if $d = 4$, and if d_1 or d_2 equals 3, then S is automatically transient, since the simple random walk on $\mathbb{Z}^3$ is transient. Likewise if d is larger than 5, then for any choice of d_1 and d_2 , the resulting process will be transient. Thus the question of recurrence vs. transience is only interesting in dimension less than or equal to 4. In dimension 3 there are two versions: $d_1 = 1$ and $d_2 = 2$ or $d_1 = 2$ and $d_2 = 1$. We conjecture that in both cases S will be transient, also because it is a 3-dimensional process. Proving this seems nontrivial, but notice that a possible intermediate step between the dimension 3 and 4 could be to consider the analogue problem on the discrete 3-dimensional Heisenberg group, which is generated by 2 elements (and their inverses), yet balls of radius r have size order r^4 .

Let us make some comments now on the 2-dimensional case. As a 2-dimensional process, we believe that $M(1, 1)$ is recurrent. Observe however that this is not true when starting from any configuration of visited sites. Indeed if we start with a vertical line of visited sites, then the process will be trapped in this line, and if the line does not include the origin, the process will not return there. It is also not difficult to construct starting environments such that the first coordinate of the process will tend almost surely toward $+\infty$. For example if the initial configuration is the “trumpet” $\{(x, y): |y| < e^x\}$ then the walker will drift to infinity in the x direction.¹ Of course it is not possible for the random walk to create these environments in finite time, so it is not an obstacle for recurrence, but it may be interesting to keep this in mind. Another problem concerns the limiting shape of the range (i.e. the set of visited sites) of the process. Based on heuristics and some simulations, we believe that it is a vertical interval. This problem is closely related to the question of evaluating the size of the range R_n at time n . Indeed the horizontal displacement of the process at time n is of order $\sqrt{\#R_n}$, whereas its vertical displacement is always of order $\sqrt{n}$. So another formulation of the problem would be to show that R_n is sublinear. By the way we mention a related question. Assume that at each step, one can decide, conditionally on the past, to move the first coordinate or the second coordinate (and then perform a 1-dimensional simple random walk step). Then what is the best strategy to maximize the range? In particular is it possible for the range to be of size roughly n , or at least significantly larger than $n/\ln n$, which is the size of the range of the simple random walk?

A possible generalization of our model would be to consider multi-excited versions, in the spirit of Zerner [13]. In this case one should first decompose d as $d = d_1 + \dots + d_m$, for some $m \geq 2$ and $d_i \geq 1$, with $i \leq m$. Then at i th visit to a site only the i th component of S performs a simple random walk step, if $i < m$, and at further visits only the m th component moves. In dimension 4 for instance the case $d_1 = 2$ and $d_2 = d_3 = 1$ seems interesting and nontrivial. Another interesting case is $d \geq 3$ and $d_i = 1$ for each $i \leq d$ (even the case d very large seems nontrivial).

A related problem which appeared in draft versions of this paper was solved by Y. Peres and P. Sousi (private communication) who proved, among other things, the following. Let μ_1 and μ_2 be any two symmetric laws on $\mathbb{Z}^d$, $d \geq 3$, and assume the support of μ_1 and μ_2 both generate $\mathbb{Z}^d$. Decide that at first visit to a site the jump of the process has law μ_1 , and at further visits it has law μ_2 . Then this process is transient.

2. Proof of the theorem

The theorem is a direct consequence of the following proposition:

Proposition 1. *There exists a constant $C > 0$ such that for any $n > 1$,*

$$\mathbb{P}[0 \in \{S_n, \dots, S_{2n}\}] \leq C \left(\frac{\ln \ln n}{\ln n} \right)^2.$$

¹ We will not prove any of these claims, as they are somewhat off-topic.

Indeed assuming this proposition we get

$$\sum_{k \geq 0} \mathbb{P}[0 \in \{S_{2k}, \dots, S_{2k+1}\}] < +\infty,$$

and we can conclude by using the Borel–Cantelli Lemma. So all we have to do is to prove this proposition.

Proof of Proposition 1. For any $n \geq 1$, denote by r_n the cardinality of the range of S at time n . The next lemma will be needed:

Lemma 1. For any $M > 0$, there exists a constant $C > 0$, such that

$$\mathbb{P}[n/(C \ln n)^2 \leq r_n \leq 99n/100] = 1 - o(n^{-M}).$$

Proof. Note first that for any k , if S_k and S_{k+1} were not already visited in the past, then $S_{k+2} = S_k$ with probability at least $1/4$. In particular for any k , there is probability at least $1/4$ for the event

$$E_k := \{S \text{ is not at a fresh site at one of the times } k, k+1 \text{ or } k+2\},$$

to occur. In other words, if $\varepsilon_k = 1_{E_k}$, then $\mathbb{E}[\varepsilon_{3k} \mid \varepsilon_0, \dots, \varepsilon_{3(k-1)}] \geq 1/4$ for all k . Then a standard use of the Azuma–Hoeffding inequality gives that

$$\mathbb{P}\left[\sum_{k=0}^{\lfloor n/3 \rfloor} \varepsilon_{3k} \leq n/100\right] = o(n^{-M}),$$

which gives the desired upper bound on r_n .

We now prove the lower bound. Let $c > 0$ be fixed. Let $(U_n, n \geq 0)$ be a simple random walk on $\mathbb{Z}^2$. For any $n \geq 1$ and $x \in \mathbb{Z}^2$, denote by $N_n(x)$ the number of visits of U to x before time n . A simple and standard calculation (see e.g. [9, Proposition 4.2.4]) shows that there exists a constant $C > 0$ such that the probability to not visit x in the next n steps after a given visit is $\geq C/\log n$. Using the strong Markov property one gets that the probability to make $k+1$ visits by time n is $\leq \exp(-Ck/\log n)$ and hence there exists some $C' > 0$ depending on M such that

$$\mathbb{P}[N_n(x) \geq C'(\ln n)^2] = o(n^{-M-2}).$$

Moreover since U makes nearest-neighbor jumps, before time n it stays in a ball of radius n . Thus if $N_n^* = \sup_x N_n(x)$, then

$$\mathbb{P}[N_n^* \geq C'(\ln n)^2] = n^2 \times o(n^{-M-2}) = o(n^{-M}).$$

Thus if $r_{n,U}$ is the size of the range of U at time n , we get

$$\mathbb{P}[r_{n,U} \leq n/(C'(\ln n)^2)] = o(n^{-M}).$$

Let's come back to the original process $S = (X, Y)$ now. We just observe that at time n one of the X or Y component performed $n/2$ steps. Since each of these components is a simple random walk, we deduce from the previous estimate, that before time n , X or Y will visit at least $n/(2C'(\ln n)^2)$ sites, with probability at least $1 - o(n^{-M})$. This gives the desired lower bound for r_n and concludes the proof of the lemma. $\square$

We can finish now the proof of Proposition 1. As noticed in the introduction, observe that the X and Y components are time changed simple random walks. Specifically we have the equality in law:

$$((X_k, Y_k), k \geq 1) = ((U(r_{k-1}), V(k - r_{k-1})), k \geq 1), \quad (1)$$

where U and V are two independent simple random walks on $\mathbb{Z}^2$, and where by abuse of notation we also denote by r_k the size of the range of the (U, V) process at k th step. More precisely r_k may be defined recursively by $r_0 = 1$ and for $k \geq 1$,

$$r_k := \#\{(U(0), V(0)), \dots, (U(r_{k-1}), V(k - r_{k-1}))\}.$$

The proof of (1) is immediate by induction.

By using now Lemma 1 and the independence of U and V , we get

$$\mathbb{P}[0 \in \{S_n, \dots, S_{2n}\}] \leq \mathbb{P}[0 \in \{U(n/(C \ln n)^2), \dots, U(2n)\}] \mathbb{P}[0 \in \{V(n/100), \dots, V(2n)\}] + o(n^{-M}).$$

Thus Proposition 1 follows from the following lemma:

Lemma 2. Let U be the simple random walk on $\mathbb{Z}^2$ and let $t \in [n/(\ln n)^3, 2n]$. Then

$$\mathbb{P}[0 \in \{U(t), \dots, U(2n)\}] = \mathcal{O}\left(\frac{\ln \ln n}{\ln n}\right). \quad (2)$$

Proof. This lemma is standard, but we give a proof for the reader's convenience. First let $|\cdot|$ denote the L^1 norm on $\mathbb{R}^2$ (i.e. $|(x_1, x_2)| = |x_1| + |x_2|$ for all $(x_1, x_2) \in \mathbb{R}^2$). Since $t \geq n/(\ln n)^3$, it is well known (see e.g. [3, Theorem 1, p. 542]) that

$$\mathbb{P}\left[|U(t)| \leq \frac{\sqrt{n}}{(\ln n)^3}\right] = \mathcal{O}((\ln n)^{-1}).$$

Moreover it is known (see e.g. [8, Proposition 1.6.7]) that

$$\mathbb{P}_x[\tau_0 < \tau_{|x|(\ln |x|)^4}] = \mathcal{O}\left(\frac{\ln \ln |x|}{\ln |x|}\right),$$

where $\mathbb{P}_x$ denotes the law of U starting from x and for any $r \geq 0$,

$$\tau_r = \inf\{k > 0: r < |U(k)| \leq r + 1\}.$$

But if $|x| \geq \sqrt{n}/(\ln n)^3$, then $\sqrt{n} \ln n = \mathcal{O}(|x|(\ln |x|)^4)$. On the other hand (see e.g. [9, Proposition 2.1.2])

$$\mathbb{P}[\tau_{\sqrt{n} \ln n} \leq 2n] = \mathcal{O}(n^{-1}).$$

Notice finally that if $|x| \geq \sqrt{n}/(\ln n)^3$, then

$$\frac{\ln \ln |x|}{\ln |x|} = \mathcal{O}\left(\frac{\ln \ln n}{\ln n}\right).$$

The lemma follows by using Markov's property at time $t = n/(\ln n)^3$,

$$\begin{aligned} \mathbb{P}[0 \in \{U(t), \dots, U(2n)\}] &\leq \mathbb{P}\left[|U(t)| \leq \frac{\sqrt{n}}{(\ln n)^3}\right] + \max_{|x| > \sqrt{n}/(\ln n)^3} \mathbb{P}_x(\tau_0 \leq 2n) \\ &\leq \mathcal{O}((\ln n)^{-1}) + \max_{|x| > \sqrt{n}/(\ln n)^3} \mathbb{P}_x[\tau_0 < \tau_{|x|(\ln |x|)^4}] + \mathbb{P}[\tau_{|x|(\ln |x|)^4} \leq 2n] \\ &\leq \mathcal{O}((\ln n)^{-1}) + \mathcal{O}\left(\frac{\ln \ln n}{\ln n}\right) + \mathcal{O}(n^{-1}) = \mathcal{O}\left(\frac{\ln \ln n}{\ln n}\right) \end{aligned}$$

as required. $\square$

The proof of Theorem 1 is now finished. $\square$

Remark 1. A slight modification of the proof actually shows that $M(2, 2)$ is transient for any finite initial configuration of visited sites. One just has to replace r_{k-1} in (1) by $r_{k-1} - n_{k-1}$, where n_k is the number of sites (different from the origin) which are considered as visited at time 0, and effectively visited by the (U, V) process in the k first steps. Since $\sup_k n_k$ is finite this does not change the rest of the proof. This is of course not always the case if this configuration is infinite. For instance if we decide that all sites of the form $(0, 0, *, *)$ are already visited at time 0, then the X component will never move and $M(2, 2)$ will not be transient.

Acknowledgements

This work was done while B.S. was a visitor at the Weizmann Institute and he thanks the institute for its kind hospitality.

References

- [1] G. Amir, I. Benjamini, G. Kozma, Excited random walk against a wall, *Probab. Theory Related Fields* 140 (2008) 83–102.
- [2] I. Benjamini, D.B. Wilson, Excited random walk, *Electron. Commun. Probab.* 8 (2003) 86–92 (electronic).
- [3] W. Feller, *An Introduction to Probability Theory and Its Applications*, vol. II, second ed., John Wiley & Sons, 1971.
- [4] M. Holmes, Excited against the tide: A random walk with competing drifts, preprint, arXiv:0901.4393.
- [5] H. Kesten, O. Raimond, Br. Schapira, Random walks with occasionally modified transition probabilities, preprint, arXiv:0911.3886.
- [6] E. Kosygina, M.P.W. Zerner, Positively and negatively excited random walks on integers, with branching processes, *Electron. J. Probab.* 13 (2008) 1952–1979.
- [7] G. Kozma, Problem session, in: *Non-classical interacting random walks*, Oberwolfach report 27/2007, http://www.mfo.de/programme/schedule/2007/21/OWR_2007_27.pdf.
- [8] G.F. Lawler, *Intersections of Random Walks*, Probab. Appl., Birkhäuser Boston, Inc., Boston, MA, 1991, 219 pp.
- [9] G.F. Lawler, V. Limic, *Random Walk: A Modern Introduction*, Cambridge Stud. Adv. Math., vol. 123, Cambridge Univ. Press, Cambridge, 2010.
- [10] M. Menshikov, S. Popov, A. Ramirez, M. Vachkovskaia, On a general many-dimensional excited random walk, preprint, arXiv:1001.1741.
- [11] F. Merkl, S.W.W. Rolles, Linearly edge-reinforced random walks, in: *IMS Lecture Notes, Monogr. Ser. Dynamics & Stochastics*, vol. 48, Inst. Math. Statist., Beachwood, OH, 2006, pp. 66–77.
- [12] R. Pemantle, A survey of random processes with reinforcement, *Probab. Surv.* 4 (2007) 1–79 (electronic).
- [13] M.P.W. Zerner, Multi-excited random walks on integers, *Probab. Theory Related Fields* 133 (2005) 98–122.