

From repeated to continuous quantum interactions

Stéphane ATTAL¹ and Yan PAUTRAT²

¹ Institut G. Desargues
Université Claude Bernard, Lyon I
21, av. Claude Bernard
69622 Villeurbanne cedex, France

² Laboratoire de Mathématiques
Université Paris-Sud
Bâtiment 425
91405 Orsay Cedex, France

Abstract

We consider the general physical situation of a quantum system \mathcal{H}_0 interacting with a chain of exterior systems $\bigotimes_{\mathbb{N}^*} \mathcal{H}$, one after the other, during a small interval of time \hbar and following some Hamiltonian H on $\mathcal{H}_0 \otimes \mathcal{H}$. We discuss the passage to the limit to continuous interactions ($\hbar \rightarrow 0$) in a setup which allows to compute the limit of this Hamiltonian evolution in a single state space: a continuous field of exterior systems $\bigotimes_{\mathbb{R}^+} \mathcal{H}$. Surprisingly, the passage to the limit shows the necessity for 3 different time scales in H . The limit evolution equation is shown to spontaneously produce quantum noises terms: we obtain a quantum Langevin equation as limit of the Hamiltonian evolution. For the very first time, these quantum Langevin equations are obtained as the effective limit from repeated to continuous interactions and not only as a model. These results justify the usual quantum Langevin equations considered in continual quantum measurement or in quantum optics. Physically, the typical hamiltonian allowing this passage to the limit shows up three different parts which correspond to the free evolution, an analogue of a weak coupling limit part and a scattering interaction part. We apply these results to give an Hamiltonian description of the von Neumann measurements. We also consider the approximation of continuous time quantum master equations by discrete time ones; in particular we show how any Lindblad generator is naturally obtained as the limit of completely positive maps.

Contents

I. Introduction

II. Discrete dynamics on the atom chain

II.1 Repeated quantum interactions

II.2 Structure of the atom chain

II.3 Unitary dilations of completely positive maps

III. From the atom chain to the atom field

III.1 Structure of the atom field

III.2 Quantum noises

III.3 Embedding and approximation by the atom chain

III.4 Quantum Langevin equations

IV. Convergence theorems

IV.1 Convergence to quantum Langevin equations

IV.2 Typical Hamiltonian

IV.3 Hamiltonian description of von Neumann measurements

IV.4 An example

IV.5 From completely positive maps to Lindbladians

I. Introduction

Quantum Langevin equations as a model for quantum open systems have been considered for at least 40 years (for example [FKM], [FLO], [AFL]). They have been given many different meanings in terms of several definitions of quantum noises or quantum Brownian motions (for example [G-Z], [H-P], [GSI]). One of the most developed and useful mathematical languages developed for that purpose is the quantum stochastic calculus of Hudson and Parthasarathy and their quantum stochastic differential equations ([H-P]). The quantum Langevin equations they allow to consider have been used very often to model typical situations of quantum open systems: continual quantum measurement ([Ba1], [B-B]), quantum optics ([F-R], [FRS] [Ba2]), electronic transport [BRSW], thermalization ([M-R], [L-M]), etc.

The physical justification for considering such quantum Langevin equations is often given in terms of some particular approximations of the true Hamiltonian interaction dynamics: rotating wave approximation, Markov approximation, large band approximation (cf [G-Z] chapter 11).

They are also often justified as natural dilations of quantum master equations on the small system. That is, for any (good) semigroup of completely positive maps on the small system (with Lindblad generator \mathcal{L}), one can dilate the small system with an appropriate Fock space and obtain an explicit quantum stochastic differential equation on the whole space. The unique solution of this equation is a unitary evolution in interaction picture such that the trace on the small system of the induced evolution yields the original semigroup. This corresponds, at the quantum level, to the well-known way of realizing a concrete Markov process from

a given semigroup (or generator) by adding a noise space to the (classical) system space and solving an adequate stochastic differential equation.

Some quantum stochastic differential equations have also been obtained in the so-called *stochastic limit* from explicit Hamiltonian dynamics ([A-L], [AGL], [ALV]). These results show some similarity with the ones described here, but they are actually quite different in the type of limit considered. In those articles the weak coupling regime and the convergence is convergence of correlation functions of processes living in different spaces; we study a continuous limit and obtain weak or strong convergence of operators in a single here.

In this article we consider the effective Hamiltonian dynamics describing the repeated interactions, during short time intervals of length h , of a small system \mathcal{H}_0 with a chain of exterior systems $\otimes_{\mathbb{N}^*} \mathcal{H}$. We embed all these chains as particular subspaces, attached to the parameter h , of a continuous field

$$\bigotimes_{\mathbb{R}^+} \mathcal{H}$$

in such a way that the subspaces associated to the chain increase and fill the field when h tends to 0. This framework may seem to specialize to the case of a zero-temperature exterior system; actually, it also applies to the case of positive temperature, using the cyclic (GNS) representation of the given state (see [AJ2]).

By developing an appropriate language for the chain $\otimes_{\mathbb{N}^*} \mathcal{H}$ and for the field $\bigotimes_{\mathbb{R}^+} \mathcal{H}$ and by describing the discrete time Hamiltonian evolution generated by the repeated interactions, we are able to pass to the limit when $h \rightarrow 0$ and to prove that the limit evolution equation is the solution of a quantum stochastic differential equation. This limit is obtained in the strong topology of operators and in a single space: the continuous field $\bigotimes_{\mathbb{R}^+} \mathcal{H}$, and implies the weak convergence of the Heisenberg evolutions of any observable.

Of course, such a limit cannot be obtained without assumptions on the elementary interaction Hamiltonian H . This is similar to the central limit theorem: a random walk gives a trivial limit when its time step h goes to zero and it is only when suitably renormalized (by a factor \sqrt{h}) that it yields a Gaussian. Other normalizations give either trivial limits or no limit at all. In our Hamiltonian context the situation is going to be the same. For a non-trivial limit of these repeated interactions to exist, we will need the Hamiltonian H to satisfy some renormalization properties. The surprise here is that the necessary renormalization factor is not global, it is different following some parts of the Hamiltonian operator. We identify three different time scales in H : one of order 1, one of order \sqrt{h} , one of order h .

We describe a large class of Hamiltonians which seems to be typical for the above conditions to be satisfied. These typical Hamiltonians are a combination of a free evolution part and two parts which are reminiscent of a weak coupling limit-like interaction part and a scattering-like interaction part. The precise relation of our limits with the more classical weak coupling is studied in [AJ1]. Another way

to interpret these different time scales is to see them as a renormalization of the field operators of the photon chain. In the limit we observe a quantum Langevin which shows up both diffusion-like terms and Poisson-like terms. The combination of the two limits here shows an effective Hamiltonian for the small system which contains the surprising term

$$V^*D^{-2}(\sin D - D)V$$

which comes from the simultaneous presence of all the different parts of the Hamiltonian. Notice a possible extension of our results: only the case of time-independent coupling is discussed here but results for time-dependent ones can easily be deduced from our proofs.

This article is structured as follows. In section II we present the exact mathematical model of repeated quantum interactions and end up with the associated evolution equation (subsection II.1). We then introduce a mathematical setup for the study of the space $\bigotimes_{N^*} \mathcal{H}$ which will help much for passing to the continuous field. In particular this includes a particular choice of an orthonormal basis of the phase space and a particular choice of a basis for the operators on that phase space (subsection II.2). Finally we show how the typical evolution equations obtained in II.1 are the general model for the unitary dilation of any given discrete semigroup of completely positive maps (subsection II.3).

Section III is devoted to presenting the whole formalism of the continuous atom field. In subsection III.1 we present the space which is candidate for representing the continuous field limit of the atom chain. It is actually a particular Fock space on which we develop an unusual structure which shows up the required properties. In subsection III.2 we present the natural quantum noises on the continuous field and the associated quantum stochastic integrals, the quantum Ito formula and the quantum stochastic differential equations. In subsection III.3 we concretely realize the atom chain of section II as a strict subspace of the atom field. Not only do we realize it as a subspace, but also realize the action of its basic operators inside the atom field. All these atom chain subspaces are related to a partition of \mathbb{R}^+ . When the diameter of the partition goes to 0, we show that the corresponding subspace completely fills the continuous field and the basic operators of the chain converge to the quantum noises of the field (with convenient normalizations). Finally, considering the projection of the continuous atom field onto an atom chain subspace, we state a formula for the projection of a general quantum stochastic integral.

In section IV all the pieces of the puzzle fit together. By computing the projection on the atom chain of a quantum stochastic differential equation we show that the typical evolution equation of repeated interactions converges in the field space to the solution of a quantum Langevin equation, assuming the fact that the associated Hamiltonian satisfies some particular renormalization property corresponding to three different time scales. It is to that result and to some of its extensions that subsection IV.1 is devoted. In subsection IV.2 we describe a family of Hamiltonians which seems to be typical of the conditions obtained above. We

show that this family of Hamiltonians describes altogether free evolution, weak coupling limit-like and scattering-like terms. Computing the associated quantum Langevin equation at the limit, we obtain an effective Hamiltonian on \mathcal{H}_0 which contains a new term. This new term appears only when weak coupling and low density limits terms are in presence together in the Hamiltonian. In subsection IV.3, we apply these results to describe the von Neumann measurement apparatus in the Hamiltonian framework of repeated quantum interactions. In subsection IV.4 we explicitly compute a simple example. In subsection IV.5 we show that our approximation theorem puts into evidence a natural way that completely positive maps have to converge to Lindblad generators.

II. Discrete dynamics on the atom chain

II.1 Repeated quantum interactions

Here we give a precise description of our physical model: repeated quantum interactions. All Hilbert spaces that we consider in the sequel are supposed to be separable.

We consider a small quantum system \mathcal{H}_0 and another quantum system \mathcal{H} which represents either a piece of environment, a measuring apparatus or incoming photons . . . We consider the space $\mathcal{H}_0 \otimes \mathcal{H}$ in order to couple the two systems, an Hamiltonian H on $\mathcal{H}_0 \otimes \mathcal{H}$ which describes the interaction and the associated unitary evolution during the interval $[0, h]$ of time:

$$\mathbb{L} = e^{-ihH}.$$

This single interaction is therefore described in the Schrödinger picture by

$$\rho \mapsto \mathbb{L} \rho \mathbb{L}^*$$

and in the Heisenberg picture by

$$X \mapsto \mathbb{L}^* X \mathbb{L}.$$

Now, after this first interaction, we repeat it but this time coupling the same \mathcal{H}_0 with a new copy of \mathcal{H} . This means that the new copy was kept isolated until then; similarly the previously considered copy of \mathcal{H} will remain isolated for the rest of the experience. One can think of many physical examples where this situations arises: in repeated quantum measurement where a family of identical measurement devices are presented one after the other before the system (or a single device is refreshed after every use), in quantum optics where a sequence of independent atoms arrives one after the other to interact with a field in some cavity for a short time. More generally it can be seen as a good model if it is assumed that perturbations in \mathcal{H} due to the interaction are dissipated after every time h .

The sequence of interactions can be described in the following way: the state space for the whole system is

$$\mathcal{H}_0 \otimes \bigotimes_{\mathbb{N}^*} \mathcal{H}$$

Index for a few lines only the copies of \mathcal{H} as $\mathcal{H}_1, \mathcal{H}_2, \dots$. Define then a unitary operator \mathbb{L}_n as the canonical ampliation to $\mathcal{H}_0 \otimes \mathcal{H}_1 \otimes \mathcal{H}_2 \otimes \dots$ of the operator which acts as \mathbb{L} on $\mathcal{H}_0 \otimes \mathcal{H}_n$; that is, \mathbb{L}_n acts as the identity on copies of \mathcal{H} other than \mathcal{H}_n .

The effect of the n -th interaction in the Schrödinger picture writes then

$$\rho \mapsto \mathbb{L}_n \rho \mathbb{L}_n^*,$$

for every density matrix ρ , so that the effect of the n first interactions is

$$\rho \mapsto u_n \rho u_n^*$$

where $(u_n)_{n \in \mathbb{N}}$ is a sequence in $\mathcal{B}(\mathcal{H}_0 \otimes \bigotimes_{\mathbb{N}^*} \mathcal{H})$ which satisfies the equations

$$\begin{cases} u_{n+1} = \mathbb{L}_{n+1} u_n \\ u_0 = I. \end{cases} \quad (1)$$

It is evolution equations such as (1) that we are going to study in this article.

II.2 Structure of the atom chain

We here describe some useful mathematical structure on the space $\bigotimes_{\mathbb{N}^*} \mathcal{H}$ which will constitute the main ingredient of our approach.

Let us fix a particular Hilbertian basis $(X^i)_{i \in \Lambda \cup \{0\}}$ for the Hilbert space \mathcal{H} , where we assume (for notational purposes) that $0 \notin \Lambda$. This particular choice of notations is motivated by physical interpretations: indeed, we see the X^i , $i \in \Lambda$, as representing for example the different possible excited states of an atom. The vector X^0 represents the “ground state” or “vacuum state” of the atom and will usually be denoted Ω .

Let $\mathbb{T}\Phi$ be the tensor product $\bigotimes_{\mathbb{N}^*} \mathcal{H}$ with respect to the stabilizing sequence Ω . In other words, this means simply that an orthonormal basis of $\mathbb{T}\Phi$ is given by the family

$$\{X_A; A \in \mathcal{P}_{\mathbb{N}^*, \Lambda}\}$$

where

- the set $\mathcal{P}_{\mathbb{N}^*, \Lambda}$ is the set of finite subsets

$$\{(n_1, i_1), \dots, (n_k, i_k)\}$$

of $\mathbb{N}^* \times \Lambda$ such that the n_i 's are mutually different. Another way to describe the set $\mathcal{P}_{\mathbb{N}^*, \Lambda}$ is to identify it to the set of sequences $(A_n)_{n \in \mathbb{N}^*}$ with values in $\Lambda \cup \{0\}$ which take a value different from 0 only finitely often.

- X_A denotes the vector

$$\Omega \otimes \dots \otimes \Omega \otimes X^{i_1} \otimes \Omega \otimes \dots \otimes \Omega \otimes X^{i_2} \otimes \dots$$

where X^{i_1} appears in n_1 -th copy of $\mathcal{H} \dots$

The physical signification of this basis is easy to understand: we have a chain of atoms, indexed by \mathbb{N}^* . The space $\mathbb{T}\Phi$ is the state space of this chain, the vector X_A with $A = \{(n_1, i_1), \dots, (n_k, i_k)\}$ representing the state in which exactly

k atoms are excited: atom n_1 in the state i_1 , etc, all other atoms being in the ground state.

This particular choice of a basis gives $\mathbb{T}\Phi$ a particular structure. If we denote by $\mathbb{T}\Phi_{[n]}$ the space generated by the X_A such that $A \subset \{1, \dots, n\} \times \Lambda$ and by $\mathbb{T}\Phi_{[m]}$ the one generated by the X_A such that $A \subset \{m, m+1, \dots\} \times \Lambda$, we get an obvious natural isomorphism between $\mathbb{T}\Phi$ and $\mathbb{T}\Phi_{[n-1]} \otimes \mathbb{T}\Phi_{[n]}$ given by

$$[f \otimes g](A) = f(A \cap \{1, \dots, n-1\} \times \Lambda) g(A \cap \{n, \dots\} \times \Lambda).$$

Put $\{a_j^i; i, j \in \Lambda \cup \{0\}\}$ to be the natural basis of $\mathcal{B}(\mathcal{H})$, that is,

$$a_j^i(X^k) = \delta_{ik} X^j.$$

We denote by $a_j^i(n)$ the natural ampliation of the operator a_j^i to $\mathbb{T}\Phi$ which acts on the copy number n as a_j^i and the identity elsewhere. That is, in terms of the basis X_A ,

$$a_j^i(n)X_A = \mathbb{1}_{(n,i) \in A} X_{A \setminus (n,i) \cup (n,j)}$$

if neither i nor j is zero, and

$$a_0^i(n)X_A = \mathbb{1}_{(n,i) \in A} X_{A \setminus (n,i)},$$

$$a_j^0(n)X_A = \mathbb{1}_{(n,0) \in A} X_{A \cup (n,j)},$$

$$a_0^0(n)X_A = \mathbb{1}_{(n,0) \in A} X_A,$$

where $(n, 0) \in A$ actually means “for any i in Λ , $(n, i) \notin A$ ”.

II.3 Unitary dilation of completely positive maps

The evolution equations

$$u_n = \mathbb{I}L_n \dots \mathbb{I}L_1$$

obtained in the physical setup of repeated quantum interactions are actually of mathematical interest on their own for they provide a canonical way of dilating discrete semigroups of completely positive maps into unitary automorphisms.

The mathematical setup is the same. Let $\mathbb{I}L$ be any operator on $\mathcal{H}_0 \otimes \mathcal{H}$. Let $\mathbb{T}\Phi = \otimes_{\mathbb{N}^*} \mathcal{H}$ and $(\mathbb{I}L_n)_{n \in \mathbb{N}^*}$ be defined as in the above section. We then consider the associated evolution equations

$$u_n = \mathbb{I}L_n \dots \mathbb{I}L_1 \tag{1}$$

with $u_0 = I$.

The following result is obvious.

Proposition 1. – *The solution $(u_n)_{n \in \mathbb{N}}$ of (1) is made of unitary (resp. isometric, contractive) operators if and only if $\mathbb{I}L$ is unitary (resp. isometric, contractive).* ■

Note that if $\mathbb{I}L$ is unitary, then the mappings

$$j_n(H) = u_n^* H u_n$$

are automorphisms of $\mathcal{B}(\mathcal{H}_0 \otimes \mathcal{H})$.

Let \mathbb{E}_0 be the partial trace on \mathcal{H}_0 defined by

$$\langle \phi, \mathbb{E}_0(H) \psi \rangle = \langle \phi \otimes \Omega, H \psi \otimes \Omega \rangle$$

for all $\phi, \psi \in \mathcal{H}_0$ and every operator H on $\mathcal{H}_0 \otimes \mathbb{T}\Phi$.

Unitary dilations of completely positive semigroups are obtained in the following theorem. Recall that, by Kraus' theorem, any completely positive operator ℓ on $\mathcal{B}(\mathcal{H}_0)$ is of the form

$$\ell(X) = \sum A_i^* X A_i$$

where the summation ranges over $\Lambda \cup \{0\}$, the A_i are bounded operators and the sum is strongly convergent. Conversely, any such operator is completely positive.

Remark: Of course the Kraus form of an operator is *a priori* indifferent to the specificity of the value $i = 0$. The special role played by one of the indices will appear later on.

Theorem 2. – *Let \mathbb{I} be any unitary operator on $\mathcal{H}_0 \otimes \mathcal{H}$. Consider the coefficients $(\mathbb{I}_j^i)_{i,j \in \Lambda \cup \{0\}}$, which are operators on \mathcal{H}_0 , of the matrix representation of \mathbb{I} in the basis $\Omega, X^i, i \in \Lambda$ of \mathcal{H} .*

Then, for any $X \in \mathcal{B}(\mathcal{H}_0)$ we have

$$\mathbb{E}_0[j_n(X \otimes I)] = \ell^n(X)$$

where ℓ is the completely positive map on $\mathcal{B}(\mathcal{H}_0)$ given by

$$\ell(X) = \sum_{i \in \Lambda \cup \{0\}} (\mathbb{I}_i^0)^* X \mathbb{I}_i^0.$$

Conversely, consider any completely positive map

$$\ell(X) = \sum_{i \in \Lambda \cup \{0\}} A_i^* X A_i$$

on $\mathcal{B}(\mathcal{H}_0)$ such that $\ell(I) = I$. Then there exists a unitary operator \mathbb{I} on $\mathcal{H}_0 \otimes \mathcal{H}$ such that the associated unitary family of automorphisms

$$j_n(H) = u_n^* H u_n$$

satisfies

$$\mathbb{E}_0[j_n(X \otimes I)] = \ell^n(X),$$

for all $n \in \mathbb{N}$.

Proof

Consider $\mathbb{I} = (\mathbb{I}_j^i)_{i,j \in \Lambda \cup \{0\}}$ such as in the above statements. Consider the unitary family

$$u_n = \mathbb{I}_n \dots \mathbb{I}_1.$$

Note that

$$u_{n+1} = \mathbb{I}_{n+1} u_n.$$

Put $j_n(H) = u_n^* H u_n$ for every operator H on $\mathcal{H}_0 \otimes \mathcal{H}$. Then, for any operator X on \mathcal{H}_0 we have

$$j_{n+1}(X \otimes I) = u_n^* \mathbb{L}_{n+1}^*(X \otimes I) \mathbb{L}_{n+1} u_n.$$

When considered as a matrix of operators on \mathcal{H}_0 , in the basis $\Omega, X^i, i \in \Lambda$ of \mathcal{H} , the matrix associated to $X \otimes I$ is of diagonal form. We get

$$\begin{aligned} \mathbb{L}_{n+1}^*(X \otimes I) \mathbb{L}_{n+1} &= \\ &= \begin{pmatrix} (\mathbb{L}_0^0)^* & (\mathbb{L}_1^0)^* & \cdots \\ (\mathbb{L}_0^1)^* & (\mathbb{L}_1^1)^* & \cdots \\ \vdots & \vdots & \ddots \end{pmatrix} \begin{pmatrix} X & 0 & \cdots \\ 0 & X & \cdots \\ \vdots & \vdots & \ddots \end{pmatrix} \begin{pmatrix} \mathbb{L}_0^0 & \mathbb{L}_1^0 & \cdots \\ \mathbb{L}_0^1 & \mathbb{L}_1^1 & \cdots \\ \vdots & \vdots & \ddots \end{pmatrix} \end{aligned}$$

which is the matrix $\mathbf{L}_{n+1}(X) = (B_j^i(X))_{i,j \in \Lambda \cup \{0\}}$ with

$$B_j^i(X) = \sum_{k \in \Lambda \cup \{0\}} (\mathbb{L}_k^j)^* X \mathbb{L}_k^i.$$

Note that the operator $\mathbf{L}_{n+1}(X)$ acts non trivially only on the tensor product of \mathcal{H}_0 with the $(n+1)$ -th copy of \mathcal{H} . When represented as an operator on

$$\mathcal{H}_0 \otimes \mathbb{T}\Phi_{n+1}] = (\mathcal{H}_0 \otimes \mathbb{T}\Phi_n]) \otimes \mathcal{H}$$

as a matrix with coefficients in $\mathcal{B}(\mathcal{H}_0 \otimes \mathbb{T}\Phi_n])$ it writes exactly in the same way as above, just replacing $B_j^i(X)$ (which belongs to $\mathcal{B}(\mathcal{H}_0)$) by

$$B_j^i(X) \otimes I_{|\mathbb{T}\Phi_n]}.$$

Also note that, as can be proved by an easy induction, the operator u_n acts on $\mathcal{H}_0 \otimes \mathbb{T}\Phi_n]$ only. As an operator on $\mathcal{H}_0 \otimes \mathbb{T}\Phi_{n+1}]$ it is represented by a diagonal matrix. Thus $j_{n+1}(X) = u_n^* \mathbf{L}_{n+1}(X) u_n$ can be written on $\mathcal{H}_0 \otimes \mathbb{T}\Phi_{(n+1)}] = \mathcal{H}_0 \otimes \mathbb{T}\Phi_n] \otimes \mathcal{H}$ as a matrix of operators on $\mathcal{H}_0 \otimes \mathbb{T}\Phi_n]$ by

$$(j_{n+1}(X \otimes I))_j^i = j_n(B_j^i(X) \otimes I).$$

Note that $B_0^0(X) = \sum_{i \in \Lambda \cup \{0\}} (\mathbb{L}_i^0)^* X \mathbb{L}_i^0$ which is the mapping $\ell(X)$ of the statement.

Put $T_n(X) = \mathbb{E}_0[j_n(X \otimes I)]$. We have, for all $\phi, \psi \in \mathcal{H}_0$

$$\begin{aligned} \langle \phi, T_{n+1}(X) \psi \rangle &= \langle \phi \otimes \Omega, j_{n+1}(X \otimes I) \psi \otimes \Omega \rangle \\ &= \langle \phi \otimes \Omega, (j_n(B_j^i(X) \otimes I))_{i,j} \psi \otimes \Omega \rangle \\ &= \langle \phi \otimes \Omega_{\mathbb{T}\Phi_n]} \otimes \Omega_{\mathcal{H}}, (j_n(B_j^i(X) \otimes I))_{i,j} \psi \otimes \Omega_{\mathbb{T}\Phi_n]} \otimes \Omega_{\mathcal{H}} \rangle \\ &= \langle \phi \otimes \Omega_{\mathbb{T}\Phi_n]}, j_n(B_0^0(X) \otimes I) \psi \otimes \Omega_{\mathbb{T}\Phi_n]} \rangle \\ &= \langle \phi, T_n(\ell(X)) \psi \rangle. \end{aligned}$$

This proves that $T_{n+1}(X) = T_n(\ell(X))$ and the first part of the theorem is proved.

Conversely, consider a decomposition of a completely positive map ℓ of the form

$$\mathcal{L}(X) = \sum_{i \in \Lambda \cup \{0\}} A_i^* X A_i$$

for a family $(A_i)_{i \in \Lambda \cup \{0\}}$ of bounded operators on \mathcal{H}_0 such that

$$\sum_{i \in \Lambda \cup \{0\}} A_i^* A_i = I.$$

We claim that there exists a unitary operator \mathbb{L} on $\mathcal{H}_0 \otimes \mathcal{H}$ of the form

$$\mathbb{L} = \begin{pmatrix} A_0 & \dots & \dots \\ A_1 & \dots & \dots \\ \vdots & \vdots & \ddots \end{pmatrix}.$$

Indeed, the condition $\sum_{i \in \Lambda \cup \{0\}} A_i^* A_i = I$ guarantees that the first columns of \mathbb{L} are made of orthonormal vectors of $\mathcal{H}_0 \otimes \mathcal{H}$. We can thus complete the matrix by completing it into an orthonormal basis of $\mathcal{H}_0 \otimes \mathcal{H}$. This makes out a unitary matrix \mathbb{L} the coefficients of which we denote by $(A_j^i)_{i,j \in \Lambda \cup \{0\}}$. Note that $A_i^0 = A_{i+1}$. We now conclude easily by the first part of the theorem. ■

III From the atom chain to the atom field

III.1 Structure of the atom field

We now describe the structure of the continuous version of $\mathbb{T}\Phi$. The structure we are going to present here is rather original and not much expanded in the literature. It is very different from the usual presentation of quantum stochastic calculus ([H-P]), but it actually constitutes a very natural language for our purpose: approximation of the atom field by atom chains. This approach is taken from [At1]. We first start with a heuristical discussion. By a continuous version of the atom chain $\mathbb{T}\Phi$ we mean a Hilbert space with a structure which makes it the space

$$\Phi = \bigotimes_{\mathbb{R}^+} \mathcal{H}.$$

We have to give a meaning to the above notation. This could be achieved by invoking the framework of continuous tensor products of Hilbert spaces (see [Gui]), but we prefer to give a self-contained presentation which fits better with our approximation procedure.

Let us make out an idea of what it should look like by mimicking, in a continuous time version, what we have described in $\mathbb{T}\Phi$.

The countable orthonormal basis X_A , $A \in \mathcal{P}_{\mathbb{N}^*, \Lambda}$ is replaced by a continuous orthonormal basis $d\chi_\sigma$, $\sigma \in \mathcal{P}_{\mathbb{R}, \Lambda}$, where $\mathcal{P}_{\mathbb{R}, \Lambda}$ is the set of finite subsets of $\mathbb{R}^+ \times \Lambda$. With the same idea as for $\mathbb{T}\Phi$, this means that each copy of \mathcal{H} is equipped with an orthonormal basis $(\Omega, d\chi_t^i; i \in \Lambda)$ (where t is the parameter attached to the copy we are looking at). The orthonormal basis above is just the one obtained by specifying a finite number of sites t_1, \dots, t_n which are going to be excited, the other ones being supposed to be in the fundamental state Ω , and by specifying their level of excitation.

The representation of an element f of $\mathbb{T}\Phi$:

$$f = \sum_{A \in \mathcal{P}_{\mathbb{N}^*, \Lambda}} f(A) X_A$$

$$\|f\|^2 = \sum_{A \in \mathcal{P}_{\mathbb{N}^*, \Lambda}} |f(A)|^2$$

is replaced by an integral version of it in Φ :

$$f = \int_{\mathcal{P}_{\mathbb{R}, \Lambda}} f(\sigma) d\chi_\sigma$$

$$\|f\|^2 = \int_{\mathcal{P}_{\mathbb{R}, \Lambda}} |f|^2 d\sigma.$$

This last integral has to be explained: the measure $d\sigma$ is a ‘‘Lebesgue measure’’ on $\mathcal{P}_{\mathbb{R}, \Lambda}$, as will be explained later. From now on, the notation \mathcal{P} will denote, depending on the context, spaces of the type $\mathcal{P}_{\mathbb{N}^*, \Lambda}$ or $\mathcal{P}_{\mathbb{R}, \Lambda}$.

A good basis of operators acting on Φ can be obtained by mimicking the operators $a_j^i(n)$ of $\mathbb{T}\Phi$. We will here have a set of infinitesimal operators $da_j^i(t)$, $i, j \in \Lambda \cup \{0\}$, acting on the ‘‘t-th’’ copy of \mathcal{H} by:

$$da_0^0(t) d\chi_\sigma = d\chi_\sigma dt \mathbb{1}_{(t,0) \in \sigma}$$

$$da_i^0(t) d\chi_\sigma = d\chi_{\sigma \cup \{(t,i)\}} \mathbb{1}_{(t,0) \in \sigma}$$

$$da_0^i(t) d\chi_\sigma = d\chi_{\sigma \setminus \{(t,i)\}} dt \mathbb{1}_{(t,i) \in \sigma}$$

$$da_j^i(t) d\chi_\sigma = d\chi_{\sigma \setminus \{(t,i)\} \cup \{(t,j)\}} \mathbb{1}_{(t,i) \in \sigma}$$

for all $i, j \in \Lambda$. Similarly as in the discrete-time case, $(t, 0) \in \sigma$ means ‘‘for any i in Λ , $(t, i) \notin \sigma$ ’’.

We shall now describe a rigorous setup for the above heuristic discussion. We recall the structure of the bosonic Fock space Φ and its basic structure (cf [At1] for more details and [At3] for a complete study of the theory and its connections with classical stochastic processes).

Let \mathcal{H} be, as before, a Hilbert space with an orthonormal basis X^i , $i \in \Lambda \cup \{0\}$ and let \mathcal{H}' be the closed subspace generated by vectors X^i , $i \in \Lambda$ (or simply said, the orthogonal of X^0).

Let $\Phi = \Gamma_s(L^2(\mathbb{R}^+, \mathcal{H}'))$ be the symmetric (bosonic) Fock space over the space $L^2(\mathbb{R}^+, \mathcal{H}')$. We shall here give a very efficient presentation of that space, the so-called *Guichardet interpretation* of the Fock space.

Let \mathcal{P} ($= \mathcal{P}_{\mathbb{R}, \Lambda}$) be the set of finite subsets $\{(s_1, i_1), \dots, (s_n, i_n)\}$ of $\mathbb{R}^+ \times \Lambda$ such that the s_i are two by two different. Then $\mathcal{P} = \cup_n \mathcal{P}_n$ where \mathcal{P}_n is the subset of \mathcal{P} made of n -elements subsets of $\mathbb{R}^+ \times \Lambda$. By ordering the \mathbb{R}^+ -part of the elements of $\sigma \in \mathcal{P}_n$, the set \mathcal{P}_n can be identified to the increasing simplex $\Sigma_n = \{0 < t_1 < \dots < t_n\} \times \Lambda$ of $\mathbb{R}^n \times \Lambda$. Thus \mathcal{P}_n inherits a measured space structure from the Lebesgue measure on \mathbb{R}^n times the counting measure on Λ . This also gives a measure structure on \mathcal{P} if we specify that on $\mathcal{P}_0 = \{\emptyset\}$ we put the

measure δ_\emptyset . Elements of \mathcal{P} are often denoted by σ , the measure on \mathcal{P} is denoted $d\sigma$. The σ -field obtained this way on \mathcal{P} is denoted \mathcal{F} .

We identify any element $\sigma \in \mathcal{P}$ with a family $\{\sigma_i, i \in \Lambda\}$ of (mutually disjoint) subsets of \mathbb{R}^+ where

$$\sigma_i = \{s \in \mathbb{R}^+; (s, i) \in \sigma\}.$$

The *Fock space* Φ is the space $L^2(\mathcal{P}, \mathcal{F}, d\sigma)$. An element f of Φ is thus a measurable function $f : \mathcal{P} \rightarrow \mathbb{C}$ such that

$$\|f\|^2 = \int_{\mathcal{P}} |f(\sigma)|^2 d\sigma < \infty.$$

One can define, in the same way, $\mathcal{P}_{[a,b]}$ and $\Phi_{[a,b]}$ by replacing \mathbb{R}^+ with $[a, b] \subset \mathbb{R}^+$. These spaces are canonically identified to subspaces of Φ by identifying $\Phi_{[a,b]}$ with

$$\{f \in \Phi; f(\sigma) = 0 \text{ unless } \sigma \subset [a, b]\}.$$

As in discrete time, there is a natural isomorphism between $\Phi_{[0,t]} \otimes \Phi_{[t,+\infty[}$ given by $h \otimes g \mapsto f$ where $f(\sigma) = h(\sigma \cap [0, t] \times \Lambda)g(\sigma \cap (t, +\infty[\times \Lambda)$. This isomorphism is the usual “exponential property” of Fock spaces.

We shall use the following notations:

$$\Phi_t = \Phi_{[0,t]}, \quad \Phi_{[t} = \Phi_{[t,+\infty[}.$$

Define Ω to be the *vacuum vector*, that is, $\Omega(\sigma) = \delta_\emptyset(\sigma)$.

We now define a particular family of curves in Φ , which is going to be of great importance here. Define $\chi_t^i \in \Phi$ by

$$\chi_t^i(\sigma) = \begin{cases} \mathbb{1}_{[0,t]}(s) & \text{if } \sigma = \{(s, i)\} \\ 0 & \text{otherwise.} \end{cases}$$

Then notice that for all $t \in \mathbb{R}^+$ we have that χ_t^i belongs to $\Phi_{[0,t]}$. We actually have much more than that: we have

$$\chi_t^i - \chi_s^i \in \Phi_{[s,t]} \text{ for all } s \leq t.$$

This last property can be checked immediately from the definitions, and it is going to be of great importance in our construction. Also notice that χ_t^i and χ_s^j are orthogonal elements of Φ as soon as $i \neq j$. As we will see later on, apart from trivialities, the curves $(\chi_t^i)_{t \geq 0}$ are the only ones to share these properties.

These properties allow to define the so-called *Ito integral* on Φ . Indeed, let $g = \{(g_t^i)_{t \geq 0}, i \in \Lambda\}$ be families of elements of Φ indexed by both \mathbb{R}_+ and Λ , such that

- i) $t \mapsto \|g_t^i\|$ is measurable, for all i ,
- ii) $g_t^i \in \Phi_{[0,t]}$ for all t ,
- iii) $\sum_{i \in \Lambda} \int_0^\infty \|g_t^i\|^2 dt < \infty$

then one says that g is *Ito integrable* and we define its *Ito integral*

$$\sum_{i \in \Lambda} \int_0^\infty g_t^i d\chi_t^i$$

to be the limit in Φ of

$$\sum_{i \in \Lambda} \sum_{j \in \mathcal{N}} \tilde{g}_{t_j}^i \otimes (\chi_{t_{j+1}}^i - \chi_{t_j}^i) \quad (2)$$

where $\mathcal{S} = \{t_j, j \in \mathcal{N}\}$ is a partition of \mathbb{R}^+ which is understood to be refining and to have its diameter tending to 0, and $(\tilde{g}_t^i)_i$ is an Ito integrable family in Φ , such that for each i , $t \mapsto \tilde{g}_t^i$ is a step process, and which converges to $(g^i)_i$ in $L^2(\mathbb{R}^+ \times \mathcal{P})$.

Note that by assumption we always have that $\tilde{g}_{t_j}^i$ belongs to $\Phi_{[t_j, t_{j+1}]}$ and $\chi_{t_{j+1}}^i - \chi_{t_j}^i$ belongs to $\Phi_{[t_j, t_{j+1}]}$, hence the tensor product symbol in (2).

Also note that, as an example, one can take

$$\tilde{g}_t^i = \frac{1}{t_{j+1} - t_j} \int_{t_j}^{t_{j+1}} P_{t_j} g_s^i ds$$

where P_t is the orthogonal projection onto $\Phi_{[0, t]}$.

One then obtains the following properties ([At1], Proposition 1.4), where the notation $\vee \sigma$ means $\max \sigma$:

Theorem 3. – *The Ito integral $I(g) = \sum_i \int_0^\infty g_t^i d\chi_t^i$, of an Ito integrable family $g = (g^i)_{i \in \Lambda}$, is the element of Φ given by*

$$I(g)(\sigma) = \begin{cases} 0 & \text{if } \sigma = \emptyset \\ g_{\vee \sigma}^i(\sigma \setminus (\vee \sigma, i)) & \text{if } \vee \sigma \in \sigma_i. \end{cases}$$

It satisfies the Ito isometry formula:

$$\|I(g)\|^2 = \left\| \sum_{i \in \Lambda} \int_0^\infty g_t^i d\chi_t^i \right\|^2 = \sum_{i \in \Lambda} \int_0^\infty \|g_t^i\|^2 dt. \quad (3)$$

■

In particular, consider a family $f = (f^i)_{i \in \Lambda}$ which belongs to $L^2(\mathcal{P}_1) = L^2(\mathbb{R}^+ \times \Lambda)$, then the family $(f^i(t)\Omega)$, $t \in \mathbb{R}^+$, $i \in \Lambda$, is clearly Ito integrable. Computing its Ito integral we find that

$$I(f) = \sum_{i \in \Lambda} \int_0^\infty f^i(t)\Omega d\chi_t^i$$

is the element of the first particle space of the Fock space Φ associated to the function f , that is,

$$I(f)(\sigma) = \begin{cases} f^i(s) & \text{if } \sigma = \{s\}_i \\ 0 & \text{otherwise.} \end{cases}$$

Let us define the “adjoint” mapping of the Ito integral. For all $f \in \Phi$, all i in Λ and all $t \in \mathbb{R}^+$, consider the following mapping on \mathcal{P} :

$$[D_t^i f](\sigma) = f(\sigma \cup \{(t, i)\}) \mathbb{1}_{\sigma \subset [0, t[}.$$

We then have the following result ([At1], Theorem 1.6).

Theorem 4. [Fock space predictable representation property] – For all $f \in \Phi$, all $i \in \Lambda$ and for almost all $t \in \mathbb{R}^+$, the mapping $D_t^i f$ belongs to $\Phi = L^2(\mathcal{P})$. Furthermore, the family $(D^i f)_i$ is always Ito integrable and we have the representation

$$f = f(\emptyset)\Omega + \sum_{i \in \Lambda} \int_0^\infty D_t^i f d\chi_t^i \quad (4)$$

with the isometry formula

$$\|f\|^2 = |f(\emptyset)|^2 + \sum_{i \in \Lambda} \int_0^\infty \|D_t^i f\|^2 dt. \quad (5)$$

■

As an immediate corollary we get the following.

Corollary 5. – The representation (4) of f is unique. In particular, if $g \in \Phi$ is of the form

$$g = c\Omega + \sum_{i \in \Lambda} \int_0^\infty h_t^i d\chi_t^i$$

then for almost all t , all i in Λ ,

$$D_t^i g = h_t^i.$$

■

Let $f \in L^2(\mathcal{P}_n)$, one can easily define the *iterated Ito integral* on Φ :

$$I_n(f) = \int_{\mathcal{P}_n} f(\sigma) d\chi_\sigma$$

by iterating the definition of the Ito integral:

$$I_n(f) = \sum_{i_1, \dots, i_n \in \Lambda} \int_0^\infty \int_0^{t_1} \dots \int_0^{t_{n-1}} f^{i_1, \dots, i_n}(t_1, \dots, t_n) \Omega d\chi_{t_1}^{i_1} \dots d\chi_{t_n}^{i_n}.$$

We obtain this way an element of Φ which is actually the representant of f in the n -particle subspace of Φ , that is

$$[I_n(f)](\sigma) = \begin{cases} f^{i_1, \dots, i_n}(t_1, \dots, t_n) & \text{if } \sigma = \{t_1\}_{i_1} \cup \dots \cup \{t_n\}_{i_n} \\ 0 & \text{otherwise.} \end{cases}$$

For any $f \in \mathcal{P}$ we put

$$\int_{\mathcal{P}} f(\sigma) d\chi_\sigma$$

to denote the series of iterated Ito integrals

$$f(\emptyset)\Omega + \sum_{n=1}^\infty \sum_{i_1, \dots, i_n \in \Lambda} \int_0^\infty \int_0^{t_1} \dots \int_0^{t_{n-1}} f^{i_1, \dots, i_n}(t_1, \dots, t_n) \Omega d\chi_{t_1}^{i_1} \dots d\chi_{t_n}^{i_n}.$$

We then have the following representation ([At1], Theorem 1.7).

Theorem 6. [Fock space chaotic representation property]—Any element f of Φ admits an abstract chaotic representation

$$f = \int_{\mathcal{P}} f(\sigma) d\chi_{\sigma} \quad (6)$$

satisfying the isometry formula

$$\|f\|^2 = \int_{\mathcal{P}} |f(\sigma)|^2 d\sigma \quad (7).$$

This representation is unique. ■

The above theorem is the exact expression of the heuristics we wanted in order to describe the space

$$\Phi = \bigotimes_{\mathbb{R}^+} \mathcal{H}.$$

Indeed, we have, for each $t \in \mathbb{R}^+$, a family of elementary orthonormal elements Ω , $d\chi_t^i$, $i \in \Lambda$ (a basis of \mathcal{H}) whose (tensor) products $d\chi_{\sigma}$ form a continuous basis of Φ (formula (6)) and, even more, form an orthonormal continuous basis (formula (7)).

The attentive reader will have noticed that the only property of the curves χ^i that we really used is the fact that $\chi_t^i - \chi_s^i$ belongs to $\Phi_{[s,t]}$ for all $s \leq t$. One can naturally wonder if there exists another such family, which will then allow another Ito integral and furnish another continuous basis for Φ via another chaotic expansion property.

Of course there are obvious curves that can be obtained from the χ^i : for any function f on \mathbb{R}^+ and any $g \in L^2(\mathbb{R}^+ \times \Lambda)$ put

$$y_t = f(t)\Omega + \sum_i \int_0^t g^i(s)\Omega d\chi_s^i$$

for all $t \in \mathbb{R}^+$. Then one easily checks that (y_{\cdot}) satisfies the same property, namely, $y_t - y_s$ belongs to $\Phi_{[s,t]}$ for all $s \leq t$. But clearly the Ito integration theory obtained from y is the same as the one from χ , except that scalar factors $g^i(s)$ will appear in the integration.

One can wonder if there exist more complicated examples, giving rise to a different Ito integration. The following result shows that there are no more examples. In particular, there is only one Ito integral, one chaotic expansion and one natural continuous basis ([At1], Theorem 1.8).

Theorem 7.—Let $(y_t)_{t \geq 0}$ be a curve in Φ such that $y_t - y_s$ belongs to $\Phi_{[s,t]}$ for all $s \leq t$. Then there exist a function f on \mathbb{R}^+ and $g \in L^2(\mathbb{R}^+ \times \Lambda)$ such that

$$y_t = f(t)\Omega + \sum_i \int_0^t g^i(s)\Omega d\chi_s^i$$

for all $t \in \mathbb{R}^+$.



III.2 Quantum noises

The space Φ we have constructed is the natural space for defining quantum noises. These quantum noises are the natural, continuous-time, extensions of the basis operators $a_j^i(n)$ we met in the atom chain $\mathbb{T}\Phi$.

As indicated in the heuristic discussion above, we shall deal with a family of infinitesimal operators $da_j^i(t)$ on Φ which act on the continuous basis $d\chi_\sigma$ in the same way as their discrete-time counterparts $a_j^i(n)$ act on the X_A . The integrated version of the above heuristic infinitesimal formulas easily gives an exact formula for the action of the operators $a_j^i(t)$ on Φ :

$$\begin{aligned} [a_i^0(t)f](\sigma) &= \sum_{\substack{s \in \sigma_i \\ s \leq t}} f(\sigma \setminus \{s\}_i), \\ [a_0^i(t)f](\sigma) &= \int_0^t f(\sigma \cup \{s\}_i) ds, \\ [a_j^i(t)f](\sigma) &= \sum_{\substack{s \in \sigma_j \\ s \leq t}} f(\sigma \setminus \{s\}_j \cup \{s\}_i) \\ [a_0^0(t)f](\sigma) &= t f(\sigma) \end{aligned}$$

for $i, j \neq 0$.

All these operators, except $a_0^0(t)$, are unbounded. Yet note that a good common domain to all these operators is

$$\mathcal{D} = \left\{ f \in \Phi ; \int_{\mathcal{P}} |\sigma| |f(\sigma)|^2 d\sigma < \infty \right\}.$$

This family of operators is characteristic and universal in a sense which is close to the one of the curves χ_t^i . Indeed, one can easily check that in the decomposition of $\Phi \simeq \Phi_s \otimes \Phi_{[s,t]} \otimes \Phi_{[t]}$, the operators $a_j^i(t) - a_j^i(s)$ are all of the form

$$I \otimes (a_j^i(t) - a_j^i(s))_{|\Phi_{[s,t]}} \otimes I.$$

This property is fundamental for the definition of the quantum stochastic integrals and, in the same way as for (χ^i) , these operator families are the only ones to share that property (cf [Coq]).

This property allows to consider Riemann sums:

$$\sum_k H_{t_k} (a_j^i(t_{k+1}) - a_j^i(t_k)) \tag{8}$$

where $\mathcal{S} = \{0 = t_0 < t_1 < \dots < t_k < \dots\}$ is a partition of \mathbb{R}^+ , where $(H_t)_{t \geq 0}$ is a family of operators on Φ such that

– each H_t is an operator of the form $H_t \otimes I$ in the tensor product space $\Phi = \Phi_{t_j} \otimes \Phi_{[t]}$ (we say that H_t is a *t-adapted operator* and that $(H_t)_{t \geq 0}$ is an *adapted process of operators*),

– $(H_t)_{t \geq 0}$ is a *step process*, that is, it is constant on intervals:

$$H_t = \sum_k H_{t_k} \mathbb{1}_{[t_k, t_{k+1}]}(t),$$

and where the operator product $H_{t_k} (a_j^i(t_{k+1}) - a_j^i(t_k))$ is actually a tensor product of operators

$$H_{t_k} \otimes (a_j^i(t_{k+1}) - a_j^i(t_k)).$$

Note that, in particular, the above “product” is commutative and does not impose any new domain constraint.

The resulting operator associated to the Riemann sum (8) is denoted by

$$\int_0^\infty H_s da_j^i(s).$$

If we denote by T the above operator and by T_t the operator

$$\int_0^t H_s da_j^i(s) = \int_0^\infty H_s \mathbb{1}_{[0, t]}(s) da_j^i(s)$$

we can then compute the action of T on a “good” vector f of its domain and we obtain (cf [A-M] for more details)

$$Tf = \sum_{k \in \Lambda} \int_0^\infty T_t D_t^k f d\chi_t^k + \int_0^\infty H_t D_t^i f d\chi_s^j \quad (9)$$

with the notations: $D_t^0 = P_t$ and $d\chi_t^0 = dt$. For general operator processes (still adapted but not step process anymore) and general f , it is equation (9) which is kept as a definition for the domain and for the action of the operator

$$T = \int_0^\infty H_s da_j^i(s).$$

The maximal domain and the explicit action of the above operator can be described but is not worth developing here. The interested reader may refer to [At3], chapter 12 or to [A-L]. There are particular domains where the definition simplifies. The one we shall use here is the case of coherent vectors.

Indeed, if ϕ is any element of $L^2(\mathbb{R}^+, \mathcal{H}')$, consider the associated coherent vector $\varepsilon(\phi)$ in Φ . That is,

$$[\varepsilon(\phi)](\sigma) = \prod_i \prod_{s \in \sigma_i} \phi_i(s).$$

Put $\phi_0(s) = 1$ for all s . If ϕ is such that

$$\int_0^t |\phi_j(s)|^{(1-\delta_{0i})(2-\delta_{0j})} \|H_s \varepsilon(\phi)\|^{2-\delta_{0j}} ds < \infty$$

then $\int_0^t H_s da_j^i(s)$ is well-defined on $\varepsilon(\phi)$ with

$$\langle \varepsilon(\psi), \int_0^t H_s da_j^i(s) \varepsilon(\phi) \rangle = \int_0^t \overline{\psi_j}(s) \phi_i(s) \langle \varepsilon(\psi), H_s \varepsilon(\phi) \rangle ds.$$

for all $\psi \in L^2(\mathbb{R}^+, \mathcal{H}')$.

III.3 Embedding and approximation by the atom chain

We now describe the way the atom chain and its basic operators can be realized as a subspace of the Fock space and a projection of the quantum noises. The subspace associated to the atom chain is attached to the choice of some partition of \mathbb{R}^+ in such a way that the expected properties are satisfied:

- the associated subspaces increase when the partition refines and they constitute an approximation of Φ when the diameter of the partition goes to 0,
- the associated basic operators are restrictions of the others when the partition increases and they constitute an approximation of the quantum noises when the diameter of the partition goes to 0.

Note that this approximation has deep interpretations in terms of approximations of n -dimensional classical noises by extremal random walks in \mathbb{R}^n whose jumps take $n + 1$ different values. This aspect is developed in [A-P].

Let $\mathcal{S} = \{0 = t_0 < t_1 < \dots < t_n < \dots\}$ be a partition of \mathbb{R}^+ and $\delta(\mathcal{S}) = \sup_i |t_{i+1} - t_i|$ be the diameter of \mathcal{S} . For \mathcal{S} being fixed, define $\Phi_n = \Phi_{[t_{n-1}, t_n]}$, $n \in \mathbb{N}^*$. We clearly have that Φ is naturally isomorphic to the countable tensor product $\otimes_{n \in \mathbb{N}^*} \Phi_n$ (which is understood to be defined with respect to the stabilizing sequence $(\Omega)_{n \in \mathbb{N}}$).

For all $n \in \mathbb{N}^*$, define for $i, j \in \Lambda$

$$\begin{aligned} X^i(n) &= \frac{\chi_{t_n}^i - \chi_{t_{n-1}}^i}{\sqrt{t_n - t_{n-1}}} \in \Phi_n, \\ a_0^i(n) &= \frac{a_0^i(t_n) - a_0^i(t_{n-1})}{\sqrt{t_n - t_{n-1}}} \circ P_{1]}, \\ a_j^i(n) &= P_{1]} \circ (a_j^i(t_n) - a_j^i(t_{n-1})) \circ P_{1]}, \\ a_i^0(n) &= P_{1]} \circ \frac{a_i^0(t_n) - a_i^0(t_{n-1})}{\sqrt{t_n - t_{n-1}}}, \\ a_0^0(n) &= P_{0]}, \end{aligned}$$

where for $i = 0, 1$, $P_{i]}$ is the orthogonal projection onto $L^2(\mathcal{P}_i)$ and where the above definitions are understood to be valid on Φ_n only, the corresponding operator acting as the identity operator I on the others Φ_m 's.

For every $A \in \mathcal{P} = \mathcal{P}_{\mathbb{N}^*, \Lambda}$, define X_A from the $X^i(n)$'s in the same way as for $T\Phi$:

$$X_A = \Omega \otimes \dots \otimes \Omega \otimes X^{i_1}(n_1) \otimes \Omega \otimes \dots \otimes \Omega \otimes X^{i_2}(n_2) \otimes \dots$$

in $\otimes_{n \in \mathbb{N}^*} \mathcal{H}_n$.

Define $T\Phi(\mathcal{S})$ to be the space of $f \in \Phi$ which are of the form

$$f = \sum_{A \in \mathcal{P}} f(A) X_A$$

(note that the condition $\|f\|^2 = \sum_{A \in \mathcal{P}} |f(A)|^2 < \infty$ is automatically satisfied).

The space $T\Phi(\mathcal{S})$ is thus clearly identifiable to the spin chain $\mathbb{T}\Phi$. The space $T\Phi(\mathcal{S})$ is a closed subspace of Φ . We denote by $P_{\mathcal{S}}$ the operator of orthogonal projection from Φ onto $T\Phi(\mathcal{S})$. One can prove for example that the projection of an exponential vector is an ‘‘exponential vector’’ of the embedded toy Fock space: indeed, a direct computation shows that for any ϕ in $L^2(\mathbb{R}^+, \mathcal{H}')$,

$$(P_{\mathcal{S}} \varepsilon(\phi))(A) = \prod_i \prod_{n \in A_i} \tilde{\phi}_i(n)$$

where the function $\tilde{\phi}$ belongs to $l^2(\mathbb{N}^*, \mathcal{H}')$ and is defined by

$$\tilde{\phi}_i(n) = \frac{1}{\sqrt{t_n - t_{n-1}}} \int_{t_{n-1}}^{t_n} \phi_i(s) ds.$$

We will denote by $e(\tilde{\phi})$ such a discrete time version of a coherent vector.

We shall now check that the above operators $a_j^i(n)$ act on $\mathbb{T}\Phi(\mathcal{S})$ in the same way as the basic operators of $\mathbb{T}\Phi$.

Proposition 8. – *We have, for all $i, j \in \Lambda$*

$$\begin{cases} a_0^i(n) X^j(n) = \delta_{ij} \Omega \\ a_0^i \Omega = 0 \end{cases}$$

$$\begin{cases} a_j^i(n) X^k(n) = \delta_{ik} X^j(n) \\ a_j^i \Omega = 0 \end{cases}$$

$$\begin{cases} a_i^0(n) X^j(n) = 0 \\ a_i^0(n) \Omega = X^i(n) \end{cases}$$

$$\begin{cases} a_0^0(n) X^k(n) = 0 \\ a_j^i \Omega = \Omega. \end{cases}$$

Proof

This is a direct application of the definitions and computations using equation (9), cf [At2] for details. For example:

$$\begin{aligned} a_0^i(n) X^j(n) &= \frac{1}{t_n - t_{n-1}} (a_0^i(t_n) - a_0^i(t_{n-1})) \int_{t_{n-1}}^{t_n} \Omega d\chi_t^j \\ &= \frac{1}{t_n - t_{n-1}} \left[\sum_{k \in \Lambda} \int_{t_{n-1}}^{t_n} (a_0^i(t) - a_0^i(t_{n-1})) \Omega d\chi_t^k + \int_{t_{n-1}}^{t_n} \delta_{ij} \Omega dt \right] \\ &= \frac{1}{t_n - t_{n-1}} (0 + (t_n - t_{n-1}) \delta_{ij} \Omega) = \delta_{ij} \Omega. \end{aligned}$$

And so on for the other cases. ■

Thus the action of the operators a_j^i on the $X^i(n)$ is exactly the same as the action of the corresponding operators on the spin chain of section II; the operators

$a_j^i(n)$ act on $T\Phi(\mathcal{S})$ exactly in the same way as the corresponding operators do on $\mathbb{T}\Phi$. We have completely embedded the toy Fock space in the Fock space.

The action of operators $a_j^i(n)$ on discrete exponential vectors as defined above will be most useful in the sequel. The following lemma is deduced immediately from Proposition 8.

Lemma 9. – *For any ϕ, ψ in $L^2(\mathbb{R}_+, \mathcal{H})$ and for any t_n -adapted operator H_n the bracket*

$$\left\langle e(\tilde{\phi}), H_n a_j^i(n+1) e(\tilde{\psi}) \right\rangle$$

is equal to

$$\overline{\tilde{\phi}}_j(n+1) \tilde{\psi}_i(n+1) \left\langle e(\tilde{\phi}_{[n]}), H_n e(\tilde{\psi}_{[n]}) \right\rangle \left\langle e(\tilde{\phi}_{[n+2]}), e(\tilde{\psi}_{[n+2]}) \right\rangle.$$

Where $\phi_{[n]}, \psi_{[n]}, \phi_{[n+2]}, \psi_{[n+2]}$ are the restrictions of ϕ, ψ to $\{1, \dots, n\}, \{n+2, \dots\}$ respectively.

This lemma is the basis for our future computations involving discrete-time quantum stochastic integrals (for more precise treatment of this subject see [Pa1] or [Pa3]).

We are now going to see that the Fock space Φ and its basic operators $a_j^i(t)$, $i, j \in \Lambda \cup \{0\}$ can be approached by the toy Fock spaces $T\Phi(\mathcal{S})$ and their basic operators $a_j^i(n)$.

We are given a sequence $(\mathcal{S}_n)_{n \in \mathbb{N}}$ of refining partitions whose diameter $\delta(\mathcal{S}_n)$ tends to 0 when n tends to $+\infty$. Let $T\Phi(n) = T\Phi(\mathcal{S}_n)$ and $P_n = P_{\mathcal{S}_n}$, for all $n \in \mathbb{N}$.

Theorem 10. –

i) *The orthogonal projectors P_n strongly converge to the identity operator I on Φ . That is, any $f \in \Phi$ can be approached in Φ by a sequence $(f_n)_{n \in \mathbb{N}}$ such that $f_n \in T\Phi(n)$ for all $n \in \mathbb{N}$.*

ii) *If $\mathcal{S}_n = \{0 = t_0^n < t_1^n < \dots < t_k^n < \dots\}$, then for all $t \in \mathbb{R}^+$, all $i, j \in \Lambda$, the operators*

$$\sum_{k; t_k^n \leq t} a_j^i(k), \quad \sum_{k; t_k^n \leq t} \sqrt{t_k^n - t_{k-1}^n} a_0^i(k), \quad \sum_{k; t_k^n \leq t} \sqrt{t_k^n - t_{k-1}^n} a_i^0(k),$$

and $\sum_{k; t_k^n \leq t} (t_k^n - t_{k-1}^n) a_0^0(k)$

converge strongly on \mathcal{D} to $a_j^i(t)$, $a_0^i(t)$, $a_i^0(t)$ and $a_0^0(t)$ respectively.

Proof

i) As the \mathcal{S}_n are refining then the $(P_n)_n$ forms an increasing family of orthogonal projection in Φ . Let $P_\infty = \vee_n P_n$. Clearly, for all $s \leq t$, we have that $\chi_t^i - \chi_s^i$ belongs to $\text{Ran} P_\infty$. But by the construction of the Ito integral and by Theorem 6, we have that the $\chi_t^i - \chi_s^i$ generate Φ . Thus $P_\infty = I$.

ii) Let us check the case of a_i^0 . A direct computation shows that, for $f \in \mathcal{D}$

$$\left[\sum_{k; t_k^n \leq t} \sqrt{t_k^n - t_{k-1}^n} a_i^0(k) f \right] (\sigma) = \sum_{k; t_k^n \leq t} \mathbb{1}_{|\sigma_i \cap [t_{k-1}^n, t_k^n]|=1} \sum_{s \in \sigma_i \cap [t_{k-1}^n, t_k^n]} f(\sigma \setminus \{s\}_i).$$

Put $t^n = \sup\{t_k^n \in \mathcal{S}_n ; t_k^n \leq t\}$. We have

$$\begin{aligned} & \left\| \left(\sum_{k; t_k^n \leq t} \sqrt{t_k^n - t_{k-1}^n} a_i^0(k) - a_i^0(t) \right) f \right\|^2 \\ &= \int_{\mathcal{P}} \left| \sum_{k; t_k^n \leq t} \mathbb{1}_{|\sigma_i \cap [t_{k-1}^n, t_k^n]|=1} \sum_{s \in \sigma_i \cap [t_{k-1}^n, t_k^n]} f(\sigma \setminus \{s\}_i) - \sum_{s \in \sigma_i \cap [0, t]} f(\sigma \setminus \{s\}_i) \right|^2 d\sigma \\ &\leq 2 \int_{\mathcal{P}} \left| \sum_{s \in \sigma \cap [t^n, t]} f(\sigma \setminus \{s\}_i) \right|^2 d\sigma + 2 \int_{\mathcal{P}} \left| \sum_{k; t_k^n \leq t} \mathbb{1}_{|\sigma_i \cap [t_{k-1}^n, t_k^n]| \geq 2} \right. \\ &\quad \left. \times \sum_{s \in \sigma_i \cap [t_{k-1}^n, t_k^n]} f(\sigma \setminus \{s\}_i) \right|^2 d\sigma. \end{aligned}$$

For any fixed σ , the terms inside each of the integrals above converge to 0 when n tends to $+\infty$. Furthermore we have, for n large enough,

$$\begin{aligned} \int_{\mathcal{P}} \left| \sum_{s \in \sigma \cap [t^n, t]} f(\sigma \setminus \{s\}_i) \right|^2 d\sigma &\leq \int_{\mathcal{P}} |\sigma| \sum_{\substack{s \in \sigma \\ s \leq t+1}} |f(\sigma \setminus \{s\}_i)|^2 d\sigma \\ &= \int_0^{t+1} \int_{\mathcal{P}} (|\sigma| + 1) |f(\sigma)|^2 d\sigma ds \\ &\leq (t+1) \int_{\mathcal{P}} (|\sigma| + 1) |f(\sigma)|^2 d\sigma \end{aligned}$$

which is finite for $f \in \mathcal{D}$;

$$\begin{aligned} & \int_{\mathcal{P}} \left| \sum_{k; t_k^n \leq t} \mathbb{1}_{|\sigma_i \cap [t_{k-1}^n, t_k^n]| \geq 2} \sum_{s \in \sigma_i \cap [t_{k-1}^n, t_k^n]} f(\sigma \setminus \{s\}_i) \right|^2 d\sigma \\ &\leq \int_{\mathcal{P}} \left(\sum_{k; t_k^n \leq t} \mathbb{1}_{|\sigma_i \cap [t_{k-1}^n, t_k^n]| \geq 2} \left| \sum_{s \in \sigma_i \cap [t_{k-1}^n, t_k^n]} f(\sigma \setminus \{s\}_i) \right| \right)^2 d\sigma \\ &\leq \int_{\mathcal{P}} \left(\sum_{k; t_k^n \leq t} \sum_{s \in \sigma_i \cap [t_{k-1}^n, t_k^n]} |f(\sigma \setminus \{s\}_i)| \right)^2 d\sigma \\ &= \int_{\mathcal{P}} \left(\sum_{\substack{s \in \sigma_i \\ s \leq t^n}} |f(\sigma \setminus \{s\}_i)| \right)^2 d\sigma \\ &= \int_{\mathcal{P}} |\sigma| \sum_{\substack{s \in \sigma_i \\ s \leq t^n}} |f(\sigma \setminus \{s\}_i)|^2 d\sigma \\ &\leq (t+1) \int_{\mathcal{P}} (|\sigma| + 1) |f(\sigma)|^2 d\sigma < \infty \end{aligned}$$

in the same way as above. So we can apply Lebesgue's theorem. This proves the result.

The other cases are treated in the same way. See [At2] for details. ■

We have fulfilled our duties: not only does the space $\mathbb{T}\Phi(\mathcal{S})$ recreate $\mathbb{T}\Phi$ and its basic operators as a subspace of Φ and a projection of its quantum noises, but, when $\delta(\mathcal{S})$ tends to 0, this realization constitutes an approximation of the space Φ and of its quantum noises.

To any operator H on Φ we can associate the projected operator $P_{\mathcal{S}}HP_{\mathcal{S}}$ which acts on the atom chain only and which approximates H (if H is bounded for example).

We wish to compute the corresponding projections of the quantum stochastic integral operators. We reduce our computations to the case where integrals are of the type $H = \int_0^\infty H_t^\epsilon da_j^i(s)$, with $(i, j) \neq (0, 0)$, and satisfy the following conditions **(HS)**:

- the operator H is bounded and
- the integrands $H_j^i(t)$ are bounded for all t and $t \mapsto \|H_j^i(t)\|$ is square integrable if one of i or j is zero, essentially bounded otherwise.

Even though they are rather restrictive, these hypotheses will suffice for our needs.

The following result is a consequence of the theory and the computations developed in [Pa1] and [Pa2] (adapted here to the case of higher multiplicity) and we do not reproduce the proof here. It is also stated in the following form in [Pa3], chapter 4.

Theorem 11. – *Let $(i, j) \neq (0, 0)$ be fixed. Let $H = \int H_j^i(t) da_j^i(t)$ be a quantum stochastic integral on Φ that satisfies the assumptions **(HS)**. Then $P_{\mathcal{S}}HP_{\mathcal{S}}$ is an operator on $\mathbb{T}\Phi$ of the form*

$$\sum_{k,l} \sum_n h_l^k(n) a_l^k(n+1)$$

where the sum is over all couples (k, l) in $(\Lambda \cup \{0\})^2$ different from $(0, 0)$ and is meaningful in the weak sense. The operators h_l^k are given by :

- if both i and j are nonzero,

$$h_l^k(n) = \delta_{ki} \delta_{lj} \frac{1}{t_{n+1} - t_n} P_{\mathcal{S}} \int_{t_n}^{t_{n+1}} P_{t_n} H_j^i(t) dt$$

- if $i = 0$,

$$h_l^0(n) = \delta_{lj} \frac{1}{\sqrt{t_{n+1} - t_n}} P_{\mathcal{S}} \int_{t_n}^{t_{n+1}} P_{t_n} H_j^0(t) dt$$

and for all $k \neq 0$,

$$h_l^k(n) = \delta_{lj} \frac{1}{t_{n+1} - t_n} P_{\mathcal{S}} \int_{t_n}^{t_{n+1}} P_{t_n} H_j^0(t) (a_k^0(t) - a_k^0(t_n)) dt$$

– if $j = 0$,

$$h_0^k(n) = \delta_{ki} \frac{1}{\sqrt{t_{n+1} - t_n}} P_S \int_{t_n}^{t_{n+1}} P_{t_n} H_0^i(t) dt$$

and for all $l \neq 0$,

$$h_l^k(n) = \delta_{ki} \frac{1}{t_{n+1} - t_n} P_S \int_{t_n}^{t_{n+1}} P_{t_n} (a_0^l(t) - a_0^l(t_n)) H_0^i(t) dt.$$

■

III.4 Quantum Langevin equations

In this article what we call quantum Langevin equation is actually a restricted version of what is usually understood in the literature (cf [G-Z]); by this we mean that we here study the so-called quantum stochastic differential equations as defined by Hudson and Parthasarathy and heavily studied by further authors.

This type of quantum noise perturbation of the Schrödinger equation is exactly the type of equation which we will get as the continuous limit of our Hamiltonian description of repeated quantum interactions.

The aim of quantum stochastic differential equations is to study equations of the form

$$dU_t = \sum_{i,j \in \Lambda \cup \{0\}} L_j^i U_t da_j^i(t), \quad (10)$$

with initial condition $U_0 = I$. The above equation has to be understood as an integral equation

$$U_t = I + \int_0^t \sum_{i,j \in \Lambda \cup \{0\}} L_j^i U_s da_j^i(s),$$

for operators on $\mathcal{H}_0 \otimes \Phi$, the operators L_j^i being bounded operators on \mathcal{H}_0 alone which are amplified to $\mathcal{H}_0 \otimes \Phi$.

The main motivation and application of that kind of equation is that it gives an account of the interaction of the small system \mathcal{H}_0 with the bath Φ in terms of quantum noise perturbation of a Schrödinger-like equation. Indeed, the first term of the equation

$$dU_t = L_0^0 U_t dt + \dots$$

describes the induced dynamics on the small system, all the other terms are quantum noises terms.

One of the main application of equations such as (10) is that they give explicit constructions of unitary dilations of semigroups of completely positive maps of $\mathcal{B}(\mathcal{H}_0)$ (see [H-P]). Let us here only recall one of the main existence, uniqueness and boundedness theorems connected to equations of the form (10). The literature is huge about those equations; we refer to [Par] for the result we mention here.

Remark : The Hilbert-Schmidt type of summability condition we use here (in the case where \mathcal{H} is infinite dimensionnal) is not quite the usual one used in

the theory of quantum stochastic differential equations, but it is the one we need for our convergence theorems.

Theorem 12. – Let \mathcal{H}_0 be a separable Hilbert space and let L_j^i , $i, j \in \Lambda \cup \{0\}$, be bounded operators on \mathcal{H}_0 . Assume that

$$\|L\| = \left(\sum_{i,j \in \Lambda \cup \{0\}} \|L_j^i\|^2 \right)^{1/2} < +\infty,$$

then the quantum stochastic differential equation

$$U_t = I + \sum_{i,j} \int_0^t L_j^i U_t da_j^i(t)$$

admits a unique solution defined on the space of coherent vectors.

Furthermore, if the coefficients L_j^i are of the form

$$L_0^0 = -(iH + \frac{1}{2} \sum_{k \in \Lambda} L_k^* L_k)$$

$$L_j^0 = L_j$$

$$L_0^i = - \sum_{k \in \Lambda} L_k^* S_i^k$$

$$L_j^i = S_j^i - \delta_{ij} I.$$

for a self-adjoint operator H , some operators L_i , $i \in \Lambda$ and some operators S_j^i , $i, j \in \Lambda$ such that $(S_j^i)_{i,j \in \Lambda}$ is unitary, then the solution $(U_t)_{t \geq 0}$ is made of unitary operators. ■

IV Convergence theorems

IV.1 Convergence to quantum Langevin equations

We are now ready to put all the pieces of the jigsaw puzzle together and prove that the Hamiltonian dynamic associated to repeated quantum interactions spontaneously converges to a quantum Langevin equation under some normalization conditions on the Hamiltonian. Notice that for the first convergence theorem we no longer assume that $\mathbb{L}(h)$ has been conveniently constructed for our needs; in particular \mathbb{L} is not assumed to be unitary. We shall come back to the unitary case later on.

Let h be a parameter in \mathbb{R}^+ , which is thought of as representing a small time interval. Let $\mathbb{L}(h)$ be an operator on $\mathcal{H}_0 \otimes \mathcal{H}$, with coefficients $\mathbb{L}_j^i(h)$ as a matrix of operators on \mathcal{H}_0 . Let $u_n(h)$ be the associated solution of

$$u_{n+1}(h) = \mathbb{L}_{n+1}(h)u_n(h)$$

with the same notation as in section II.3. In the following we will drop dependency in h and write simply \mathbb{L} , or u_n . Besides, we denote

$$\varepsilon_{ij} = \frac{1}{2}(\delta_{0i} + \delta_{0j})$$

for all i, j in $\Lambda \cup \{0\}$. That is,

$$\varepsilon_{i0} = e_{0j} = \frac{1}{2}, \quad \varepsilon_{ij} = 0, \quad \varepsilon_{00} = 1.$$

Note that from now on we take the embedding of $\mathbb{T}\Phi$ in Φ for granted and we consider, without mentioning it, all the repeated quantum interactions to happen in $\mathbb{T}\Phi(\hbar)$, the subspace of Φ associated to the partition $\mathcal{S} = \{t_i = ih; i \in \mathbb{N}\}$. We also make the convention that the default summation sets for sums is $\Lambda \cup \{0\}$, e.g. \sum_i is $\sum_{i \in \Lambda \cup \{0\}}$.

The following theorem is our general convergence result.

Theorem 13. – Assume that there exist bounded operators L_j^i , $i, j \in \Lambda \cup \{0\}$ on \mathcal{H}_0 such that

$$\sum_{i, j \in \Lambda \cup \{0\}} \|L_j^i\|^2 < +\infty$$

and

$$\lim_{\hbar \rightarrow 0} \sum_{i, j \in \Lambda \cup \{0\}} \left\| \frac{\mathbb{L}_j^i(\hbar) - \delta_{ij}I}{\hbar^{\varepsilon_{ij}}} - L_j^i \right\|^2 = 0$$

for all $j \in \Lambda \cup \{0\}$. Assume that the quantum stochastic differential equation

$$dU_t = \sum_{i, j} L_j^i U_t da_j^i(t)$$

with initial condition $U_0 = I$ admits a unique solution $(U_t)_{t \geq 0}$ which is a process of bounded operators with locally uniform norm bound.

Then, for all t , for every ϕ, ψ in $L^\infty([0, t])$, the quantity the quantity

$$\langle a \otimes \varepsilon(\phi), P_{\mathcal{S}} u_{[t/\hbar]} P_{\mathcal{S}} b \otimes \varepsilon(\psi) \rangle$$

converges to

$$\langle a \otimes \varepsilon(\phi), U_t b \otimes \varepsilon(\psi) \rangle$$

when \hbar goes to 0.

Moreover, the convergence is uniform for a, b in any bounded ball of \mathcal{H} , uniform for t in a bounded interval of \mathbb{R}_+ .

Remarks

– This is where we particularize the index zero : the above hypotheses of convergence simply mean that, among the coefficients of \mathbb{L} ,

$$(\mathbb{L}_0^0(\hbar) - I)/\hbar \text{ converges,}$$

$$\mathbb{L}_j^i(\hbar)/\sqrt{\hbar} \text{ converges if either } i \text{ or } j \text{ is zero,}$$

$$\mathbb{L}_j^i(\hbar) - \delta_{i,j} \text{ converges if neither } i \text{ nor } j \text{ is zero.}$$

We here meet the announced three time scales appearing in the Hamiltonian. We shall discuss the physical meaning of these normalizations in next section.

– In the case where the operator \mathbb{L} is unitary and satisfies the convergence assumptions of the above theorem, then one can see that the limiting operators L_j^i are of the form given in the second part of Theorem 11.

– In that case, the solution $(U_t)_{t \in \mathbb{R}_+}$ enjoys a particular algebraic property which we won't define here: it is a *cocycle* (see [H-P] or chapter 6 of [Pa3]). This property traduces the fact that the evolution of the system is, in the limit, memory-less.

Consider the quantum stochastic differential equation on $\mathcal{H}_0 \otimes \Phi$:

$$dU_s = \sum_{i,j} L_j^i U_s da_j^i(s)$$

where the L_j^i are the bounded operators on the initial space \mathcal{H}_0 given by our assumptions.

We consider that h is fixed and the associated partition $\mathcal{S} = \{0 = t_0 < t_1 = h < \dots < t_k = kh < \dots\}$ is also fixed. Note that we have chosen a regular partition only for simplicity and that all our results hold with general partitions when the mesh size tends to 0. We fix some bounded interval $[0, T]$ of \mathbb{R}_+ .

We will proceed by successive simplifications. Consider the operator on the atom chain defined by

$$w_k = P_{\mathcal{S}} U_{t_k} P_{\mathcal{S}}.$$

The following lemma will be used over and again: as a first application note that, together with obvious estimates it allows us to prove the convergence in Theorem 13 for almost all t only.

Lemma 14. – For any $r < s$, for any vectors $a \otimes \varepsilon(\phi)$, $b \otimes \varepsilon(\psi)$ such that

$$\phi, \psi \in L^2(\mathbb{R}^+; \mathcal{H}') \cap L^\infty(\mathbb{R}^+; \mathcal{H}'),$$

we have

$$| \langle a \otimes \varepsilon(\phi), (U_s - U_r) b \otimes \varepsilon(\psi) \rangle | \leq C \|a\| \|b\| (s - r)$$

where C depends only on $\|L\|$, defined in Theorem 12, and on the L^2 and L^∞ norms of ϕ and ψ .

Proof

$$\begin{aligned} & | \langle a \otimes \varepsilon(\phi), (U_s - U_r) b \otimes \varepsilon(\psi) \rangle | \\ & \leq \sum_{i,j} \int_r^s |\bar{\phi}_i(u)| |\psi_j(u)| | \langle a \otimes \varepsilon(\phi), L_j^i U_u b \otimes \varepsilon(\psi) \rangle | du \\ & \leq \|L\| \int_s^t \|\phi(u)\| \|\psi(u)\| \|a \otimes \varepsilon(\phi)\| \|U_u b \otimes \varepsilon(\psi)\| du \end{aligned}$$

from which the estimate follows, using the fact that $\|\phi(u)\| \leq \|\phi\|_\infty$ and $\|\psi(u)\| \leq \|\psi\|_\infty$, the L_j^i are bounded and U is locally uniformly bounded. \blacksquare

The following lemma shows that $(w_k)_k$ converges to $(U_t)_{t \geq 0}$, in the same weak sense as in the theorem, as h goes to 0.

Lemma 15. – For any $t_k < s$, any vectors $a \otimes \varepsilon(\phi)$, $b \otimes \varepsilon(\psi)$ with

$$\phi, \psi \in L^2(\mathbb{R}^+; \mathcal{H}') \cap L^\infty(\mathbb{R}^+; \mathcal{H}'),$$

we have

$$|\langle a \otimes \varepsilon(\phi), (w_k - U_s)b \otimes \varepsilon(\psi) \rangle| \leq C \|a\| \|b\| ((s - t_k) + \|(I - P_S)\varepsilon(\phi)\| + \|(I - P_S)\varepsilon(\psi)\|).$$

where C depends only on $\|L\|$ and on the L^2 and L^∞ norms of ϕ and ψ .

Proof

We have

$$\begin{aligned} |\langle a \otimes \varepsilon(\phi), (w_k - U_s)b \otimes \varepsilon(\psi) \rangle| &\leq \\ &\leq |\langle a \otimes \varepsilon(\phi), (U_{t_k} - U_s)b \otimes \varepsilon(\psi) \rangle| \\ &\quad + |\langle a \otimes \varepsilon(\phi), (P_S U_{t_k} P_S - U_{t_k} P_S)b \otimes \varepsilon(\psi) \rangle| \\ &\quad + |\langle a \otimes \varepsilon(\phi), (U_{t_k} P_S - U_{t_k})b \otimes \varepsilon(\psi) \rangle| \\ &\leq \sum_{i,j} \int_{t_k}^s |\bar{\phi}_i(u)| |\psi_j(u)| |\langle a \otimes \varepsilon(\phi), L_j^i U_u b \otimes \varepsilon(\psi) \rangle| du \\ &\quad + \|(I - P_S)a \otimes \varepsilon(\phi)\| \|U_{t_k} P_S b \otimes \varepsilon(\psi)\| \\ &\quad + \|U_{t_k}^* a \otimes \varepsilon(\phi)\| \|(I - P_S)b \otimes \varepsilon(\psi)\| \end{aligned}$$

and we conclude as in the previous lemma. ■

We can now prove Theorem 13.

Proof of Theorem 13

Let $\omega_j^i(h)$ be such that

$$\mathbb{L}_j^i(h) - \delta_{ij}I = h^{\varepsilon_{ij}}(L_j^i + \omega_j^i(h))$$

for all i, j in $\Lambda \cup \{0\}$. In particular we have that,

$$\sum_{i,j \in \Lambda \cup \{0\}} \|\omega_j^i(h)\|^2$$

converges to 0 when h tends to 0.

Consider the solution $(u_n)_{n \in \mathbb{N}^*}$ of

$$u_{n+1} = \mathbb{L}_{n+1} u_n$$

with the notations of section II.3. Note that if A denotes the matrix $\mathbb{L} - (\delta_{ij}I)_{i,j}$ we then have

$$u_{n+1} - u_n = A_{n+1} u_n.$$

Let F be the matrix $(h^{\varepsilon_{ij}} L_j^i + \widehat{\delta}_{ij} h L_0^0)_{i,j}$ where

$$\widehat{\delta}_{ij} = \begin{cases} 1 & \text{if } i = j \text{ and } (i, j) \neq (0, 0), \\ 0 & \text{if } i \neq j \text{ or } (i, j) = (0, 0) \end{cases}$$

and consider the solution $(v_n)_{n \in \mathbb{N}}$ of the equation

$$v_{n+1} - v_n = F_{n+1} v_n.$$

Note that

$$A_{n+1} = \sum_{i,j} A_j^i a_j^i(n+1)$$

and similarly for F_{n+1} .

Also note that $a_j^i(n+1)$ commutes with u_n (*resp.* v_n), for they do not act on the same part of the space $\mathbb{T}\Phi$. Thus we get another useful way to write the above equations in terms of the basis $a_j^i(n)$:

$$u_{n+1} - u_n = \sum_{i,j} A_j^i u_n a_j^i(n+1).$$

and

$$v_{n+1} - v_n = \sum_{i,j} \left(h^{\varepsilon_{ij}} L_j^i + \widehat{\delta}_{ij} h L_0^0 \right) v_n a_j^i(n+1).$$

From the above lemma it is enough to prove the convergence to zero of $u_n - w_n$. We actually start with $w_n - v_n$.

From the fact that

$$U_{t_{k+1}} - U_{t_k} = \sum_{i,j} \int_{t_k}^{t_{k+1}} L_j^i U_s da_j^i(s)$$

and thanks to the formulas for projections of Fock space integrals onto the toy Fock space in Theorem 11, one obtains the following expression for $w_{k+1} - w_k$ (be careful that the $da_0^0(t)$ integrals gives rise to $a_i^i(k)$ terms for *all* i , for $I = \sum_i a_i^i$):

$$\begin{aligned} w_{k+1} - w_k &= \sum_{i,j \neq (0,0)} h^{\varepsilon_{ij}} L_j^i \left(\frac{1}{h} P_S \int_{t_k}^{t_{k+1}} P_{t_k} U_t dt \right) a_j^i(k+1) P_S \\ &+ \sum_i h L_0^0 \left(\frac{1}{h} P_S \int_{t_k}^{t_{k+1}} U_t dt \right) a_i^i(k+1) P_S \\ &+ \sum_{i \in \Lambda} \sum_{j \in \Lambda} P_S \left(\frac{1}{h} \int_{t_k}^{t_{k+1}} \left(L_0^i P_{t_k} U_t (a_0^j(t) - a_0^j(t_k)) \right) dt \right. \\ &\left. + \frac{1}{h} \int_{t_k}^{t_{k+1}} \left(L_j^0 P_{t_k} U_t (a_i^0(t) - a_i^0(t_k)) \right) dt \right) a_j^i(k+1) P_S. \end{aligned}$$

As a consequence

$$\begin{aligned} w_n - v_n &= \sum_{k < n} \sum_{i,j \neq (0,0)} h^{\varepsilon_{ij}} L_j^i a_j^i(k+1) \left(\frac{1}{h} P_S \int_{t_k}^{t_{k+1}} P_{t_k} U_t dt P_S - v_k \right) \\ &+ \sum_{k < n} \sum_i h L_0^0 a_i^i(k+1) \left(\frac{1}{h} P_S \int_{t_k}^{t_{k+1}} U_t dt P_S - v_k \right) \\ &+ \sum_{k < n} \sum_{i \in \Lambda} \sum_{j \in \Lambda} P_S \left(\frac{1}{h} \int_{t_k}^{t_{k+1}} \left(L_0^i P_{t_k} U_t (a_0^j(t) - a_0^j(t_k)) \right) dt \right. \\ &\left. + \frac{1}{h} \int_{t_k}^{t_{k+1}} \left(L_j^0 P_{t_k} U_t (a_i^0(t) - a_i^0(t_k)) \right) dt \right) P_S a_j^i(k+1). \quad (11) \end{aligned}$$

We first wish to replace $\frac{1}{h} P_S \int_{t_k}^{t_{k+1}} P_{t_k} U_t dt$ or $\frac{1}{h} P_S \int_{t_k}^{t_{k+1}} U_t dt$ in the first two terms by w_k . Lemma 14 allows us to estimate the error term. Consider two essentially

bounded functions $\phi, \psi \in L(\mathbb{R}^+)$ and two vectors a, b in the unit ball \mathcal{H}_1 of \mathcal{H} . We expand the first terms in (11) and estimate each term :

$$\begin{aligned} & \left| \langle a \otimes e(\tilde{\phi}), \sum_{k < n} h^{\varepsilon_{ij}} L_j^i \left(\frac{1}{h} P_S \int_{t_k}^{t_{k+1}} P_{t_k} U_t dt - w_k \right) a_j^i(k+1) b \otimes e(\tilde{\psi}) \rangle \right| \\ & \leq \sum_{k < n} \left| \tilde{\phi}_j(k) \right| \left| \tilde{\psi}_i(k) \right| \left| \langle a \otimes e(\tilde{\phi}_{[k]}), P_S h^{\varepsilon_{ij}} L_j^i \frac{1}{h} \int_{t_k}^{t_{k+1}} (U_s - U_{t_k}) ds b \otimes e(\tilde{\psi}_{[k]}) \rangle \right|, \end{aligned}$$

where we have omitted a uniformly bounded factor

$$\left\langle a \otimes e(\tilde{\phi}_{[k+2]}), b \otimes e(\tilde{\psi}_{[k+2]}) \right\rangle.$$

From the fact that $\sum_i \left| \tilde{\phi}_i(k) \right| \leq \sqrt{h} \|\phi\|_\infty$, $\sum_i \left| \tilde{\psi}_i(k) \right| \leq \sqrt{h} \|\psi\|_\infty$ we obtain an estimation of the error term of the form

$$C \|a\| \|b\| \sqrt{h}$$

for some constant C which depends only on $\|L\|$ and on the L^2 and L^∞ norms of ϕ and ψ .

On the other hand, the error term obtained by replacing $\frac{1}{h} P_S \int_{t_k}^{t_{k+1}} U_t dt$ by w_k in (11) is clearly dominated by Ch in norm because $\sum_i h L_0^i a_i^i(k+1)$ is just $h L_0^0$.

We now seek to evaluate the third sum in (11); for that consider again two functions ϕ, ψ in \mathcal{M} and two vectors a, b in the unit ball \mathcal{H}_1 of \mathcal{H} ; we have, up to an uniformly bounded factor,

$$\begin{aligned} & \sum_{i,j \in \Lambda} \left| \langle a \otimes \varepsilon(\phi), \frac{1}{h} L_0^i P_S \left(\int_{t_k}^{t_{k+1}} P_{t_k} U_s (a_0^j(s) - a_0^j(t_k)) ds \right) a_j^i(k+1) P_S b \otimes \varepsilon(\psi) \rangle \right| \\ & = \sum_{i,j \in \Lambda} \left| \tilde{\phi}_j(k) \tilde{\psi}_i(k) \right| \left| \langle a \otimes e(\tilde{\phi}_{[k]}), \frac{1}{h} L_0^i \int_{t_k}^{t_{k+1}} U_s (a_0^j(s) - a_0^j(t_k)) b \otimes e(\tilde{\psi}_{[k]}) ds \rangle \right| \\ & \leq Ch^{3/2}, \end{aligned}$$

for $a_0^j(s) - a_0^j(t_k)$ is bounded on Φ_{t_k} , with norm $\sqrt{s - t_k}$. One obtains similarly

$$\begin{aligned} & \sum_{i,j \in \Lambda} \left| \langle a \otimes \varepsilon(\phi), \frac{1}{h} L_j^0 P_S \int_{t_k}^{t_{k+1}} P_{t_k} U_s (a_i^0(s) - a_i^0(t_k)) ds a_j^i(k+1) P_S b \otimes \varepsilon(\psi) \rangle \right| \\ & \leq Ch^{3/2}, \end{aligned}$$

so that the third sum in (11) is bounded by $C \sqrt{h} 2t \|\phi\|_\infty \|\psi\|_\infty$.

$$\begin{aligned} & \text{We have shown that, putting } F_k = \sum_{i,j} (h^{\varepsilon_{ij}} L_j^i + \hat{\delta}_{ij} h L_0^i) a_j^i(k+1), \\ & \left| \langle a \otimes \varepsilon(\phi), P_S (w_n - v_n) P_S b \otimes \varepsilon(\psi) \rangle \right| \\ & = \sum_{k < n} \left| \langle a \otimes e(\tilde{\phi}), F_k (w_k - v_k) b \otimes e(\tilde{\psi}) \rangle \right| + o(1) \quad (12) \end{aligned}$$

where $o(1)$ is a term which converges to zero as h goes to zero *uniformly for a, b in \mathcal{H}_1 and for t in a bounded interval*. That uniform convergence property will be important in the sequel.

Expanding F_k in equation (12) gives

$$\begin{aligned} & \left| \langle a \otimes \varepsilon(\phi), P_S(w_n - v_n)P_S b \otimes \varepsilon(\psi) \rangle \right| \\ & \leq \sum_{i,j} \sum_{k < n} h^{\varepsilon_{ij}} \left| \tilde{\phi}_j(k) \right| \left| \tilde{\psi}_i(k) \right| \left| \langle (L_j^i)^* a \otimes e(\tilde{\phi}_k), (w_k - v_k)b \otimes e(\tilde{\psi}_k) \rangle \right| \\ & \quad + \sum_{i \in \Lambda} \sum_{k < n} h \left| \tilde{\phi}_i(k) \right| \left| \tilde{\psi}_i(k) \right| \left| \langle (L_0^0)^* a \otimes e(\tilde{\phi}_k), (w_k - v_k)b \otimes e(\tilde{\psi}_k) \rangle \right| \\ & \quad + o(1), \end{aligned}$$

where in each term we have omitted an uniformly bounded factor and the notation $o(1)$ indicates a function of h which converges to zero as h goes to zero.

Since ϕ and ψ are essentially bounded, the quantities $\sum_{i,j} h^{\varepsilon_{ij}} |\tilde{\phi}_j(k)| |\tilde{\psi}_i(k)|$ and $\sum_{i \in \Lambda} h |\tilde{\phi}_i(k)| |\tilde{\psi}_i(k)|$ are again of order h at most, with an estimate which is independent on k . Besides, remark that normalizing all operators would only imply an additional constant factor, so that we can assume all the L_j^i to be contractions. In that case the above implies that for all n ,

$$\begin{aligned} & \sup_{a,b \in \mathcal{H}_1} \left| \langle a \otimes e(\tilde{\phi}_n), (w_n - v_n)b \otimes e(\tilde{\psi}_n) \rangle \right| \\ & \leq hC \sum_{k < n} \sup_{a,b \in \mathcal{H}_1} \left| \langle a \otimes e(\tilde{\phi}_k), (w_k - v_k)b \otimes e(\tilde{\psi}_k) \rangle \right| + o(1) \end{aligned}$$

for some constant C . Here we have used our earlier remark that all convergences are uniform in $a, b \in \mathcal{H}_1$. This implies that

$$\sup_{a,b \in \mathcal{H}_1} \left| \langle a \otimes e(\tilde{\phi}_n), (w_n - v_n)b \otimes e(\tilde{\psi}_n) \rangle \right| \leq (1 + Ch)^n \times o(1)$$

and since nh converges to t , the quantity $(1 + Ch)^n$ is bounded so that

$$\sup_{a,b \in \mathcal{H}_1} \left| \langle a \otimes e(\tilde{\phi}_n), (w_n - v_n)b \otimes e(\tilde{\psi}_n) \rangle \right|$$

converges to zero as h goes to zero.

We have proved the desired convergence property for the process $(w_k)_{k \geq 0}$. Now we will prove that

$$\sup_{a,b \in \mathcal{H}_1} \left| \langle a \otimes e(\tilde{\phi}_n), (u_n - v_n)b \otimes e(\tilde{\psi}_n) \rangle \right|$$

converges to zero as h goes to zero. We have

$$\begin{aligned} u_n - v_n &= \sum_{k < n} F_k(u_k - v_k) + \sum_{k < n} \left(\sum_{i,j} h^{\varepsilon_{ij}} (\omega_j^i(h) - h\widehat{\delta}_{ij}L_0^0) a_j^i(k+1) \right) u_k \\ &= \sum_{k < n} \left(F_k + \sum_{i,j} h^{\varepsilon_{ij}} (\omega_j^i(h) - h\widehat{\delta}_{ij}L_0^0) a_j^i(k+1) \right) (u_k - v_k) \\ & \quad + \sum_{k < n} \left(\sum_{i,j} h^{\varepsilon_{ij}} (\omega_j^i(h) - h\widehat{\delta}_{ij}L_0^0) a_j^i(k+1) \right) v_k \end{aligned} \tag{13}$$

Replace for simplicity $\omega_j^i(h)$ by $\omega_j^i(h) - h\widehat{\delta}_{ij}L_0^0$ (which does not change the validity of the earlier estimates). The interest of the second form lies therein, that the term without recurring $u_k - v_k$ can now be estimated thanks to our previous result:

$$\left| \langle a \otimes e(\tilde{\phi}_k), \sum_{k < n} \left(\sum_{i,j} h^{\varepsilon_{ij}} \omega_j^i(h) a_j^i(k+1) \right) v_k b \otimes e(\tilde{\psi}_k) \rangle \right|$$

is bounded by

$$\sum_{k < n} \left(\sum_{i,j} h^{\varepsilon_{ij}} \left| \tilde{\phi}_j(k) \right| \left| \tilde{\psi}_i(k) \right| \left\| \omega_j^i(h) \right\| \right) \sup_{a,b \in \mathcal{H}_1} \left| \langle a \otimes e(\tilde{\phi}_k], v_k b \otimes e(\tilde{\psi}_k] \rangle \right| \quad (14)$$

with

$$\begin{aligned} \sup_{a,b \in \mathcal{H}_1} \left| \langle a \otimes e(\tilde{\phi}_k], v_k b \otimes e(\tilde{\psi}_k] \rangle \right| \\ \leq \sup_{a,b \in \mathcal{H}_1} \left(\left| \langle a \otimes e(\tilde{\phi}_k], (v_k - w_k) b \otimes e(\tilde{\psi}_k] \rangle \right| \right. \\ \left. + \left| \langle a \otimes e(\tilde{\phi}_k], w_k b \otimes e(\tilde{\psi}_k] \rangle \right| \right) \end{aligned}$$

and our estimate shows that the first term on the right-hand-side converges to zero uniformly in k as h goes to zero. The second term is, in turn, bounded by $\|\varepsilon(\phi)\| \|\varepsilon(\psi)\|$ since any w_k is $P_{\mathcal{S}} U_{t_k} P_{\mathcal{S}}$ and as such has bounded norm.

Thanks to our assumptions on the perturbative operators, the bound (14) converges to zero as h goes to zero, uniformly for a, b in \mathcal{H}_1 .

Besides, for the recurring term in (13) we easily obtain as before

$$\begin{aligned} \sup_{a,b \in \mathcal{H}_1} \left| \langle a \otimes e(\tilde{\phi}_n], \sum_{k < n} \left(F_k + \sum_{i,j} h^{\varepsilon_{ij}} \omega_j^i(h) a_j^i(k+1) \right) (w_k - v_k) b \otimes e(\tilde{\psi}_n] \rangle \right| \\ \leq h C \sum_{k < n} \sup_{a,b \in \mathcal{H}_1} \left| \langle a \otimes e(\tilde{\phi}_k], (v_k - w_k) b \otimes e(\tilde{\psi}_k] \rangle \right|, \end{aligned}$$

thanks to the fact that the ω_j^i are assumed to have norms which converge to zero uniformly in h . We conclude as in the previous case.

This ends the proof. ■

Under some additional assumptions, which are verified in many applications, we can very much improve the convergence.

Theorem 16. – *Consider the same assumptions and the same notations as in Theorem 13. If $\|u_k\|_{\mathbb{T}\Phi}$ (the norm of operators on $\mathbb{T}\Phi$) is locally uniformly bounded, then $P_{\mathcal{S}} u_{[t/h]} P_{\mathcal{S}}$ converges weakly to U_t on all $\mathcal{H}_0 \otimes \Phi$. If furthermore $\|u_k\|_{\Phi}$ (the norm of operators on Φ) is locally uniformly bounded, then $u_{[t/h]}$ converges weakly to U_t on all $\mathcal{H}_0 \otimes \Phi$.*

Proof

Theorem 13 allows us to perform a $\epsilon/3$ argument with an approximation of any vectors of $\mathcal{H}_0 \otimes \Phi$ by combinations of vectors $a \otimes \varepsilon(\phi)$, $b \otimes \varepsilon(\psi)$ with essentially bounded functions ϕ, ψ . ■

One of the main application of this last theorem is the case when the matrices $\mathbb{L}(h)$ give rise at the limit to a matrix L such as in Theorem 12 (the case of a unitary solution $(U_t)_{t \geq 0}$). We shall show that the associated discrete evolution $(u_n)_{n \in \mathbb{N}}$ satisfy the conditions of the above theorem, so that the convergence of $u_{[t/h]}$ towards U_t is weak.

Theorem 17. – Consider a matrix \mathbb{L} on $\mathcal{H}_0 \otimes \mathcal{H}$ with coefficients

$$\mathbb{L}_0^0 = I - h(iH + \frac{1}{2} \sum_k L_k^* L_k) + h\omega_0^0$$

$$\mathbb{L}_j^0 = \sqrt{h}L_j + h\omega_j^0$$

$$\mathbb{L}_0^i = -\sqrt{h} \sum_k L_k^* S_j^k + h\omega_0^i$$

$$\mathbb{L}_j^i = I + S_j^i - \delta_{ij}I + h\omega_j^i$$

where H is a bounded self-adjoint operator, $(S_j^i)_{i,j \in \Lambda}$, is unitary and L_i , $i \in \Lambda$ are operators on \mathcal{H}_0 , where the coefficients ω_j^i are such that $\sum_{i,j} \|\omega_j^i\|^2$ is uniformly bounded and $\|\omega_0^0(h)\|$ converge to 0 when h tends to 0.

Then the solution $(u_n)_{n \in \mathbb{N}}$ of

$$u_{n+1} = \mathbb{L}_{n+1} u_n$$

is made of invertible operators which are locally uniformly bounded in norm.

In particular $u_{[t/h]}$ converges weakly to the solution U_t of (X).

Proof

A straightforward computation shows the special form of \mathbb{L} induces many cancellations when computing the coefficients of $\mathbb{L}^* \mathbb{L} - I$ and of $\mathbb{L} \mathbb{L}^* - I$, and that they are of order h . Thus for h small enough the operators $\mathbb{L}^* \mathbb{L}$, $\mathbb{L} \mathbb{L}^*$ and thus \mathbb{L} are invertible. Thus so are the operators u_n .

Furthermore the above estimates show that $\|\mathbb{L}\| \leq \sqrt{1 + Ch}$. This easily gives the locally uniform boundedness of $(u_n)_{n \in \mathbb{N}}$ and thus the desired weak convergence. ■

Specializing some more will allow us to answer the natural question of convergence of Heisenberg evolutions of observables:

Corollary 18. – If the operator \mathbb{L} is unitary and satisfies the conditions of Theorem 13 then the solution $(U_t)_{t \in \mathbb{R}_+}$ of (E) is unitary. In this case the convergence of $u_{[t/h]}$ to U_t is strong and for all bounded operator X on $\mathcal{H}_0 \otimes \Phi$, all t , the sequence $u_{[t/h]}^* X u_{[t/h]}$ converges weakly to $U_t^* X U_t$ as h tends to zero.

Proof

It is easy to see from the above conditions that the operators $(L_j^i)_{i,j}$ satisfy for all i, j

$$L_j^i + L_j^{i*} + \sum_{k \in \Lambda} L_k^i L_k^{j*} = 0$$

$$L_j^i + L_j^{i*} + \sum_{k \in \Lambda} L_i^k L_j^{k*} = 0$$

which implies that the equation (E) is of the form which has unitary solutions (see Theorem 12).

By Theorem 17 and Proposition 1, for all t , $u_{[t/h]}$ is a sequence of contractions that converges weakly to a unitary operator, so that strong convergence also holds.

It is now straightforward to prove the statement regarding convergence of $u_{[t/h]}^* X u_{[t/h]}$ to $U_t^* X U_t$ for bounded X . \blacksquare

IV.2 Typical Hamiltonian

We are now coming back to the initial physical motivations of Theorems 13, 16 and 17. These theorems show up quite strong conditions on the unitary operator $\mathbb{L} = e^{-ihH}$ and a natural question now is: what kind of Hamiltonian H will produce such conditions on \mathbb{L} ? What is the typical Hamiltonian for repeated quantum interactions which will produce quantum Langevin equations at the continuous limit?

In this section we answer partly that question. We answer it as we exhibit a large family of such Hamiltonians and we conjecture that they are the typical ones. We do not fully answer the question for we are not able to prove that they are the only ones.

We keep here the notations of section II.1-II.3.

On $\mathcal{H}_0 \otimes \mathcal{H}$ consider the following Hamiltonian

$$H = H_0 \otimes I + I \otimes H_S + \frac{1}{\sqrt{h}} \sum_{i \in \Lambda} (V_i \otimes a_i^0 + V_i^* \otimes a_i^0) + \frac{1}{h} \sum_{i,j \in \Lambda} D_{ij} \otimes a_j^i \quad (15)$$

where H_0 , V_i and D_{ij} are bounded operators on \mathcal{H}_0 (with H_0 hermitian and $D_{ij} = D_{ji}^*$), and H_S is hermitian on \mathcal{H} and contains the basis elements X^i in its domain.

Put \mathcal{H}' to be the closed subspace of \mathcal{H} generated by the basis elements X^i , $i \in \Lambda$; that is, the orthogonal of $X^0 = \Omega$. Consider the ‘‘column operator’’

$$V = \begin{pmatrix} V_1 \\ V_2 \\ \vdots \end{pmatrix}$$

as an operator from \mathcal{H}_0 to $\mathcal{H}_0 \otimes \mathcal{H}'$. Assume that this operator is bounded.

The adjoint of V is then the ‘‘row operator’’

$$V^* = (V_1 \quad V_2 \quad \dots)$$

from $\mathcal{H}_0 \otimes \mathcal{H}'$ to \mathcal{H}_0 .

Define the ‘‘matrix operator’’ $D = (D_{ij}^i)_{ij}$ as an operator from $\mathcal{H}_0 \otimes \mathcal{H}'$ to $\mathcal{H}_0 \otimes \mathcal{H}'$. We also assume D to be bounded.

Let us relate the above Hamiltonian with the usual literature on weak coupling and low density limits.

Recall that, in the literature about weak coupling limit, the bath is usually made of several harmonic oscillators (a Fock space) with associated creation operators $a^*(g)$ and annihilation operators $a(g)$, where g runs over a Hilbert space \mathcal{H} . The Hamiltonian which is considered is then of the form

$$H = H_0 \otimes I + I \otimes H_S + \lambda (V \otimes a^*(g) + V^* \otimes a(g))$$

The part $V \otimes a^*(g) + V^* \otimes a(g)$ corresponds to the typical dipole Hamiltonian usually considered in the weak coupling limit or van Hove limit (cf [Dav], [D-J]).

This Hamiltonian meets (15) when considering an orthonormal basis (e_i) of \mathcal{H} and when one writes

$$(V \otimes a^*(g) + V^* \otimes a(g)) = \sum_i (\langle e_i, g \rangle V \otimes a^*(e_i) + \langle g, e_i \rangle V^* \otimes a(e_i))$$

Our time renormalization term $1/\sqrt{\hbar}$ does not actually corresponds to the usual time renormalisation for the weak coupling limit ($t/\lambda^2 = \tau$), we are more in a situation which looks like a strong coupling limit. Actually, as is shown in [AJ1], they are intimately related, when considering the setup of weak coupling limit for repeated interactions.

On the other hand, interaction Hamiltonians of the form $D \otimes a^*(f)a(h) + D^* \otimes a^*(h)a(f)$ are of scattering type. They present some important similarity with the typical interaction Hamiltonians considered in the low density limit (cf [APV], [AFL]), but we are not able to show that this similarity contains anything deeper.

We are now back to our general Hamiltonian (15).

We shall take compact notations for the unitary quantum Langevin equations of Theorem 10. Consider the equation

$$dU_t = \sum_{i,j} L_j^i U_t da_j^i(t) \tag{16}$$

with

$$L_0^0 = -(iK + \frac{1}{2} \sum_{k \in \Lambda} L_k^* L_k)$$

$$L_j^0 = L_j$$

$$L_0^i = - \sum_{k \in \Lambda} L_k^* S_i^k$$

$$L_j^i = S_j^i - \delta_{ij} I,$$

where K is a bounded self-adjoint operator and $(S_j^i)_{i,j \in \Lambda}$ is unitary.

We write W for the column operator

$$\begin{pmatrix} L_1 \\ L_2 \\ \vdots \end{pmatrix}$$

and S for the matrix operator $(S_j^i)_{i,j}$. Then, with obvious notations

$$\begin{aligned} L_0^0 &= -(iK + \frac{1}{2}W^*W) \\ L_0^1 &= W \\ L_0^2 &= -W^*S \\ L_1 &= S - I. \end{aligned}$$

With the Hamiltonian H given by (15), put

$$U = e^{-ihH}$$

and consider the evolution equation for repeated interactions associated to U :

$$u_{n+1} = U_{n+1}u_n.$$

Theorem 19. – *The solution $(u_n)_{n \in \mathbb{N}^*}$ of the discrete time evolution equation converges strongly in $\mathcal{H}_0 \otimes \Phi$ to the solution $(U_t)_{t \geq 0}$ of the quantum Langevin equation*

$$dU_t = \sum_{i,j \in \Lambda \cup \{0\}} L_j^i U_t da_j^i(t)$$

where, with the same notations as above

$$\begin{aligned} K &= H_0 + \langle \Omega, H_S \Omega \rangle I + V^* D^{-2} (\sin D - D) V \\ W &= D^{-1} (e^{-iD} - I) V \\ S &= e^{-iD}. \end{aligned}$$

Proof

Put $k_j^i = \langle X^j, H_S X^i \rangle$ for all $i, j \in \Lambda \cup \{0\}$. We consider the column operator

$$|k\rangle = \begin{pmatrix} k_1^0 I \\ k_2^0 I \\ \vdots \end{pmatrix},$$

the row operator

$$\langle k| = (k_0^1 I \quad k_0^2 I \quad \dots)$$

and the matrix operator

$$k = (k_j^i I)_{i,j \in \Lambda}.$$

They all are bounded operators.

Put $\tilde{H} = H_0 + k_0^0 I$ and $M = (M_j^i)_{i,j \in \Lambda}$ with $M_j^i = \delta_{ij} H_0 + k_j^i I$. We then have

$$H = \begin{pmatrix} \tilde{H} & \frac{1}{\sqrt{\hbar}} V^* + \langle k| \\ \frac{1}{\sqrt{\hbar}} V + |k\rangle & \frac{1}{\hbar} D + M \end{pmatrix}$$

as an operator on $\mathcal{H}_0 \otimes \mathcal{H}$ which is decomposed as an operator on $\mathcal{H}_0 \otimes (\mathbb{C}\Omega \oplus \mathcal{H}')$. In particular

$$hH = \begin{pmatrix} h\tilde{H} & \sqrt{h}V^* + h\langle k | \\ \sqrt{h}V + h|k\rangle & D + hM \end{pmatrix}.$$

Let α be a bound for $\|\tilde{H}\|, \|V\|, \|D\|, \|M\|, \|\langle k | \|, \| |k\rangle \|$.

Lemma 20. – For all $m \in \mathbb{N}$ we have

$$(hH)^m = \begin{pmatrix} hA_m + h^{3/2}R_m^1 & \sqrt{h}B_m + hR_m^2 \\ \sqrt{h}C_m + hR_m^3 & D_m + hR_m^4 \end{pmatrix}$$

with

$$\|X_m\| \leq \alpha^m$$

for all $X = A, B, C, D$ and

$$\|R_m^i\| \leq 7^{m-1}\alpha^m$$

for all $i = 1, 2, 3, 4$; with

$$\begin{aligned} A_{m+1} &= V^*C_m, & A_0 &= I, A_1 = \tilde{H} \\ B_{m+1} &= V^*D_m, & B_0 &= 0, B_1 = V^* \\ C_{m+1} &= DC_m, & C_0 &= 0, C_1 = V \\ D_{m+1} &= DD_m, & D_0 &= I. \end{aligned}$$

Proof of the lemma

For $m = 0$ the statements are clearly satisfied. For $m = 1$ we find the announced A_1, B_1, C_1, D_1 and $R_1^1 = 0, R_1^2 = \langle k |, R_1^3 = |k\rangle, R_1^4 = M$. The norm estimates are then clearly satisfied. Now, applying the induction hypothesis and computing $(hH)^{m+1}$ from $(hH)^m$ we get

$$\begin{aligned} A_{m+1} &= V^*C_m & \text{and } \|A_{m+1}\| &\leq \alpha\alpha^m = \alpha^{m+1} \\ R_{m+1}^1 &= \sqrt{h}\tilde{H}A_m + h\tilde{H}R_m^1 + V^*R_m^3 + \langle k |C_m + \sqrt{h}\langle k |R_m^3 \\ \|R_{m+1}^1\| &\leq 5(7^{m-1}\alpha^{m+1}) \leq 7^m\alpha^{m+1} \\ B_{m+1} &= V^*D_m & \text{and } \|B_{m+1}\| &\leq \alpha\alpha^m = \alpha^{m+1} \\ R_{m+1}^2 &= \sqrt{h}\tilde{H}B_m + h\tilde{H}R_m^2 + \sqrt{h}V^*R_m^4 + \langle k |D_m + h\langle k |R_m^4 \\ \|R_{m+1}^2\| &\leq 5(7^{m-1}\alpha^{m+1}) \leq 7^m\alpha^{m+1} \\ C_{m+1} &= DC_m & \text{and } \|C_{m+1}\| &\leq \alpha\alpha^m = \alpha^{m+1} \\ R_{m+1}^3 &= \sqrt{h}VA_m + hVR_m^1 + DR_m^3 + \sqrt{h}MC_m + hMR_m^3 \\ \|R_{m+1}^3\| &\leq 5(7^{m-1}\alpha^{m+1}) \leq 7^m\alpha^{m+1} \\ D_{m+1} &= DD_m & \text{and } \|D_{m+1}\| &\leq \alpha\alpha^m = \alpha^{m+1} \\ R_{m+1}^4 &= VB_m + hVR_m^2 + \sqrt{h}|k\rangle B_m + h|k\rangle R_m^2 + DR_m^4 + MD_m + hMR_m^4 \\ \|R_{m+1}^4\| &\leq 7(7^{m-1}\alpha^{m+1}) \leq 7^m\alpha^{m+1}. \end{aligned}$$

The lemma is proved.

In particular, we get

$$\begin{aligned} D_m &= D^m \\ B_m &= V^* D^{m-1} \\ C_m &= D^{m-1} V \\ A_m &= V^* D^{m-2} V. \end{aligned}$$

By the norm estimates of Lemma 20 we have that the series

$$\sum_m \frac{(-i)^m}{m!} R_m^j$$

are norm convergent. Let us denote by R^k their respective limit.

We get

$$\begin{aligned} \mathbb{L} &= \sum_m \frac{(-i)^m}{m!} (hH)^m = \\ &= \begin{pmatrix} I - ih\tilde{H} + h \sum_{m=2}^{\infty} \frac{(-i)^m}{m!} V^* D^{m-2} V & \sqrt{h} \sum_{m=1}^{\infty} \frac{(-i)^m}{m!} V^* D^{m-1} \\ +h^{3/2} R^1 & +hR^2 \\ \sqrt{h} \sum_{m=1}^{\infty} \frac{(-i)^m}{m!} D^{m-1} V & I + \sum_{m=1}^{\infty} \frac{(-i)^m}{m!} D^m \\ +hR^3 & +hR_m^4 \end{pmatrix}. \end{aligned}$$

This gives

$$U = \begin{pmatrix} I - ih\tilde{H} + hV^* D^{-2} (e^{-iD} - I + iD)V & \sqrt{h} V^* D^{-1} (e^{-iD} - I) \\ +h^{3/2} R^1 & +hR^2 \\ \sqrt{h} D^{-1} (e^{-iD} - I)V & I + (e^{-iD} - I) \\ +hR^3 & +hR_m^4 \end{pmatrix}.$$

We are exactly in the conditions of Corollary 18 and we thus get the strong convergence to the solution of

$$dU_t = \sum_{i,j \in \Lambda \cup \{0\}} L_j^i U_t da_j^i(t)$$

where

$$\begin{aligned} L_0^0 &= -i\tilde{H} + V^* D^{-2} (e^{-iD} - I + iD)V \\ L_{\cdot}^0 &= D^{-1} (e^{-iD} - I)V \\ L_0^{\cdot} &= V^* D^{-1} (e^{-iD} - I) \\ L_{\cdot}^{\cdot} &= e^{-iD} - I. \end{aligned}$$

If we put $W = L^0$ and $S = e^{-iD}$, we then get

$$-W^* S = -V^* (e^{iD} - I) D^{-1} S = V^* D^{-1} (e^{-iD} - I) = L_0^{\cdot}$$

and

$$\begin{aligned} -\frac{1}{2} W^* W &= -\frac{1}{2} V^* (e^{iD} - I) D^{-1} D^{-1} (e^{-iD} - I) V \\ &= -\frac{1}{2} V^* D^{-2} (\cos D - I) V. \end{aligned}$$

This shows that

$$L_0^0 = -i\tilde{H} - iV^*D^{-2}(\sin D - D)V - \frac{1}{2}W^*W.$$

The theorem is proved. ■

Let us interpret the above theorem in terms of weak coupling and low density limit again. If in the above Hamiltonian we consider no term D_{ij} , that is,

$$H = H_0 \otimes I + I \otimes H_S + \frac{1}{\sqrt{\hbar}} \sum_{i \in \Lambda} (V_i \otimes a_i^0 + V_i^* \otimes a_i^0)$$

then we are in a similar situation than the one of a weak coupling limit, with its typical dipole interaction Hamiltonian. The quantum Langevin equation we obtain in Theorem 16 then simplifies to

$$\begin{aligned} dU_t = - \left(iH_0 + i\langle \Omega, H_S \Omega \rangle I + \frac{1}{2} \sum_i V_i^* V_i \right) U_t dt + \\ + \sum_i V_i U_t da_i^0(t) - \sum_i V_i^* U_t da_i^0(t). \end{aligned}$$

This is also typical of the “diffusion” type of quantum Langevin equation one can meet in the literature for this kind of limit. The fact that only creation and annihilation quantum noises are involved is here the quantum analogue of a classical stochastic differential equation with Brownian noise. Note that if the V_i 's are such that $V_i^* = -V_i$ then the above quantum Langevin equation becomes a classical stochastic differential equation with Brownian noises:

$$dU_t = - \left(iH_0 + i\langle \Omega, H_S \Omega \rangle I + \frac{1}{2} \sum_i V_i^2 \right) U_t dt + \sum_i V_i U_t dW_i(t).$$

We refer to [At1] for a complete discussion on the classical stochastic interpretations of the quantum noises.

On the other hand, if in the Hamiltonian we consider no term V_i , that is,

$$H = H_0 \otimes I + I \otimes H_S + \frac{1}{\hbar} \sum_{i,j \in \Lambda} D_{ij} \otimes a_j^i$$

then the quantum Langevin equation we obtain in Theorem 19 simplifies to

$$dU_t = -i(H_0 + \langle \Omega, H_S \Omega \rangle I)U_t dt + \sum_{i,j \in \Lambda} (S_j^i - \delta_{ij})U_t da_j^i(t)$$

where $S = e^{-iD}$.

This is a rather classical type of Langevin equation which actually describes a scattering of the bath.

The surprise in our setup in the apparition of the term

$$V^*D^{-2}(\sin D - D)V$$

only when both the limits are in presence in the Hamiltonian.

Indeed, the term

$$L_0^0 = -(iK + \frac{1}{2}W^*W)$$

is the driving term of the dynamic associated to the quantum Langevin equation. In some sense it is the generator of the dynamic on \mathcal{H}_0 . The part $\frac{1}{2}W^*W$ is representative of the dissipation from \mathcal{H}_0 to the bath. The part iK is an effective Hamiltonian on \mathcal{H}_0 . The apparition of this new contribution $V^*D^{-2}(\sin D - D)V$ is new in this situation.

Instead of interpreting the typical hamiltonian

$$H = H_0 \otimes I + I \otimes H_S + \frac{1}{\sqrt{\hbar}} \sum_{i \in \Lambda} (V_i \otimes a_i^0 + V_i^* \otimes a_0^i) + \frac{1}{\hbar} \sum_{i,j \in \Lambda} D_{ij} \otimes a_j^i$$

as being \hbar dependent, one can view the above expression as a usual constant Hamiltonian

$$H = H_0 \otimes I + I \otimes H_S + \sum_{i \in \Lambda} (V_i \otimes a_i^0 + V_i^* \otimes a_0^i) + \sum_{i,j \in \Lambda} D_{ij} \otimes a_j^i$$

but for a photon field which has be renormalized:

$$a_i^0 \mapsto \frac{1}{\sqrt{\hbar}} a_i^0.$$

Indeed this renormalization implies

$$a_0^i \mapsto \frac{1}{\sqrt{\hbar}} a_0^i$$

and

$$a_j^i = a_0^i a_j^0 \mapsto \frac{1}{\hbar} a_j^i.$$

IV.3 Hamiltonian description of von Neumann measurements

Our setup and approach allows to construct an Hamiltonian description for the usual von Neumann measurement procedure (collapse of the wave packet postulate).

On some quantum system state space \mathcal{H}_0 consider an observable A with discrete spectrum (maybe infinite). Let P_1, P_2, \dots denote the associated spectral (orthogonal) projections, with $\sum_k P_k = I$.

We want to give a model for the action on \mathcal{H}_0 of an exterior measurement apparatus which measures the observable A . That is, the action of the measurement apparatus on \mathcal{H}_0 should be to transform any observable X of \mathcal{H}_0 into

$$\sum_k P_k X P_k.$$

Let \mathcal{H} be a Hilbert space with one more dimension than the number of projectors P_k involved above (infinite dimensional if the P_k 's are in infinite number). On \mathcal{H} consider an orthonormal basis $\Omega = X_0, X_1, X_2, \dots$ and the associated creation

operators a_k^0 and annihilation operators a_k^k , as in sections II.1-II.3. Consider the following Hamiltonian on $\mathcal{H}_0 \otimes \mathcal{H}$:

$$H = \frac{1}{\sqrt{\hbar}} \sum_k (iP_k \otimes a_k^0 - iP_k \otimes a_k^k).$$

Let $U = e^{-ihH}$ and consider the process

$$u_{n+1} = \mathbb{L}_{n+1} u_n$$

of repeated quantum interactions on $\mathcal{H}_0 \otimes \bigotimes_{\mathbb{N}^*} \mathcal{H}$.

Theorem 21. – *The repeated quantum interaction process $(u_n)_{n \in \mathbb{N}^*}$ converges weakly in $\mathcal{H}_0 \otimes \bigotimes_{\mathbb{R}^+} \mathcal{H}$, when h tends to 0, to the solution $(U_t)_{t \geq 0}$ of the quantum Langevin equation*

$$dU_t = -\frac{1}{2}U_t dt + \sum_k (iP_k U_t da_k^0(t) - iP_k U_t da_k^k(t)).$$

Furthermore, for any observable X on \mathcal{H}_0 , the partial trace along the field $\bigotimes_{\mathbb{R}^+} \mathcal{H}$, in the vacuum state, of $U_t^*(X \otimes I)U_t$ converges, when t tends to $+\infty$, to

$$\sum_k P_k X P_k.$$

Proof

In the basis $\Omega = X_0, X_1, X_2, \dots$ for \mathcal{H} the Hamiltonian H writes as

$$H = \begin{pmatrix} 0 & -i\frac{1}{\sqrt{\hbar}}P_1 & -i\frac{1}{\sqrt{\hbar}}P_2 & \dots \\ i\frac{1}{\sqrt{\hbar}}P_1 & 0 & 0 & \dots \\ i\frac{1}{\sqrt{\hbar}}P_2 & 0 & 0 & \dots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

In particular, as an easy direct computation shows, we have

$$\mathbb{L} = \begin{pmatrix} \cos \sqrt{\hbar} I & -\sin \sqrt{\hbar} P_1 & -\sin \sqrt{\hbar} P_2 & \dots \\ \sin \sqrt{\hbar} P_1 & \cos \sqrt{\hbar} P_1 & 0 & \dots \\ \sin \sqrt{\hbar} P_2 & 0 & \cos \sqrt{\hbar} P_2 & \dots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

This unitary matrix clearly satisfies the conditions of Corollary 18 (one could also have directly applied Theorem 19 to the Hamiltonian), we thus have the weak convergence to the announced quantum Langevin equation

$$dU_t = -\frac{1}{2}U_t dt + \sum_k (iP_k U_t da_k^0(t) - iP_k U_t da_k^k(t)),$$

with $U_0 = I$.

Now, consider the evolution under $(U_t)_{t \geq 0}$ of a system observable X :

$$U_t^*(X \otimes I)U_t$$

and the partial trace along the field $\bigotimes_{\mathbb{R}^+} \mathcal{H}$ in the vacuum state:

$$P_t(X) = \langle \Omega, U_t^*(X \otimes I)U_t\Omega \rangle$$

with the notation

$$\langle a, \langle \Omega, A\Omega \rangle b \rangle_{\mathcal{H}_0} = \langle a \otimes \Omega, A(b \otimes \Omega) \rangle_{\mathcal{H}_0 \otimes \bigotimes_{\mathbb{R}^+} \mathcal{H}}$$

for any operator A on $\mathcal{H}_0 \otimes \bigotimes_{\mathbb{R}^+} \mathcal{H}$, any $a, b \in \mathcal{H}_0$.

It is well known from the usual theory of quantum stochastic differential equations (cf [H-P]), that $(P_t)_{t \geq 0}$ is then a semigroup, on $\mathcal{B}(\mathcal{H}_0)$, of completely positive maps with Lindblad generator

$$\mathcal{L}(X) = -\frac{1}{2} \sum_k (P_k X + X P_k - 2P_k X P_k).$$

One could also have obtained this ‘‘quantum master equation’’ by simply computing the discrete time quantum master equation associated to $(u_n)_{n \in \mathbb{N}^*}$, using Theorem 2 and then by passing to the limit $h \rightarrow 0$ to recover $(P_t)_{t \geq 0}$ (see section IV.5). Our approach has the advantage to also describe the exact equation for the interaction with the bath.

We thus get

$$\mathcal{L}(X) = \sum_k P_k X P_k - X$$

so that

$$\mathcal{L}^2(X) = -\mathcal{L}(X)$$

and

$$P_t(X) = e^{t\mathcal{L}}(X) = (I + (1 - e^{-t})\mathcal{L})(X).$$

That is,

$$P_t(X) = (1 - e^{-t}) \sum_k P_k X P_k + e^{-t} X$$

which converges to

$$\sum_k P_k X P_k$$

when t tends to $+\infty$. ■

We thus have proved that a von Neumann measurement apparatus can be described in an Hamiltonian setup by, first considering an Hamiltonian description of a repeated quantum interaction, second, passing to the limit to continuous quantum interactions ($h \rightarrow 0$) and third, passing to the limit to large times ($t \rightarrow +\infty$).

The authors are grateful to Prof. K.B. Sinha for bringing to their attention this application.

IV.4 An example

We shall here follow a very basic example. It is actually the simplest non-trivial physical example and it already gives very interesting consequences.

Assume $\mathcal{H}_0 = \mathcal{H} = \mathbb{C}^2$ that is, both are two-level systems with basis states Ω (the fundamental state) and X (the excited state).

During the small amount of time h the two systems are in contact and they evolve in the following way:

- if the states of the two systems are the same (both fundamental or both excited) then nothing happens;
- if they are different (one fundamental and the other one excited) then they can either be exchanged or stay as they are.

In the base $\{\Omega \otimes \Omega, \Omega \otimes X, X \otimes \Omega, X \otimes X\}$ the operator \mathbb{L} is taken to be of the form

$$\mathbb{L} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos \alpha & -\sin \alpha & 0 \\ 0 & \sin \alpha & \cos \alpha & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

The associated Hamiltonian is thus

$$H = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & -i\alpha/h & 0 \\ 0 & i\alpha/h & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

so that $\mathbb{L} = e^{-ihH}$.

For the choice $\alpha = \sqrt{h}$, that is,

$$H = \frac{1}{\sqrt{h}} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & -i & 0 \\ 0 & i & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

we get

$$\mathbb{L} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos \sqrt{h} & -\sin \sqrt{h} & 0 \\ 0 & \sin \sqrt{h} & \cos \sqrt{h} & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

Repeating this interaction leads to an Hamiltonian equation of the form

$$u_{n+1} = \mathbb{L}_{n+1} u_n$$

with coefficients

$$\begin{aligned} \mathbb{L}_0^0 &= \begin{pmatrix} 1 & 0 \\ 0 & \cos \sqrt{h} \end{pmatrix}, \quad \mathbb{L}_1^0 = \begin{pmatrix} 0 & \sin \sqrt{h} \\ 0 & 0 \end{pmatrix}, \\ \mathbb{L}_0^1 &= \begin{pmatrix} 0 & 0 \\ -\sin \sqrt{h} & 0 \end{pmatrix}, \quad \mathbb{L}_1^1 = \begin{pmatrix} \cos \sqrt{h} & 0 \\ 0 & 1 \end{pmatrix} \end{aligned}$$

for \mathbb{L} . We then have

$$\lim_{h \rightarrow 0} \frac{\mathbb{L}_0^0 - I}{h} = \begin{pmatrix} 0 & 0 \\ 0 & -1/2 \end{pmatrix}$$

$$\begin{aligned}\lim_{h \rightarrow 0} \frac{\mathbb{I}_1^0}{\sqrt{h}} &= \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \\ \lim_{h \rightarrow 0} \frac{\mathbb{I}_0^1}{\sqrt{h}} &= \begin{pmatrix} 0 & 0 \\ -1 & 0 \end{pmatrix} \\ \lim_{h \rightarrow 0} \mathbb{I}_1^1 - I &= \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}.\end{aligned}$$

We are exactly in the condition for applying Theorem 15 since for all h the matrix $\mathbb{I}(h)$ is unitary. We get that for all t , $u_{[t/h]}$ converges weakly to U_t where $(U_t)_{t \in \mathbb{R}_+}$ is the unitary solution of

$$dU_t = -\frac{1}{2}V^*V U_t dt + VU_t da_1^0(t) - V^*U_t da_0^1(t)$$

with $V = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$. This equation is the well-known quantum Langevin equation associated to the spontaneous decay into the ground state in the Wigner-Weisskopf model for the two-level atom (see [M-R] for example).

IV.5 From completely positive maps to Lindbladians

Recall, from section II.3 that the solution $(u_n)_{n \in \mathbb{N}}$ of the equation

$$u_{n+1} = \mathbb{I}_{n+1} u_n$$

with $u_0 = I$ induces a completely positive evolution on the small system. Namely, in the Heisenberg picture, one has for any a, b in \mathcal{H}_0 , any X in $\mathcal{B}(\mathcal{H}_0)$,

$$\langle a \otimes \Omega, u_n^* X u_n b \otimes \Omega \rangle = \langle a, \ell^n(X) b \rangle$$

where

$$\ell(X) = \sum_{i \in \mathbb{N}} (\mathbb{I}_i^0)^* X \mathbb{I}_i^0.$$

Theorem 21. – Let $\mathbb{I}(h) = (\mathbb{I}_j^i(h))_{i,j=0,\dots,n}$ be a family of matrices such that $\sum_i \mathbb{I}_i^0(h)^* \mathbb{I}_i^0(h) = I$ and such that

- $(\mathbb{I}_0^0(h) - I)/h$ converges to some L_0^0 ,
- \mathbb{I}_i^0/\sqrt{h} converges to some L_i^0 for all $i = 1, \dots, n$.

Then there exists a self-adjoint operator H in $\mathcal{B}(\mathcal{H}_0)$ such that for all $t \in \mathbb{R}^+$,

$$\ell^{[t/h]} \longrightarrow e^{t\mathcal{L}}$$

in operator norm, where \mathcal{L} is the Lindblad generator

$$\mathcal{L}(X) = i[H, X] + \frac{1}{2} \sum_{i \in \mathbb{N}} \left(2L_i^{0*} X L_i^0 - L_i^{0*} L_i^0 X - X L_i^{0*} L_i^0 \right)$$

and ℓ is defined above.

Proof

It is a straightforward computation that

$$\ell(X) = X + h(L_0^{0*} X + X L_0^0) + h \sum_{i=1}^n L_i^{0*} X L_i^0 + o(h \|X\|).$$

The equality $\ell(I) = I$ entails

$$(L_0^{0*} + L_0^0) + \sum_{i \in \mathcal{N}} L_i^{0*} L_i^0 = 0$$

so that

$$i \left(L_0^0 + \frac{1}{2} \sum_{i \in \mathcal{N}} L_i^{0*} L_i^0 \right)$$

is self-adjoint. We denote it by H . Then ℓ is of the form

$$\ell(X) = X + h \left(i[H, X] + \frac{1}{2} \sum_{i \in \mathcal{N}} \left(2L_i^{0*} X L_i^0 - L_i^{0*} L_i^0 X - X L_i^{0*} L_i^0 \right) \right) + o(h \|X\|),$$

which is, with the notations of the statements

$$\ell(X) = X + h\mathcal{L}(X) + o(h \|X\|).$$

The above mentioned convergence is therefore clear. ■

As a consequence, it is very easy to obtain approximations of solutions of continuous-time master equations:

$$\frac{dX_t}{dt} = \mathcal{L}(X_t)$$

by solutions of discrete-time ones:

$$x_{n+1} = \ell(x_n).$$

Yet notice that the master equation gives no information whatsoever on the interaction between the small system and the environment or on the environment itself. On the other hand, the associated quantum Langevin equation contains the information of the whole system; this justifies our effort.

Notice that, in the above proposition, no hypothesis is needed on the other coefficients of the matrix $\mathbb{L}(h)$. Their properties actually depend on the choice of additional features of the matrices $\mathbb{L}(h)$, for example their unitarity. The possibility of choosing $\mathbb{L}(h)$ to be unitary and obtain in the end the desired Lindbladian \mathcal{L} is described by Parthasarathy in exercises 29.12 and 29.13 of [Par].

What's more, these manipulations show that the hypotheses of convergence of Theorem 17 are not as artificial as it seems, and are not only convenient assumptions we set up in order to obtain the right convergence. Indeed, to $\mathbb{L}(h)$ is associated both a dynamic on the observables, as we have seen, and an evolution τ , defined by

$$\langle a, \tau_n b \rangle = \langle a \otimes \Omega, u_n b \otimes \Omega \rangle$$

for all a, b in \mathcal{H}_0 , which turns out to be

$$\tau_n = (\mathbb{L}_0^0)^n.$$

If one assumes that $\tau_{[t/h]}$ converges for almost all t and that \mathbb{L}_0^0 is assumed to be continuous at $h = 0$ then the assumption on \mathbb{L}_0^0 in Theorem 17 is to be fulfilled; this

implies that $\sum_{i=1}^n \mathbb{L}_0^{i*} \mathbb{L}_0^i = -h(L_0^{0*} + L_0^0) + o(h)$, so that the other assumptions of convergence of Theorem 17 are natural.

The other conditions described in Theorem 13 are in turn necessary if one wants the process $(U_t)_{t \geq 0}$ obtained in the limit to be unitary (see Theorem 12) or alternatively the matrices $\mathbb{L}(h)$ to be sufficiently close to unitarity.

Remark:

All the constructions and proofs of this article rely heavily on the fact that the reference state for the photon chain was chosen to be a pure state, that is a vector state Ω . This assumption is here essential for the construction of our countable and continuous tensor products. The generalization of this work to the case of an arbitrary state (i.e. a density matrix) has been explored in [AJ2].

References

- [AFL]: Accardi L., Frigerio A., Lu Y.G. “Quantum Langevin equation in the weak coupling limit”, *Quantum probability and applications V* (Heidelberg, 1988), 1–16, Lecture Notes in Math., 1442, Springer, Berlin, 1990.
- [AGL]: Accardi L., Gough J., Lu Y. G. “On the stochastic limit for quantum theory”, *Proceedings of the XXVII Symposium on Mathematical Physics* (Toruń, 1994). *Rep. Math. Phys.* 36 (1995), no. 2-3, 155–187.
- [ALV]: Accardi L., Lu Y. G., Volovich I. “*Quantum theory and its stochastic limit*”, Springer-Verlag, Berlin, 2002.
- [A-L]: Accardi L., Lu Y. G. “The low density limit and the quantum Poisson process”, *Probability theory and mathematical statistics*, Vol. I (Vilnius, 1989), 1–27, ”Mokslas”, Vilnius, 1990.
- [APV]: Accardi L., Pechen A.N., Volovich I.V. “A stochastic golden rule and quantum Langevin equation for low density limit”, *preprint*
- [AFL]: Accardi L., Frigerio A., Lu Y. G. “The weak coupling limit as a quantum functional central limit”, *Comm. Math. Phys.* 131 (1990), no. 3, 537–570.
- [At1]: Attal S. “Extensions of the quantum stochastic calculus” *Quantum Probability Communications XI*, 1-38, World Scientific, 2003.
- [At2]: Attal S. “Approximating the Fock space with the toy Fock space” *Séminaire de Probabilités XXXVI*, 477–491, Lecture Notes in Math., 1801, Springer, Berlin, 2003.
- [At3]: Attal S. “*Quantum Noise Theory*”, book in preparation, Springer verlag.
- [A-L]: Attal S., Lindsay J.M. “Quantum stochastic calculus with maximal operator domains”, *The Annals of Probability*, to appear.
- [AJ1]: Attal S., Joye A. “Weak coupling and continuous limits for repeated quantum interactions”, *preprint*.
- [AJ2]: Attal S., Joye A. “The Langevin equation for a quantum heat bath”, *preprint*.
- [A-M]: Attal S., Meyer P.-A. “Interprétation probabiliste et extension des intégrales stochastiques non commutatives”, *Séminaire de Probabilités XXVII*, 312–327, Lecture Notes in Math., 1557, Springer, Berlin, 1993.
- [A-P]: Attal S., Pautrat Y. “From $(n + 1)$ -level atom chains to n dimensional noises”, *Annales de l’Institut Henri Poincaré* to appear.

- [Ba1]: Barchielli A. “Measurement theory and stochastic differential equations in quantum mechanics”, *Phys. Rev. A* (3) 34 (1986), no. 3, 1642–1649.
- [Ba2]: Barchielli A. “Some stochastic differential equations in quantum optics and measurement theory: the case of counting processes”, *Stochastic evolution of quantum states in open systems and in measurement processes* (Budapest, 1993), 1–14, World Sci. Publishing, River Edge, NJ, 1994.
- [B-B]: Barchielli A., Belavkin V.P. “Measurements continuous in time and a posteriori states in quantum mechanics”, *J. Phys. A* 24 (1991), no. 7, 1495–1514.
- [BRSW]: Bellissard J., Rebolledo R., Spohner D., von Waldenfels W. “The Quantum Flow of Electronic transport”, *preprint*.
- [Coq]: Coquio A., “Why are there only three quantum noises?”, *Probab. Theory Related Fields* 118 (2000), no. 3, 349–364.
- [Dav]: Davies E.B., Markovian master equations. *Comm. Math. Phys.* 39 (1974), 91–110.
- [D-J]: Dereziński J., Jakšić V. “On the nature of Fermi Golden Rule for open quantum systems”, *J. Stat. Phys.*, to appear.
- [F-R]: Fagnola F., Rebolledo R. “A view on stochastic differential equations derived from quantum optics”, *Stochastic models* (Guanajuato, 1998), 193–214, *Aportaciones Mat. Investig.*, 14, Soc. Mat. Mexicana, Mexico, 1998.
- [FRS]: Fagnola F., Rebolledo R., Saavedra C. “Quantum flows associated to master equations in quantum optics”, *J. Math. Phys.* 35 (1994), no. 1, 1–12.
- [FKM]: Ford G. W., Kac M., Mazur P. “Statistical mechanics of assemblies of coupled oscillators”, *J. Mathematical Phys.* 6 1965 504–515.
- [FLO]: Ford G. W., Lewis J. T., O’Connell R. F. “Quantum Langevin equation”, *Phys. Rev. A* (3) 37 (1988), no. 11, 4419–4428.
- [G-Z]: Gardiner C. W., Zoller P. *Quantum noise. A handbook of Markovian and non-Markovian quantum stochastic methods with applications to quantum optics.*, Second edition. Springer Series in Synergetics. Springer-Verlag, Berlin, 2000.
- [GSI]: Grabert H., Schramm P., Ingold G-L “Quantum Brownian motion: the functional integral approach”, *Phys. Rep.* 168 (1988), no. 3, 115–207.
- [Gui]: Guichardet A. *Symmetric Hilbert spaces and related topics.*, Lecture Notes in Mathematics, Vol. 261. Springer-Verlag, Berlin-New York, 1972.
- [H-P]: Hudson R. L., Parthasarathy K. R. “Quantum Ito’s formula and stochastic evolutions”, *Comm. Math. Phys.* 93 (1984), no. 3, 301–323.

- [L-M]: Lindsay J.M., Maassen H. “Stochastic calculus for quantum Brownian motion of nonminimal variance—an approach using integral-sum kernel operators”, *Mark Kac Seminar on Probability and Physics Syllabus 1987–1992* (Amsterdam, 1987–1992), 97–167, CWI Syllabi, 32, Math. Centrum, Centrum Wisk. Inform., Amsterdam, 1992.
- [M-R]: Maassen H., Robinson P. “Quantum stochastic calculus and the dynamical Stark effect”, *Rep. Math. Phys.* 30 (1991), no. 2, 185–203 (1992).
- [Par]: Parthasarathy K.R. “*An introduction to quantum stochastic calculus*”, Monographs in Mathematics 85, Birkhäuser (1992).
- [Pa1]: Pautrat Y., “From Pauli matrices to quantum Ito formula”, *Mathematical Physics, Analysis and Geometry*, to appear.
- [Pa2]: Pautrat Y., “Kernel and integral representations of operators on infinite dimensional toy Fock space”, *Séminaire de Probabilités*, to appear.
- [Pa3]: Pautrat Y., “Des matrices de Pauli aux bruits quantiques”, *Thèse de doctorat de l’Université Joseph Fourier* (2003).