

Random Matrices and Their Limits

Roland Speicher

Saarland University
Saarbrücken, Germany

supported by ERC Advanced Grant
“Non-Commutative Distributions in Free Probability”



European Research Council
Co-funded by the European Commission

Section 1

Random Matrices and Operators





Oberwolfach workshop “Random Matrices” 2000

Deterministic limits of random matrices

Fundamental observation

Many random matrices show, via concentration, for $N \rightarrow \infty$ almost surely a *deterministic* behaviour.



Deterministic limits of random matrices

Fundamental observation

Many random matrices show, via concentration, for $N \rightarrow \infty$ almost surely a *deterministic* behaviour.

Interesting observation

Many random matrices show, via concentration, for $N \rightarrow \infty$ almost surely a deterministic *interesting* behaviour.



Deterministic limits of random matrices

Fundamental observation

Many random matrices show, via concentration, for $N \rightarrow \infty$ almost surely a *deterministic* behaviour.

Interesting observation

Many random matrices show, via concentration, for $N \rightarrow \infty$ almost surely a deterministic *interesting* behaviour.

Interesting observation for the operator algebraic inclined

Many random matrices show, via concentration, for $N \rightarrow \infty$ almost surely a deterministic behaviour, which can be described by interesting *operators on Hilbert spaces* (or their generated C^* -algebras or von Neumann algebras)

Random matrices and operators

Fundamental observation of Voiculescu (1991)



Limit of random matrices can often be described by “nice” and “interesting” operators on Hilbert spaces (which, in the case of several matrices, describe interesting von Neumann algebras)

Convergence of random matrices

$$\begin{array}{ccc}
 (X_1^{(N)}, \dots, X_m^{(N)}) & \longrightarrow & (x_1, \dots, x_m) \\
 \text{random matrices} & \text{almost surely} & \text{operators} \\
 (M_N(\mathbb{C}), \text{tr}) & & (\mathcal{A}, \tau), \mathcal{A} \subset B(\mathcal{H})
 \end{array}$$

Convergence of random matrices

$$\begin{array}{ccc}
 (X_1^{(N)}, \dots, X_m^{(N)}) & \longrightarrow & (x_1, \dots, x_m) \\
 \text{random matrices} & \text{almost surely} & \text{operators} \\
 (M_N(\mathbb{C}), \text{tr}) & & (\mathcal{A}, \tau), \mathcal{A} \subset B(\mathcal{H})
 \end{array}$$

Convergence in distribution

We have for all polynomials p in m non-commuting variables

$$\lim_{N \rightarrow \infty} \text{tr}[p(X_1^{(N)}, \dots, X_m^{(N)})] = \tau[p(x_1, \dots, x_m)]$$

Convergence of random matrices

$$\begin{array}{ccc}
 (X_1^{(N)}, \dots, X_m^{(N)}) & \longrightarrow & (x_1, \dots, x_m) \\
 \text{random matrices} & \text{almost surely} & \text{operators} \\
 (M_N(\mathbb{C}), \text{tr}) & & (\mathcal{A}, \tau), \mathcal{A} \subset B(\mathcal{H})
 \end{array}$$

Convergence in distribution

We have for all polynomials p in m non-commuting variables

$$\lim_{N \rightarrow \infty} \text{tr}[p(X_1^{(N)}, \dots, X_m^{(N)})] = \tau[p(x_1, \dots, x_m)]$$

Strong convergence

- convergence in distribution
- and for all polynomials p

$$\lim_{N \rightarrow \infty} \|p(X_1^{(N)}, \dots, X_m^{(N)})\| = \|p(x_1, \dots, x_m)\|$$

Our most beloved example:
independent GUE \rightarrow free semicirculars



European Research Council

Co-funded by the European Commission

Our most beloved example: independent GUE \rightarrow free semicirculars

One-matrix case: classical commuting case

Note: $X_N \rightarrow x$ means that all moments of X_N converge to corresponding moments of x , hence the distributions (in classical sense of probability measures on \mathbb{R}) converge



European Research Council
Funded by the European Commission

Our most beloved example: independent GUE \rightarrow free semicirculars

One-matrix case: classical commuting case

Note: $X_N \rightarrow x$ means that all moments of X_N converge to corresponding moments of x , hence the distributions (in classical sense of probability measures on \mathbb{R}) converge

... so we can just look on pictures of distributions ...



Our most beloved example: independent GUE \rightarrow free semicirculars

One-matrix case: classical commuting case

Note: $X_N \rightarrow x$ means that all moments of X_N converge to corresponding moments of x , hence the distributions (in classical sense of probability measures on \mathbb{R}) converge

... so we can just look on pictures of distributions ...

Multi-matrix case: non-commutative case

$(X_N, Y_N) \rightarrow (x, y)$ means convergence of all moments, but this does not correspond to convergence of probability measures on \mathbb{R}^2

Our most beloved example: independent GUE \rightarrow free semicirculars

One-matrix case: classical commuting case

Note: $X_N \rightarrow x$ means that all moments of X_N converge to corresponding moments of x , hence the distributions (in classical sense of probability measures on \mathbb{R}) converge

... so we can just look on pictures of distributions ...


Multi-matrix case: non-commutative case

$(X_N, Y_N) \rightarrow (x, y)$ means convergence of all moments, but this does not correspond to convergence of probability measures on \mathbb{R}^2

... if we still want to see classical objects and pictures we can look on $p(X_N, Y_N) \rightarrow p(x, y)$ for (sufficiently many) functions p in X_N, Y_N ...


One-matrix case: classical random matrix case

X_N GUE random matrix



One-matrix case: classical random matrix case

X_N GUE random matrix



$$x = l + l^*,$$

l one-sided shift on

$$\bigoplus_{n \geq 0} \mathbb{C}e_n$$


$$le_n = e_{n+1}$$

$$l^*e_{n+1} = e_n, \quad l^*e_0 = 0$$

$$\tau(a) = \langle e_0, ae_0 \rangle$$

One-matrix case: classical random matrix case

X_N GUE random matrix



- $X_N \rightarrow x$ in distribution (Wigner 1955)

$$x = l + l^*,$$

l one-sided shift on

$$\bigoplus_{n \geq 0} \mathbb{C}e_n$$


$$le_n = e_{n+1}$$

$$l^*e_{n+1} = e_n, l^*e_0 = 0$$

$$\tau(a) = \langle e_0, ae_0 \rangle$$

One-matrix case: classical random matrix case

X_N GUE random matrix



- $X_N \rightarrow x$ in distribution (Wigner 1955)
- $\|X_N\| \rightarrow \|x\| = 2$ (Füredi, Komlós 1981)

$$x = l + l^*,$$

l one-sided shift on

$$\bigoplus_{n \geq 0} \mathbb{C}e_n$$

$$le_n = e_{n+1}$$

$$l^*e_{n+1} = e_n, l^*e_0 = 0$$

$$\tau(a) = \langle e_0, ae_0 \rangle$$

Multi-matrix case: non-commutative case

X_N, Y_N independent GUE



Multi-matrix case: non-commutative case

X_N, Y_N independent GUE

$$x = l_1 + l_1^*, y = l_2 + l_2^*$$

two copies of one-sided shift in different directions (creation and annihilation operators on full Fock space; Cuntz algebra)

$$\tau(a) = \langle \Omega, a\Omega \rangle$$

Multi-matrix case: non-commutative case

X_N, Y_N independent GUE

$$x = l_1 + l_1^*, y = l_2 + l_2^*$$

two copies of one-sided shift in different directions (creation and annihilation operators on full Fock space; Cuntz algebra)

$$\tau(a) = \langle \Omega, a\Omega \rangle$$

- $X_N, Y_N \rightarrow x, y$ in distribution (Voiculescu 1991)

Multi-matrix case: non-commutative case


X_N, Y_N independent GUE

$$p(x, y) = xy + yx + x^2$$

$$x = l_1 + l_1^*, y = l_2 + l_2^*$$

two copies of one-sided shift in different directions (creation and annihilation operators on full Fock space; Cuntz algebra)

$$\tau(a) = \langle \Omega, a\Omega \rangle$$



- $X_N, Y_N \rightarrow x, y$ in distribution (Voiculescu 1991)
- $p(X_N, Y_N) \rightarrow p(x, y)$ in distribution

Multi-matrix case: non-commutative case


X_N, Y_N independent GUE

$$p(x, y) = xy + yx + x^2$$

$$x = l_1 + l_1^*, y = l_2 + l_2^*$$

two copies of one-sided shift in different directions (creation and annihilation operators on full Fock space; Cuntz algebra)

$$\tau(a) = \langle \Omega, a\Omega \rangle$$



- $X_N, Y_N \rightarrow x, y$ in distribution (Voiculescu 1991)
- $p(X_N, Y_N) \rightarrow p(x, y)$ in distribution
- $\|p(X_N, Y_N)\| \rightarrow \|p(x, y)\|$ (Haagerup and Thorbjørnsen 2005)

Goal: Go over from Polynomials to More General Classes of Functions

For $m = 1$, one has

for all continuous f :

- $\lim_{N \rightarrow \infty} \text{tr}[f(X^{(N)})] = \tau[f(x)]$
- $\lim_{N \rightarrow \infty} \|f(X^{(N)})\| = \|f(x)\|$

Goal: Go over from Polynomials to More General Classes of Functions

For $m = 1$, one has

for all continuous f :

- $\lim_{N \rightarrow \infty} \text{tr}[f(X^{(N)})] = \tau[f(x)]$
- $\lim_{N \rightarrow \infty} \|f(X^{(N)})\| = \|f(x)\|$

Which classes of functions in non-commuting variables

- polynomials !!!

Goal: Go over from Polynomials to More General Classes of Functions

For $m = 1$, one has

for all continuous f :

- $\lim_{N \rightarrow \infty} \text{tr}[f(X^{(N)})] = \tau[f(x)]$
- $\lim_{N \rightarrow \infty} \|f(X^{(N)})\| = \|f(x)\|$

Which classes of functions in non-commuting variables

- continuous ???

- polynomials !!!

Goal: Go over from Polynomials to More General Classes of Functions

For $m = 1$, one has

for all continuous f :

- $\lim_{N \rightarrow \infty} \text{tr}[f(X^{(N)})] = \tau[f(x)]$
- $\lim_{N \rightarrow \infty} \|f(X^{(N)})\| = \|f(x)\|$

Which classes of functions in non-commuting variables

- continuous ???
- analytic \rightsquigarrow free analysis
- polynomials !!!

Goal: Go over from Polynomials to More General Classes of Functions

For $m = 1$, one has

for all continuous f :

- $\lim_{N \rightarrow \infty} \text{tr}[f(X^{(N)})] = \tau[f(x)]$
- $\lim_{N \rightarrow \infty} \|f(X^{(N)})\| = \|f(x)\|$

Which classes of functions in non-commuting variables

- continuous ???
- analytic \rightsquigarrow free analysis
- rational !!!
- polynomials !!!

Section 2

Non-Commutative Rational Functions



Non-commutative rational functions

Non-commutative rational functions (Amitsur 1966, Cohn 1971)

- A rational function $r(y_1, \dots, y_m)$ in non-commuting variables y_1, \dots, y_m is anything we can get by algebraic operations, including inverses, from y_1, \dots, y_m ,

Non-commutative rational functions

Non-commutative rational functions (Amitsur 1966, Cohn 1971)

- A rational function $r(y_1, \dots, y_m)$ in non-commuting variables y_1, \dots, y_m is anything we can get by algebraic operations, including inverses, from y_1, \dots, y_m , like

$$(4 - y_1)^{-1} + (4 - y_1)^{-1}y_2((4 - y_1) - y_2(4 - y_1)^{-1}y_2)^{-1}y_2(4 - y_1)^{-1}$$

Non-commutative rational functions

Non-commutative rational functions (Amitsur 1966, Cohn 1971)

- A rational function $r(y_1, \dots, y_m)$ in non-commuting variables y_1, \dots, y_m is anything we can get by algebraic operations, including inverses, from y_1, \dots, y_m , like

$$(4 - y_1)^{-1} + (4 - y_1)^{-1}y_2((4 - y_1) - y_2(4 - y_1)^{-1}y_2)^{-1}y_2(4 - y_1)^{-1}$$

modulo

- ▶ identifying “algebraically equivalent” expressions
- ▶ not inverting 0

Non-commutative rational functions

Non-commutative rational functions (Amitsur 1966, Cohn 1971)

- A rational function $r(y_1, \dots, y_m)$ in non-commuting variables y_1, \dots, y_m is anything we can get by algebraic operations, including inverses, from y_1, \dots, y_m , like

$$(4 - y_1)^{-1} + (4 - y_1)^{-1}y_2((4 - y_1) - y_2(4 - y_1)^{-1}y_2)^{-1}y_2(4 - y_1)^{-1}$$

modulo

- ▶ identifying “algebraically equivalent” expressions
- ▶ not inverting 0

this is not obvious to decide, e.g., one has rational identities like

$$y_2^{-1} + y_2^{-1}(y_3^{-1}y_1^{-1} - y_2^{-1})^{-1}y_2^{-1} - (y_2 - y_3y_1)^{-1} = 0$$

Non-commutative rational functions

Non-commutative rational functions (Amitsur 1966, Cohn 1971)

- A rational function $r(y_1, \dots, y_m)$ in non-commuting variables y_1, \dots, y_m is anything we can get by algebraic operations, including inverses, from y_1, \dots, y_m , like

$$(4 - y_1)^{-1} + (4 - y_1)^{-1}y_2((4 - y_1) - y_2(4 - y_1)^{-1}y_2)^{-1}y_2(4 - y_1)^{-1}$$

modulo

- ▶ identifying “algebraically equivalent” expressions
- ▶ not inverting 0

this is not obvious to decide, e.g., one has rational identities like

$$y_2^{-1} + y_2^{-1}(y_3^{-1}y_1^{-1} - y_2^{-1})^{-1}y_2^{-1} - (y_2 - y_3y_1)^{-1} = 0$$

- after all, it works and gives a skew field

$$\mathbb{C}\langle y_1, \dots, y_m \rangle \quad \text{“free field”}$$

A rational function $r(y_1, \dots, y_m)$ in non-commuting variables y_1, \dots, y_m

- can be realized more systematically by going over to matrices

$$r(y_1, \dots, y_m) = uQ^{-1}v$$

where, for some N ,

- ▶ u is $1 \times N$
- ▶ Q is $N \times N$ and invertible in $M_N(\mathbb{C}\langle y_1, \dots, y_m \rangle)$
- ▶ v is $N \times 1$
- ▶ and all entries are polynomials (can actually be chosen with degree ≤ 1)

A rational function $r(y_1, \dots, y_m)$ in non-commuting variables y_1, \dots, y_m

- can be realized more systematically by going over to matrices

$$r(y_1, \dots, y_m) = uQ^{-1}v$$

where, for some N ,

- ▶ u is $1 \times N$
- ▶ Q is $N \times N$ and invertible in $M_N(\mathbb{C}\langle y_1, \dots, y_m \rangle)$
- ▶ v is $N \times 1$
- ▶ and all entries are polynomials (can actually be chosen with degree ≤ 1)

$$\begin{aligned} & (4 - y_1)^{-1} + (4 - y_1)^{-1}y_2 \left((4 - y_1) - y_2(4 - y_1)^{-1}y_2 \right)^{-1} y_2(4 - y_1)^{-1} \\ &= \begin{pmatrix} \frac{1}{2} & 0 \end{pmatrix} \cdot \begin{pmatrix} -1 + \frac{1}{4}y_1 & \frac{1}{4}y_2 \\ \frac{1}{4}y_2 & -1 + \frac{1}{4}y_1 \end{pmatrix}^{-1} \cdot \begin{pmatrix} \frac{1}{2} \\ 0 \end{pmatrix} \end{aligned}$$

A rational function $r(y_1, \dots, y_m)$ in non-commuting variables y_1, \dots, y_m

- can be realized more systematically by going over to matrices

$$r(y_1, \dots, y_m) = uQ^{-1}v$$

where, for some N ,

- ▶ u is $1 \times N$
- ▶ Q is $N \times N$ and invertible in $M_N(\mathbb{C}\langle y_1, \dots, y_m \rangle)$
- ▶ v is $N \times 1$
- ▶ and all entries are polynomials (can actually be chosen with degree ≤ 1)

$$\begin{aligned} & (4 - y_1)^{-1} + (4 - y_1)^{-1}y_2 \left((4 - y_1) - y_2(4 - y_1)^{-1}y_2 \right)^{-1} y_2(4 - y_1)^{-1} \\ &= \begin{pmatrix} \frac{1}{2} & 0 \end{pmatrix} \cdot \begin{pmatrix} -1 + \frac{1}{4}y_1 & \frac{1}{4}y_2 \\ \frac{1}{4}y_2 & -1 + \frac{1}{4}y_1 \end{pmatrix}^{-1} \cdot \begin{pmatrix} \frac{1}{2} \\ 0 \end{pmatrix} \end{aligned}$$

- this is essentially (in the case of polynomials) the “linearization trick” which we use in free probability, for example, to calculate distributions of polynomials in free variables

Historical remark

Note that this linearization trick is a well-known idea in many other mathematical communities, known under various names like

- Higman's trick (Higman "The units of group rings", 1940)
- recognizable power series (automata theory, Kleene 1956, Schützenberger 1961)
- linearization by enlargement (ring theory, Cohn 1985; Cohn and Reutenauer 1994, Malcolmson 1978)
- descriptor realization (control theory, Kalman 1963; Helton, McCullough, Vinnikov 2006)
- linearization trick (Haagerup, Thorbjørnsen 2005 (+Schultz 2006); Anderson 2012)

Applying nc rational functions to operators

... there are a couple of issues arising ...

Not every relation in $\mathbb{C}\langle y_1, \dots, y_m \rangle$ must necessarily hold in any algebra (even if it makes sense there)

Applying nc rational functions to operators

... there are a couple of issues arising ...

Not every relation in $\mathbb{C}\langle y_1, \dots, y_m \rangle$ must necessarily hold in any algebra (even if it makes sense there)

- In $\mathbb{C}\langle y_1, y_2 \rangle$ we have

$$y_1(y_2y_1)^{-1}y_2 = 1$$

Applying nc rational functions to operators

... there are a couple of issues arising ...

Not every relation in $\mathbb{C}\langle y_1, \dots, y_m \rangle$ must necessarily hold in any algebra (even if it makes sense there)

- In $\mathbb{C}\langle y_1, y_2 \rangle$ we have

$$y_1(y_2y_1)^{-1}y_2 = 1$$

- Let v be an isometry which is not unitary, i.e. $vv^* = 1$, $v^*v \neq 1$ Then we have

$$v^*(vv^*)^{-1}v = v^*v \neq 1$$

Applying nc rational functions to operators

... there are a couple of issues arising ...

Not every relation in $\mathbb{C}\langle y_1, \dots, y_m \rangle$ must necessarily hold in any algebra (even if it makes sense there)

- In $\mathbb{C}\langle y_1, y_2 \rangle$ we have

$$y_1(y_2y_1)^{-1}y_2 = 1$$

- Let v be an isometry which is not unitary, i.e. $vv^* = 1$, $v^*v \neq 1$ Then we have

$$v^*(vv^*)^{-1}v = v^*v \neq 1$$

- Solution: By Cohn, we know that the above is the only issue, hence we are okay if we avoid non-unitary isometries (also in matrices over the algebra);

Applying nc rational functions to operators

... there are a couple of issues arising ...

Not every relation in $\mathbb{C}\langle y_1, \dots, y_m \rangle$ must necessarily hold in any algebra (even if it makes sense there)

- In $\mathbb{C}\langle y_1, y_2 \rangle$ we have

$$y_1(y_2y_1)^{-1}y_2 = 1$$

- Let v be an isometry which is not unitary, i.e. $vv^* = 1$, $v^*v \neq 1$ Then we have

$$v^*(vv^*)^{-1}v = v^*v \neq 1$$

- Solution: By Cohn, we know that the above is the only issue, hence we are okay if we avoid non-unitary isometries (also in matrices over the algebra);
 - ▶ hence only work in stably finite algebras

Applying nc rational functions to operators

... there are a couple of issues arising ...

Not every relation in $\mathbb{C}\langle y_1, \dots, y_m \rangle$ must necessarily hold in any algebra (even if it makes sense there)

- In $\mathbb{C}\langle y_1, y_2 \rangle$ we have

$$y_1(y_2y_1)^{-1}y_2 = 1$$

- Let v be an isometry which is not unitary, i.e. $vv^* = 1$, $v^*v \neq 1$ Then we have

$$v^*(vv^*)^{-1}v = v^*v \neq 1$$

- Solution: By Cohn, we know that the above is the only issue, hence we are okay if we avoid non-unitary isometries (also in matrices over the algebra);
 - ▶ hence only work in stably finite algebras
 - ▶ this is the case in our situations, where we have a tracial state

Applying nc rational functions to operators

... there are a couple of issues arising ...

- In $\mathbb{C}\langle y_1, \dots, y_m \rangle$ every $r(y_1, \dots, y_m) \neq 0$ is invertible. This is of course not true when we apply r to operators:



Applying nc rational functions to operators

... there are a couple of issues arising ...

- In $\mathbb{C}\langle y_1, \dots, y_m \rangle$ every $r(y_1, \dots, y_m) \neq 0$ is invertible. This is of course not true when we apply r to operators:
 - ▶ $r(x_1, \dots, x_m) = 0$ could happen for $0 \neq r \in \mathbb{C}\langle y_1, \dots, y_m \rangle$



Applying nc rational functions to operators

... there are a couple of issues arising ...

- In $\mathbb{C}\langle y_1, \dots, y_m \rangle$ every $r(y_1, \dots, y_m) \neq 0$ is invertible. This is of course not true when we apply r to operators:
 - ▶ $r(x_1, \dots, x_m) = 0$ could happen for $0 \neq r \in \mathbb{C}\langle y_1, \dots, y_m \rangle$
 - ▶ even if $r(x_1, \dots, x_m) \neq 0$, it does not need to be invertible in general



Applying nc rational functions to operators

... there are a couple of issues arising ...

- In $\mathbb{C}\langle y_1, \dots, y_m \rangle$ every $r(y_1, \dots, y_m) \neq 0$ is invertible. This is of course not true when we apply r to operators:
 - ▶ $r(x_1, \dots, x_m) = 0$ could happen for $0 \neq r \in \mathbb{C}\langle y_1, \dots, y_m \rangle$
 - ▶ even if $r(x_1, \dots, x_m) \neq 0$, it does not need to be invertible in general
- Possible solutions:
 - ▶ consider only r for which $r(x_1, \dots, x_m)$ is invertible as bounded operator

Applying nc rational functions to operators

... there are a couple of issues arising ...

- In $\mathbb{C}\langle y_1, \dots, y_m \rangle$ every $r(y_1, \dots, y_m) \neq 0$ is invertible. This is of course not true when we apply r to operators:
 - ▶ $r(x_1, \dots, x_m) = 0$ could happen for $0 \neq r \in \mathbb{C}\langle y_1, \dots, y_m \rangle$
 - ▶ even if $r(x_1, \dots, x_m) \neq 0$, it does not need to be invertible in general
- Possible solutions:
 - ▶ consider only r for which $r(x_1, \dots, x_m)$ is invertible as bounded operator
 - ▶ or allow also unbounded operators



Rational functions of random matrices and their limit

Proposition (Sheng Yin 2017)

Consider random matrices $(X_1^{(N)}, \dots, X_m^{(N)})$ which converge to operators (x_1, \dots, x_m) in the strong sense: for any polynomial $p \in \mathbb{C}\langle x_1, \dots, x_m \rangle$ we have

- $\lim_{N \rightarrow \infty} \text{tr}[p(X_1^{(N)}, \dots, X_m^{(N)})] = \tau[p(x_1, \dots, x_m)]$
- $\lim_{N \rightarrow \infty} \|p(X_1^{(N)}, \dots, X_m^{(N)})\| = \|p(x_1, \dots, x_m)\|$

Rational functions of random matrices and their limit

Proposition (Sheng Yin 2017)

Consider random matrices $(X_1^{(N)}, \dots, X_m^{(N)})$ which converge to operators (x_1, \dots, x_m) in the strong sense: for any polynomial $p \in \mathbb{C}\langle x_1, \dots, x_m \rangle$ we have

- $\lim_{N \rightarrow \infty} \text{tr}[p(X_1^{(N)}, \dots, X_m^{(N)})] = \tau[p(x_1, \dots, x_m)]$
- $\lim_{N \rightarrow \infty} \|p(X_1^{(N)}, \dots, X_m^{(N)})\| = \|p(x_1, \dots, x_m)\|$

Then this strong convergence remains also true for rational functions: Let $r \in \mathbb{C}\langle y_1, \dots, y_m \rangle$, such that $r(x_1, \dots, x_m)$ is defined as bounded operator.

Rational functions of random matrices and their limit

Proposition (Sheng Yin 2017)

Consider random matrices $(X_1^{(N)}, \dots, X_m^{(N)})$ which converge to operators (x_1, \dots, x_m) in the strong sense: for any polynomial $p \in \mathbb{C}\langle x_1, \dots, x_m \rangle$ we have

- $\lim_{N \rightarrow \infty} \text{tr}[p(X_1^{(N)}, \dots, X_m^{(N)})] = \tau[p(x_1, \dots, x_m)]$
- $\lim_{N \rightarrow \infty} \|p(X_1^{(N)}, \dots, X_m^{(N)})\| = \|p(x_1, \dots, x_m)\|$

Then this strong convergence remains also true for rational functions: Let $r \in \mathbb{C}\langle y_1, \dots, y_m \rangle$, such that $r(x_1, \dots, x_m)$ is defined as bounded operator. Then we have almost surely that

- $r(X_1^{(N)}, \dots, X_m^{(N)})$ is defined for sufficiently large N

Rational functions of random matrices and their limit

Proposition (Sheng Yin 2017)

Consider random matrices $(X_1^{(N)}, \dots, X_m^{(N)})$ which converge to operators (x_1, \dots, x_m) in the strong sense: for any polynomial $p \in \mathbb{C}\langle x_1, \dots, x_m \rangle$ we have

- $\lim_{N \rightarrow \infty} \text{tr}[p(X_1^{(N)}, \dots, X_m^{(N)})] = \tau[p(x_1, \dots, x_m)]$
- $\lim_{N \rightarrow \infty} \|p(X_1^{(N)}, \dots, X_m^{(N)})\| = \|p(x_1, \dots, x_m)\|$

Then this strong convergence remains also true for rational functions: Let $r \in \mathbb{C}\langle y_1, \dots, y_m \rangle$, such that $r(x_1, \dots, x_m)$ is defined as bounded operator. Then we have almost surely that

- $r(X_1^{(N)}, \dots, X_m^{(N)})$ is defined for sufficiently large N
- $\lim_{N \rightarrow \infty} \text{tr}[r(X_1^{(N)}, \dots, X_m^{(N)})] = \tau[r(x_1, \dots, x_m)]$
- $\lim_{N \rightarrow \infty} \|r(X_1^{(N)}, \dots, X_m^{(N)})\| = \|r(x_1, \dots, x_m)\|$

Proof

- by recursion on complexity of formulas with respect to inversions
- main step: controlling taking inverse, by approximations by polynomials, uniformly in approximating matrices and limit operators



Proof

- by recursion on complexity of formulas with respect to inversions
- main step: controlling taking inverse, by approximations by polynomials, uniformly in approximating matrices and limit operators

convergence for polynomials \rightsquigarrow convergence for rational functions

strong \implies strong

Proof

- by recursion on complexity of formulas with respect to inversions
- main step: controlling taking inverse, by approximations by polynomials, uniformly in approximating matrices and limit operators

convergence for polynomials \rightsquigarrow convergence for rational functions

strong \implies strong
in distribution

Proof


- by recursion on complexity of formulas with respect to inversions
- main step: controlling taking inverse, by approximations by polynomials, uniformly in approximating matrices and limit operators

convergence for polynomials \rightsquigarrow convergence for rational functions

strong \implies strong
in distribution $\not\Leftarrow$ in distribution


Distribution of random matrices and their limit for

$$r(x, y) = (4 - x)^{-1} + (4 - x)^{-1}y((4 - x) - y(4 - x)^{-1}y)^{-1}y(4 - x)^{-1}$$



Distribution of random matrices and their limit for

$$r(x, y) = (4 - x)^{-1} + (4 - x)^{-1}y((4 - x) - y(4 - x)^{-1}y)^{-1}y(4 - x)^{-1}$$



X_N, Y_N independent GUE


$$x = l_1 + l_1^*, y = l_2 + l_2^*$$

two copies of one-sided shift in different directions (creation and annihilation operators on full Fock space; Cuntz algebra)

$$\tau(a) = \langle \Omega, a\Omega \rangle$$

Distribution of random matrices and their limit for

$$r(x, y) = (4 - x)^{-1} + (4 - x)^{-1}y((4 - x) - y(4 - x)^{-1}y)^{-1}y(4 - x)^{-1}$$



X_N, Y_N independent GUE

$$x = l_1 + l_1^*, y = l_2 + l_2^*$$


two copies of one-sided shift in different directions (creation and annihilation operators on full Fock space; Cuntz algebra)

$$\tau(a) = \langle \Omega, a\Omega \rangle$$

- $r(X_N, Y_N) \rightarrow r(x, y)$ in distribution

Distribution of random matrices and their limit for

$$r(x, y) = (4 - x)^{-1} + (4 - x)^{-1}y((4 - x) - y(4 - x)^{-1}y)^{-1}y(4 - x)^{-1}$$



X_N, Y_N independent GUE

$$x = l_1 + l_1^*, y = l_2 + l_2^*$$

two copies of one-sided shift in different directions (creation and annihilation operators on full Fock space; Cuntz algebra)

$$\tau(a) = \langle \Omega, a\Omega \rangle$$

- $r(X_N, Y_N) \rightarrow r(x, y)$ in distribution
- $\|r(X_N, Y_N)\| \rightarrow \|r(x, y)\|$

Section 3

Unbounded Operators



In the limit $(x_1, \dots, x_m) \subset (\mathcal{A}, \tau)$ we are in a $\|1\|$ -situation

- unbounded operators $U(\mathcal{A})$ affiliated to vN algebra \mathcal{A} form a $*$ -algebra
- and $a \in U(\mathcal{A})$ is invertible if and only if a has no zero divisors

In the limit $(x_1, \dots, x_m) \subset (\mathcal{A}, \tau)$ we are in a $\|1\|$ -situation

- unbounded operators $U(\mathcal{A})$ affiliated to vN algebra \mathcal{A} form a $*$ -algebra
- and $a \in U(\mathcal{A})$ is invertible if and only if a has no zero divisors
(a has zero divisor: $\exists b \in U(\mathcal{A})$ with $b \neq 0$ and $ab = 0$)

In the limit $(x_1, \dots, x_m) \subset (\mathcal{A}, \tau)$ we are in a $\|1\|_1$ -situation

- unbounded operators $U(\mathcal{A})$ affiliated to vN algebra \mathcal{A} form a $*$ -algebra
- and $a \in U(\mathcal{A})$ is invertible if and only if a has no zero divisors
(a has zero divisor: $\exists b \in U(\mathcal{A})$ with $b \neq 0$ and $ab = 0$)

we expect

$r(x_1, \dots, x_m)$ for $r \in \mathbb{C}\langle y_1, \dots, y_m \rangle$ is well-defined and has no zero divisors for

- x_1, \dots, x_m free semicirculars
- more general, limit operators of “nice” random matrix models

In the limit $(x_1, \dots, x_m) \subset (\mathcal{A}, \tau)$ we are in a $\|1$ -situation

- unbounded operators $U(\mathcal{A})$ affiliated to vN algebra \mathcal{A} form a $*$ -algebra
- and $a \in U(\mathcal{A})$ is invertible if and only if a has no zero divisors
(a has zero divisor: $\exists b \in U(\mathcal{A})$ with $b \neq 0$ and $ab = 0$)

we expect

$r(x_1, \dots, x_m)$ for $r \in \mathbb{C}\langle y_1, \dots, y_m \rangle$ is well-defined and has no zero divisors for

- x_1, \dots, x_m free semicirculars
- more general, limit operators of “nice” random matrix models
(which should be operators having maximal free entropy dimension)

In the limit $(x_1, \dots, x_m) \subset (\mathcal{A}, \tau)$ we are in a II_1 -situation

- unbounded operators $U(\mathcal{A})$ affiliated to vN algebra \mathcal{A} form a $*$ -algebra
- and $a \in U(\mathcal{A})$ is invertible if and only if a has no zero divisors
(a has zero divisor: $\exists b \in U(\mathcal{A})$ with $b \neq 0$ and $ab = 0$)

we expect

$r(x_1, \dots, x_m)$ for $r \in \mathbb{C}\langle y_1, \dots, y_m \rangle$ is well-defined and has no zero divisors for

- x_1, \dots, x_m free semicirculars
- more general, limit operators of “nice” random matrix models
(which should be operators having maximal free entropy dimension)

we know (by Shlyakhtenko-Skoufranis and Mai-Speicher-Weber)

$p(x_1, \dots, x_m)$ for $p \in \mathbb{C}\langle y_1, \dots, y_m \rangle$ has no zero divisors for

- x_1, \dots, x_m free semicirculars
- x_1, \dots, x_m with maximal free entropy dimension $= m$

What do we expect of nice operators

$(X_1^{(N)}, \dots, X_m^{(N)})$ \rightarrow (x_1, \dots, x_m)
nice random matrices \rightarrow nice operators

What do we expect of nice operators

$(X_1^{(N)}, \dots, X_m^{(N)})$ \rightarrow (x_1, \dots, x_m)
 nice random matrices \rightarrow nice operators

limit operators should be without algebraic relations in a very general sense

What do we expect of nice operators

$(X_1^{(N)}, \dots, X_m^{(N)})$ \rightarrow (x_1, \dots, x_m)
 nice random matrices \rightarrow nice operators

limit operators should be without algebraic relations in a very general sense

- we know
 - ▶ no polynomial relations
 - ▶ no “local” polynomial relations

What do we expect of nice operators

$(X_1^{(N)}, \dots, X_m^{(N)})$
 nice random matrices \rightarrow (x_1, \dots, x_m)
 \rightarrow nice operators

limit operators should be without algebraic relations in a very general sense

- we know
 - ▶ no polynomial relations
 - ▶ no “local” polynomial relations
- we don't know (but would like to)
 - ▶ no rational relations
 - ▶ no “local rational” relations

What do we expect of nice operators

$(X_1^{(N)}, \dots, X_m^{(N)})$ → (x_1, \dots, x_m)
 nice random matrices → nice operators

limit operators should be without algebraic relations in a very general sense

- we know
 - ▶ no polynomial relations
 - ▶ no “local” polynomial relations
- we don't know (but would like to)
 - ▶ no rational relations
 - ▶ no “local rational” relations

Note

no polynomial relations $\not\Rightarrow$ no rational relations

Question

How much “regularity” of the distribution of $(x_1, \dots, x_m) \subset (\mathcal{A}, \tau)$ is necessary to have

$$\mathbb{C}\langle x_1, \dots, x_m \rangle \equiv \text{free field}$$

division closure in unbounded operators

Question

How much “regularity” of the distribution of $(x_1, \dots, x_m) \subset (\mathcal{A}, \tau)$ is necessary to have

$$\mathbb{C}\langle x_1, \dots, x_m \rangle \equiv \text{free field}$$

division closure in unbounded operators

Question: Consider $(X_1^{(N)}, \dots, X_m^{(N)}) \rightarrow (x_1, \dots, x_m)$

Assume that

- the random matrices are very nice (like independent GUE),
- and that $r(x_1, \dots, x_m)$ makes sense as unbounded operator.

Question

How much “regularity” of the distribution of $(x_1, \dots, x_m) \subset (\mathcal{A}, \tau)$ is necessary to have

$$\mathbb{C}\langle x_1, \dots, x_m \rangle \equiv \text{free field}$$

division closure in unbounded operators

Question: Consider $(X_1^{(N)}, \dots, X_m^{(N)}) \rightarrow (x_1, \dots, x_m)$

Assume that

- the random matrices are very nice (like independent GUE),
- and that $r(x_1, \dots, x_m)$ makes sense as unbounded operator.

When do we have almost surely:

- $r(X_1^{(N)}, \dots, X_m^{(N)})$ makes sense
- $\text{tr}[r(X_1^{(N)}, \dots, X_m^{(N)})] \rightarrow \tau[r(x_1, \dots, x_m)]$

Question

How much “regularity” of the distribution of $(x_1, \dots, x_m) \subset (\mathcal{A}, \tau)$ is necessary to have

$$\mathbb{C}\langle x_1, \dots, x_m \rangle \equiv \text{free field}$$

division closure in unbounded operators

Question: Consider $(X_1^{(N)}, \dots, X_m^{(N)}) \rightarrow (x_1, \dots, x_m)$

Assume that

- the random matrices are very nice (like independent GUE),
- and that $r(x_1, \dots, x_m)$ makes sense as unbounded operator.

When do we have almost surely:

- $r(X_1^{(N)}, \dots, X_m^{(N)})$ makes sense
- $\text{tr}[r(X_1^{(N)}, \dots, X_m^{(N)})] \rightarrow \tau[r(x_1, \dots, x_m)]$

Thank you!

Distribution of random matrices and their limit for

$$p(x_1, x_2, x_3, x_4) = x_1x_2 + x_2x_3 + x_3x_4 + x_4x_1$$

