

A SMALL SOLID BODY WITH LARGE DENSITY IN A PLANAR FLUID IS NEGLIGIBLE

JIAO HE AND DRAGOȘ IFTIMIE

ABSTRACT. In this article, we consider a small rigid body moving in a viscous fluid filling the whole \mathbb{R}^2 . We assume that the diameter of the rigid body goes to 0, that the initial velocity has bounded energy and that the density of the rigid body goes to infinity. We prove that the rigid body has no influence on the limit equation by showing convergence of the solutions towards a solution of the Navier-Stokes equations in the full plane \mathbb{R}^2 .

1. INTRODUCTION

In this paper, we consider a fluid-solid system consisting in a small smooth rigid body Ω_ε of size ε evolving in a viscous fluid filling the whole of \mathbb{R}^2 . Our aim is to determine the limit of this coupled system when the size of the rigid body ε goes to 0.

Let us describe now the fluid solid system of equations. To do that, we need to introduce some notation. We denote by u_ε , respectively p_ε , the velocity, respectively the pressure, of the fluid; they are defined on $\mathbb{R}^2 \setminus \Omega_\varepsilon$, the exterior of the smooth rigid body Ω_ε . The evolution of the rigid body $\Omega_\varepsilon(t)$ is described by h_ε , the position of its center of mass, and by θ_ε , the angle of rotation of the rigid body compared with the initial position. We have that

$$\Omega_\varepsilon(t) = h_\varepsilon(t) + \begin{pmatrix} \cos \theta_\varepsilon(t) & -\sin \theta_\varepsilon(t) \\ \sin \theta_\varepsilon(t) & \cos \theta_\varepsilon(t) \end{pmatrix} (\Omega_\varepsilon(0) - h_\varepsilon(0)).$$

The velocity of the fluid verifies the incompressible Navier-Stokes equations in the exterior of the rigid body:

$$(1) \quad \frac{\partial u_\varepsilon}{\partial t} + u_\varepsilon \cdot \nabla u_\varepsilon - \nu \Delta u_\varepsilon + \nabla p_\varepsilon = 0, \quad \operatorname{div} u_\varepsilon = 0 \quad \text{for } t > 0 \text{ and } x \in \mathbb{R}^2 \setminus \Omega_\varepsilon(t).$$

On the boundary of the rigid body we assume no-slip boundary conditions:

$$(2) \quad u_\varepsilon(t, x) = h'_\varepsilon(t) + \theta'_\varepsilon(t)(x - h_\varepsilon(t))^\perp \text{ for } t > 0 \text{ and } x \in \partial\Omega_\varepsilon(t).$$

Moreover, the velocity is assumed to vanish at infinity:

$$(3) \quad \lim_{|x| \rightarrow \infty} u_\varepsilon(t, x) = 0 \text{ for } t \geq 0.$$

Now we write down the equations of motion of the solid body. Let us denote by m_ε the mass of the solid and by J_ε the momentum of inertia of the solid. We also denote by $\sigma(u_\varepsilon, p_\varepsilon)$ the stress tensor of the fluid:

$$\sigma(u_\varepsilon, p_\varepsilon) = 2\nu D(u_\varepsilon) - p_\varepsilon I_2$$

where I_2 is the identity matrix and $D(u_\varepsilon)$ is the deformation tensor

$$D(u_\varepsilon) = \frac{1}{2} \left(\frac{\partial u_{\varepsilon,i}}{\partial x_j} + \frac{\partial u_{\varepsilon,j}}{\partial x_i} \right)_{i,j}.$$

Then the solid body $\Omega_\varepsilon(t)$ evolves according to Newton's balance law for linear and angular momenta:

$$(4) \quad m_\varepsilon h_\varepsilon''(t) = - \int_{\partial\Omega_\varepsilon(t)} \sigma(u_\varepsilon, p_\varepsilon) n_\varepsilon \quad \text{for } t > 0,$$

and

$$(5) \quad J_\varepsilon \theta_\varepsilon''(t) = - \int_{\partial\Omega_\varepsilon(t)} (\sigma(u_\varepsilon, p_\varepsilon) n_\varepsilon) \cdot (x - h_\varepsilon)^\perp \quad \text{for } t > 0.$$

Above n_ε denotes the unit normal to $\partial\Omega_\varepsilon$ which points to the interior of the rigid body Ω_ε , the orthogonal x^\perp is defined by $x^\perp = (-x_2, x_1)$ and $\sigma(u_\varepsilon, p_\varepsilon) n_\varepsilon$ denotes the matrix $\sigma(u_\varepsilon, p_\varepsilon)$ applied to the vector n_ε .

One can obtain energy estimates for this system of equations. If we formally multiply the equation of u_ε by u_ε , do some integrations by parts using also the equations of motion of the rigid body, we get the following energy estimate:

$$(6) \quad \|u_\varepsilon(t)\|_{L^2(\mathbb{R}^2 \setminus \Omega_\varepsilon)}^2 + m_\varepsilon |h_\varepsilon'(t)|^2 + J_\varepsilon |\theta_\varepsilon'(t)|^2 + 4\nu \int_0^t \|D(u_\varepsilon)\|_{L^2(\mathbb{R}^2 \setminus \Omega_\varepsilon)}^2 \\ \leq \|u_\varepsilon(0)\|_{L^2(\mathbb{R}^2 \setminus \Omega_\varepsilon)}^2 + m_\varepsilon |h_\varepsilon'(0)|^2 + J_\varepsilon |\theta_\varepsilon'(0)|^2.$$

To solve the system of equations (1)–(5), we need to impose the initial data. For the fluid part of the system we need to impose the initial velocity $u_\varepsilon(0, x)$. The two equations describing the evolution of the rigid body are second-order in time, so we need to know $h_\varepsilon(0)$, $h_\varepsilon'(0)$, $\theta_\varepsilon(0)$ and $\theta_\varepsilon'(0)$. The system of equations being translation invariant, we can assume without loss of generality that the initial position of the center of mass of the rigid body is in the origin: $h_\varepsilon(0) = 0$. Moreover, from the definition of the angle of rotation θ_ε we obviously have that $\theta_\varepsilon(0) = 0$. So we only need to impose $u_\varepsilon(0, x)$, $h_\varepsilon'(0)$ and $\theta_\varepsilon'(0)$. The initial velocity will be assumed to be square integrable only. As such, its trace on the boundary is not well-defined. Only its normal trace is defined thanks to the divergence free condition. Therefore, we need to impose the following compatibility condition on the initial velocity:

$$(7) \quad u_\varepsilon(0, x) \cdot n_\varepsilon = [h_\varepsilon'(0) + \theta_\varepsilon'(0)(x - h_\varepsilon(0))^\perp] \cdot n_\varepsilon \quad \text{on } \partial\Omega_\varepsilon(0).$$

In conclusion, to solve the system of equations (1)–(5), we need to impose that $u_\varepsilon(0, x) \in L^2(\mathbb{R}^2 \setminus \Omega_\varepsilon(0))$, that $\text{div } u_\varepsilon(0, x) = 0$ in $\mathbb{R}^2 \setminus \Omega_\varepsilon(0)$ and the compatibility condition (7). There is no condition required on $h_\varepsilon'(0)$ and $\theta_\varepsilon'(0)$ while $h_\varepsilon(0) = 0$ and $\theta_\varepsilon(0) = 0$.

To state the classical result of existence and uniqueness of solutions of (1)–(5), it is practical to extend the velocity field u_ε inside the rigid body as follows:

$$(8) \quad \tilde{u}_\varepsilon(t, x) = \begin{cases} u_\varepsilon(t, x) & \text{if } x \in \mathbb{R}^2 \setminus \Omega_\varepsilon(t) \\ h_\varepsilon'(t) + \theta_\varepsilon'(t)(x - h_\varepsilon(t))^\perp & \text{if } x \in \Omega_\varepsilon(t). \end{cases}$$

The conditions imposed on the initial data ensure that $\tilde{u}_\varepsilon(0, x)$ belongs to $L^2(\mathbb{R}^2)$ and is divergence free in \mathbb{R}^2 .

Let us denote by ρ_ε the density of the rigid body Ω_ε . We extend ρ_ε in the fluid region $\mathbb{R}^2 \setminus \Omega_\varepsilon$ by giving it value 1:

$$\tilde{\rho}_\varepsilon(t, x) = \begin{cases} 1, & x \in \mathbb{R}^2 \setminus \Omega_\varepsilon(t) \\ \rho_\varepsilon, & x \in \Omega_\varepsilon(t). \end{cases}$$

Due to the energy estimates (6), global existence of finite energy solutions of (1)–(5) have been proved in a variety of settings. The literature is vast, we give here just a few references dealing with the dimension two: in [2], [7] and [13] the authors consider the case of one or several rigid bodies moving in a bounded domain filled with a viscous fluid while in [15] the authors consider a single disk moving in a fluid filling the whole plane. The existence for the problem we are considering here was not explicitly studied in these works (because we do not assume the rigid body to be a disk), but more complicated cases have been considered in the literature: the case of a 2D bounded domain where collisions with the boundary must be taken into account (see [2], [7] and [13]) and the case of \mathbb{R}^3 with a rigid body of arbitrary shape (see for example [16] and [14]). From these results we can extract the following statement about the existence and uniqueness of solutions of (1)–(5). We use the notation $\mathbb{R}_+ = [0, \infty)$ and emphasize that the endpoint 0 belongs to \mathbb{R}_+ . This is important when we write local spaces in \mathbb{R}_+ like for instance $L^2_{loc}(\mathbb{R}_+) = \{f ; f \text{ square integrable on any interval } [0, t]\}$. We will give a formulation of the PDE in terms of the extended velocity \tilde{u}_ε .

Theorem 1. *Let $u_\varepsilon(0, x) \in L^2(\mathbb{R}^2 \setminus \Omega_\varepsilon(0))$ be divergence free and verifying the compatibility condition (7). We assume that $h_\varepsilon(0) = 0$ and $\theta_\varepsilon(0) = 0$ and we extend $u_\varepsilon(0, x)$ to $\tilde{u}_\varepsilon(0, x)$ as in (8). Then $\tilde{u}_\varepsilon(0, x)$ is divergence free and square integrable on \mathbb{R}^2 and there exists a unique global weak solution $(u_\varepsilon, h_\varepsilon, \theta_\varepsilon)$ of (1)–(5) in the following sense:*

- $u_\varepsilon, h_\varepsilon, \theta_\varepsilon$ satisfy

$$u_\varepsilon \in L^\infty(\mathbb{R}_+; L^2(\mathbb{R}^2 \setminus \Omega_\varepsilon)) \cap L^2_{loc}(\mathbb{R}_+; H^1(\mathbb{R}^2 \setminus \Omega_\varepsilon)),$$

$$h_\varepsilon \in W^{1,\infty}(\mathbb{R}_+; \mathbb{R}^2), \quad \theta_\varepsilon \in W^{1,\infty}(\mathbb{R}_+; \mathbb{R});$$

- if we define \tilde{u}_ε as in (8) then \tilde{u}_ε is divergence free with $D\tilde{u}_\varepsilon(t, x) = 0$ in $\Omega_\varepsilon(t)$ and the equations of motion are verified in the sense of distributions under the following form

$$- \int_0^\infty \int_{\mathbb{R}^2} \tilde{\rho}_\varepsilon \tilde{u}_\varepsilon \cdot (\partial_t \varphi_\varepsilon + (\tilde{u}_\varepsilon \cdot \nabla) \varphi_\varepsilon) + 2\nu \int_0^\infty \int_{\mathbb{R}^2} D(\tilde{u}_\varepsilon) : D(\varphi_\varepsilon) = \int_{\mathbb{R}^2} \tilde{\rho}_\varepsilon(0) \tilde{u}_\varepsilon(0) \cdot \varphi_\varepsilon(0).$$

for any divergence free test function $\varphi_\varepsilon \in H^1(\mathbb{R}_+ \times \mathbb{R}^2)$ compactly supported in time and such that $D\varphi_\varepsilon(t, x) = 0$ in $\Omega_\varepsilon(t)$;

Moreover, \tilde{u}_ε satisfies the following energy inequality:

$$(9) \quad \int_{\mathbb{R}^2} \tilde{\rho}_\varepsilon |\tilde{u}_\varepsilon|^2 + 4\nu \int_0^t \int_{\mathbb{R}^2} |D(\tilde{u}_\varepsilon)|^2 \leq \int_{\mathbb{R}^2} \tilde{\rho}_\varepsilon(0) |\tilde{u}_\varepsilon(0)|^2 \quad \forall t > 0.$$

As mentioned before, we are interested in describing the asymptotic behavior of this fluid-solid system when the diameter of the rigid body Ω_ε goes to 0. There are several papers dealing with this issue when the rigid body does not move with the fluid. Iftimie, Lopes Filho and Nussenzveig Lopes [9] have treated the asymptotic behavior of viscous incompressible 2D flow in the exterior of a small fixed rigid body as the size of the rigid body becomes very small, see also [1] for the case of the periodic boundary conditions. Moreover, Lacave [10] considered a two-dimensional viscous fluid in the exterior of a thin fixed rigid body shrinking to a curve and proved convergence to a solution of the Navier-Stokes equations in the exterior of a curve.

Although we are dealing here only with viscous fluids, let us mention that the case of a perfect incompressible fluid governed by the Euler equations also makes sense and the literature is richer. Let us mention a few results. Iftimie, Lopes Filho and Nussenzveig Lopes [8] have studied the asymptotic behavior of incompressible, ideal two-dimensional

flow in the exterior of a small fixed rigid body when the size of the rigid body becomes very small. Recently, Glass, Lacave and Sueur [4] have studied the case when the solid body shrinks to a point with fixed mass and circulation and is moving with the fluid. The same three authors also consider in [5] the case when the body shrinks to a massless pointwise particle with fixed circulation. In that case, the fluid-solid system converges to the vortex-wave system. In addition, Glass, Munnier and Sueur [6] considered the case of a bounded domain.

As far as we know, there is only one result dealing with the case of a small rigid body moving in a viscous fluid in dimension two. More precisely, Lacave and Takahashi [11] considered a small moving disk in a two-dimensional viscous incompressible fluid. They used a fixed-point type argument based on previously known $L^p - L^q$ decay estimates of the linear semigroup associated to the fluid-solid system (see [3]). They proved convergence towards the solution of the Navier-Stokes equations in \mathbb{R}^2 under the assumption that the rigid body is a disk of radius ε , that the density ρ_ε is constant plus some smallness assumptions on the initial data (including the smallness of the L^2 norm of the initial fluid velocity). More precisely, their result is the following.

Theorem 2 ([11]). *There exists a constant $\lambda_0 > 0$ such that if*

- $u_\varepsilon(0, x) \in L^2(\mathbb{R}^2 \setminus \Omega_\varepsilon(0))$ is divergence free and verifies the compatibility condition (7);
- the rigid body is the disk $\Omega_\varepsilon = D(h_\varepsilon, \varepsilon)$;
- the density ρ_ε is assumed to be independent of ε ;
- $\tilde{u}_\varepsilon(0, x)$ converges weakly in $L^2(\mathbb{R}^2)$ to some $u_0(x)$;
- we have the following smallness of the initial data

$$(10) \quad \|u_\varepsilon(0, x)\|_{L^2(\mathbb{R}^2 \setminus \Omega_\varepsilon(0))} + \varepsilon|h'_\varepsilon(0)| + \varepsilon^2|\theta'_\varepsilon(0)| \leq \lambda_0$$

then the global solution \tilde{u}_ε given by Theorem 1 converges weak* in $L^\infty(\mathbb{R}_+; L^2(\mathbb{R}^2)) \cap L^2_{loc}(\mathbb{R}_+; H^1(\mathbb{R}^2))$ towards the weak solution of the Navier-Stokes equations in \mathbb{R}^2 with initial data u_0 .

Although they state their result for constant density, presumably the proof can be adapted to the case where $\rho_\varepsilon \geq \rho_0$ for some $\rho_0 > 0$ independent of ε . On the other hand, the hypothesis that Ω_ε is a disk seems to be essential in the result of [11]. Indeed, a key ingredient are the estimates of [3] and the proof of that result relies heavily on the fact that Ω_ε is a disk because it uses explicit formulae valid only for the case of a disk. Moreover, it is also hard to see how the smallness condition (10) could be removed in their argument. Indeed, they use a fixed point argument and that requires smallness at some point. Let us observe that in [11] the authors also obtain uniform bounds in ε for the velocity of the disk. Therefore, they can prove that the center of mass of the disk converges to some trajectory. However, nothing can be said about this limit trajectory.

Here, we improve the result of [11] in two respects. First, the rigid body does not need to be a disk. It does not even need to be shrinking homothetically to a point like in [11]. We only assume that the diameter of the rigid body goes to 0. Second, we require no smallness assumption on the initial fluid velocity $u_\varepsilon(0, x)$. On the other hand, we need to assume that the density of the rigid body goes to infinity and we are not able to prove uniform bounds on the motion of the rigid body as in [11]. More precisely, we will prove the following result.

Theorem 3. *We assume the hypothesis of Theorem 1 and moreover*

- $\Omega_\varepsilon(0) \subset D(0, \varepsilon)$;

- the mass m_ε of the rigid body verifies that

$$(11) \quad \frac{m_\varepsilon}{\varepsilon^2} \rightarrow \infty \quad \text{as } \varepsilon \rightarrow 0;$$

- $u_\varepsilon(0, x)$ is bounded independently of ε in $L^2(\mathbb{R}^2 \setminus \Omega_\varepsilon(0))$ and $\sqrt{m_\varepsilon}h'_\varepsilon(0)$ and $\sqrt{J_\varepsilon}\theta'_\varepsilon(0)$ are bounded independently of ε ;
- $\tilde{u}_\varepsilon(0, x)$ converges weakly in $L^2(\mathbb{R}^2)$ to some $u_0(x)$ where $\tilde{u}_\varepsilon(0, x)$ is constructed as in (8).

Let $(u_\varepsilon, h_\varepsilon, \theta_\varepsilon)$ be the global solution of the system (1)–(5) given by Theorem 1. Then \tilde{u}_ε converges weak* in $L^\infty(\mathbb{R}_+; L^2(\mathbb{R}^2)) \cap L^2_{loc}(\mathbb{R}_+; H^1(\mathbb{R}^2))$ as $\varepsilon \rightarrow 0$ towards the solution of the Navier-Stokes equations in \mathbb{R}^2 with initial data u_0 .

It will be clear from the proof that the convergence of \tilde{u}_ε is stronger than stated. For instance, we shall prove that \tilde{u}_ε converges strongly in L^2_{loc} (see Section 4).

Let us remark that if the measure of Ω_ε is of order ε^2 (something which is true if the rigid body shrinks homothetically to a point, *i.e.* if $\Omega_\varepsilon(0)$ is ε times a fixed rigid body) then the hypothesis (11) means that the density ρ_ε of the rigid body goes to ∞ as $\varepsilon \rightarrow 0$.

Observe next that the boundedness of $u_\varepsilon(0, x)$ in $L^2(\mathbb{R}^2 \setminus \Omega_\varepsilon(0))$ and the boundedness of $\sqrt{m_\varepsilon}h'_\varepsilon(0)$ and $\sqrt{J_\varepsilon}\theta'_\varepsilon(0)$ imply the boundedness of $\sqrt{\tilde{\rho}_\varepsilon(0, x)}\tilde{u}_\varepsilon(0, x)$ in $L^2(\mathbb{R}^2)$. Since $\rho_\varepsilon \rightarrow \infty$ this implies that $\tilde{u}_\varepsilon(0, x)$ is bounded in $L^2(\mathbb{R}^2)$. Therefore, the weak convergence of $\tilde{u}_\varepsilon(0, x)$ to $u_0(x)$ is not really a new hypothesis.

Moreover, the boundedness of $\sqrt{\tilde{\rho}_\varepsilon(0, x)}\tilde{u}_\varepsilon(0, x)$ in $L^2(\mathbb{R}^2)$ and the energy inequality (9) imply that $\sqrt{\tilde{\rho}_\varepsilon}\tilde{u}_\varepsilon$ is bounded independently of ε in the space $L^\infty(\mathbb{R}_+; L^2(\mathbb{R}^2)) \cap L^2_{loc}(\mathbb{R}_+; H^1(\mathbb{R}^2))$. Using again that $\rho_\varepsilon \rightarrow \infty$ we deduce that \tilde{u}_ε is also bounded independently of ε in $L^\infty(\mathbb{R}_+; L^2(\mathbb{R}^2)) \cap L^2_{loc}(\mathbb{R}_+; H^1(\mathbb{R}^2))$. And this is all we need to prove the convergence of \tilde{u}_ε towards a solution of the Navier-Stokes equations in \mathbb{R}^2 . Our proof does not require that \tilde{u}_ε verifies the boundary conditions on Ω_ε , nor do we need that $D\tilde{u}_\varepsilon = 0$ in Ω_ε . We only need the above mentioned boundedness of \tilde{u}_ε and the fact that it verifies the Navier-Stokes equations (without any boundary condition) in the exterior of the disk $D(h_\varepsilon(t), \varepsilon)$. We state next a more general result.

Theorem 4. *Let v_ε be a time-dependent divergence free vector field defined on $\mathbb{R}_+ \times \mathbb{R}^2$ belonging to the space*

$$(12) \quad L^\infty(\mathbb{R}_+; L^2(\mathbb{R}^2)) \cap L^2_{loc}(\mathbb{R}_+; H^1(\mathbb{R}^2)) \cap C^0_w(\mathbb{R}_+; L^2_{loc}(\mathbb{R}^2 \setminus \overline{D}(h_\varepsilon(t), \varepsilon)))$$

and let $h_\varepsilon \in W^{1,\infty}(\mathbb{R}_+; \mathbb{R}^2)$. Assume moreover that

- v_ε is bounded independently of ε in the above space;
- $v_\varepsilon(0, x)$ converges weakly in L^2 as $\varepsilon \rightarrow 0$ to some $v_0(x)$;
- v_ε verifies the Navier-Stokes equations in the exterior of the disk $\overline{D}(h_\varepsilon(t), \varepsilon)$:

$$(13) \quad \partial_t v_\varepsilon - \nu \Delta v_\varepsilon + v_\varepsilon \cdot \nabla v_\varepsilon = -\nabla \pi_\varepsilon \quad \text{in the set } \{(t, x) ; t > 0 \text{ and } |x - h_\varepsilon(t)| > \varepsilon\}$$

for some π_ε ;

- the velocity of the center of the disk verifies that $\varepsilon|h'_\varepsilon(t)| \rightarrow 0$ in $L^\infty_{loc}(\mathbb{R}_+)$ when $\varepsilon \rightarrow 0$.

Let v be the unique solution of the Navier-Stokes equations in \mathbb{R}^2 with initial data v_0 . Then v_ε converges to v as $\varepsilon \rightarrow 0$ weak* in the space $L^\infty(\mathbb{R}_+; L^2(\mathbb{R}^2)) \cap L^2_{loc}(\mathbb{R}_+; H^1(\mathbb{R}^2))$.

Theorem 4 with $v_\varepsilon = \tilde{u}_\varepsilon$ implies Theorem 3. Indeed, we already observed above that \tilde{u}_ε has all the properties required from v_ε in Theorem 4. And the hypothesis made on the mass of the rigid body, see relation (11), in Theorem 3 implies that $\varepsilon|h'_\varepsilon(t)| \rightarrow 0$ in

$L_{\text{loc}}^\infty(\mathbb{R}_+)$ when $\varepsilon \rightarrow 0$. This can be easily seen from the energy estimate (6). Indeed, the hypothesis of Theorem 3 implies that the right-hand side of (6) is bounded uniformly in ε so $\sqrt{m_\varepsilon}h'_\varepsilon$ is uniformly bounded in t and ε . The fact that $\frac{m_\varepsilon}{\varepsilon^2} \rightarrow \infty$ and the boundedness of $\sqrt{m_\varepsilon}h'_\varepsilon$ implies that $\varepsilon h'_\varepsilon \rightarrow 0$ as $\varepsilon \rightarrow 0$ uniformly in time.

The idea of the proof of Theorem 4 is completely different from the proof given in [11]. We multiply (13) with a cut-off vanishing on the disk $D(h_\varepsilon(t), \varepsilon)$ constructed in a very particular manner. We then pass to the limit with classical compactness methods. The difficulty here is that the cut-off function itself depends on the time, so time-derivative estimates of v_ε are not so easy to obtain. Also, passing to the limit in the terms $\partial_t v$ and Δv is not obvious: the first is difficult because the time derivative is hard to control and the second one is difficult because the cut-off introduces negative powers of ε in this term.

The plan of the paper is the following. In the following section we introduce some notation and prove some preliminary results. In Section 3 we construct the special cut-off near the rigid body. The required temporal estimates are proved in Section 4. Finally, we pass to the limit in Section 5.

2. NOTATION AND PRELIMINARY RESULTS

We use the classical notation C^m for functions with m continuous derivatives and H^m the Sobolev space of functions with m square-integrable weak derivatives. The notation C_b^m stands for functions in C^m with bounded derivatives up to order m . All function spaces and norms are considered to be taken on \mathbb{R}^2 in the x variable unless otherwise specified. We define $C_{0,\sigma}^\infty$ to be the space of smooth, compactly supported and divergence free vector fields on \mathbb{R}^2 . The derivatives are always taken with respect to the variable x unless otherwise specified. The double dot product of two matrices $M = (m_{ij})$ and $N = (n_{ij})$ denotes the quantity $M : N = \sum_{i,j} m_{ij}n_{ij}$. We denote by C a generic universal constant whose value can change from one line to another.

Let $\varphi \in C_b^1(\mathbb{R}_+; C_{0,\sigma}^\infty)$. We define the stream function ψ of φ by

$$\psi(x) = \int_{\mathbb{R}^2} \frac{(x-y)^\perp}{2\pi|x-y|^2} \cdot \varphi(y) dy.$$

It is well-known that $\psi \in C_b^1(\mathbb{R}_+; C^\infty)$ and $\nabla^\perp \psi = \varphi$. The stream function ψ given above is characterized by two facts. One is that $\nabla^\perp \psi = \varphi$ and another one is that it vanishes at infinity. But in our case, the vanishing at infinity is not important since we will use compactly supported test functions. On the other hand, it is useful to have the stream function small in the neighborhood of the rigid body. We define now a modified stream function, denoted by ψ_ε , which vanishes at the center of the disk $D(h_\varepsilon(t), \varepsilon)$:

$$(14) \quad \psi_\varepsilon(t, x) = \psi(t, x) - \psi(t, h_\varepsilon(t)).$$

Observe that even if φ is constant in time, the modified stream function still depends on the time through h_ε . We collect some properties of the modified stream function in the following lemma.

Lemma 1. *The modified stream function ψ_ε has the following properties:*

- (i) *We have that $\psi_\varepsilon \in W^{1,\infty}(\mathbb{R}_+; C^\infty)$ and $\nabla^\perp \psi_\varepsilon = \varphi$.*
- (ii) *For all $t, R \geq 0$ and $x \in \mathbb{R}^2$ we have that*

$$(15) \quad \|\psi_\varepsilon(t, \cdot)\|_{L^\infty(D(h_\varepsilon(t), R))} \leq R \|\varphi(t, \cdot)\|_{L^\infty}$$

and

$$(16) \quad \|\partial_t \psi_\varepsilon(t, \cdot)\|_{L^\infty(D(h_\varepsilon(t), R))} \leq R \|\partial_t \varphi(t, \cdot)\|_{L^\infty} + |h'_\varepsilon(t)| \|\varphi(t, \cdot)\|_{L^\infty}$$

with the remark that the last relation holds true only almost everywhere in time.

Proof. Clearly $\nabla^\perp \psi_\varepsilon = \nabla^\perp \psi = \varphi$. Since $h_\varepsilon \in W^{1,\infty}(\mathbb{R}_+)$ and $\psi \in C_b^1(\mathbb{R}_+; C^\infty)$ we immediately see that $\psi_\varepsilon \in W^{1,\infty}(\mathbb{R}_+; C^\infty)$ which proves (i).

By the mean value theorem

$$(17) \quad |\psi_\varepsilon(t, x)| = |\psi(t, x) - \psi(t, h_\varepsilon(t))| \leq |x - h_\varepsilon(t)| \|\nabla \psi(t, \cdot)\|_{L^\infty} = |x - h_\varepsilon(t)| \|\varphi(t, \cdot)\|_{L^\infty}.$$

Relation (15) follows. To prove (16) we recall that h_ε is Lipschitz in time so it is almost everywhere differentiable in time. Let t be a time where h_ε is differentiable. We write

$$\begin{aligned} \partial_t \psi_\varepsilon(t, x) &= \partial_t (\psi(t, x) - \psi(t, h_\varepsilon(t))) \\ &= \partial_t \psi(t, x) - \partial_t \psi(t, h_\varepsilon(t)) - h'_\varepsilon(t) \cdot \nabla \psi(t, h_\varepsilon(t)) \end{aligned}$$

so

$$\begin{aligned} \|\partial_t \psi_\varepsilon(t, \cdot)\|_{L^\infty(D(h_\varepsilon(t), R))} &\leq \|\partial_t \psi(t, x) - \partial_t \psi(t, h_\varepsilon(t))\|_{L^\infty(D(h_\varepsilon(t), R))} + |h'_\varepsilon(t)| \|\nabla \psi(t, \cdot)\|_{L^\infty} \\ &\leq R \|\partial_t \nabla \psi(t, \cdot)\|_{L^\infty} + |h'_\varepsilon(t)| \|\varphi(t, \cdot)\|_{L^\infty} \\ &= R \|\partial_t \varphi(t, \cdot)\|_{L^\infty} + |h'_\varepsilon(t)| \|\varphi(t, \cdot)\|_{L^\infty}. \end{aligned}$$

This completes the proof of the lemma. \square

We will need to define a cut-off function near the rigid body with L^2 norm of the gradient as small as possible. This will be done in the next section. For the moment, let us recall that the function that minimizes the L^2 norm of the gradient, that vanishes for $|x| = A$ and is equal to 1 for $|x| = B$ is harmonic. So it is given by the explicit formula

$$f_{A,B} : \mathbb{R}^2 \rightarrow [0, 1], \quad f_{A,B}(x) = \begin{cases} 0 & \text{if } |x| < A \\ \frac{\ln|x| - \ln A}{\ln B - \ln A} & \text{if } A < |x| < B \\ 1 & \text{if } |x| > B. \end{cases}$$

This special cut-off has the following properties.

Lemma 2. *We have that $f_{A,B} \in W^{1,\infty}$. Moreover,*

$$\begin{aligned} \|f_{A,B}(x) - 1\|_{L^2}^2 &= \pi A^2 \left(\frac{\alpha^2}{2 \ln^2 \alpha} - \frac{1}{2 \ln^2 \alpha} - \frac{1}{\ln \alpha} \right), \\ \|\nabla f_{A,B}\|_{L^2}^2 &= \frac{2\pi}{\ln \alpha} \end{aligned}$$

and

$$\| |x| \nabla^2 f_{A,B} \|_{L^2(A < |x| < B)}^2 = \frac{4\pi}{\ln \alpha}$$

where $\alpha = \frac{B}{A}$.

Proof. The Lipschitz character of $f_{A,B}$ is obvious once we remark that $f_{A,B}$ is smooth for $|x| \neq A$ and $|x| \neq B$ and continuous across $|x| = A$ and $|x| = B$.

Next, we have that

$$\begin{aligned}
\|f_{A,B}(x) - 1\|_{L^2}^2 &= \int_{|x|<A} 1 \, dx + \int_{A<|x|<B} \left| \frac{\ln|x| - \ln B}{\ln B - \ln A} \right|^2 dx \\
&= \pi A^2 + \frac{B^2}{(\ln B - \ln A)^2} \int_{A/B<|y|<1} \ln^2|y| \, dy \\
&= \pi A^2 \left(1 + \frac{\alpha^2}{\ln^2 \alpha} \int_{1/\alpha}^1 \ln^2 r \, 2r \, dr \right) \\
&= \pi A^2 \left(\frac{\alpha^2}{2 \ln^2 \alpha} - \frac{1}{2 \ln^2 \alpha} - \frac{1}{\ln \alpha} \right).
\end{aligned}$$

From the definition of $f_{A,B}$, we compute for $A < |x| < B$

$$\nabla f_{A,B} = \frac{x}{|x|^2 \ln \alpha} \quad \text{and} \quad |\nabla^2 f_{A,B}| = \frac{\sqrt{2}}{|x|^2 \ln \alpha}.$$

So

$$\|\nabla f_{A,B}\|_{L^2}^2 = \frac{1}{\ln^2 \alpha} \int_{A<|x|<B} \frac{1}{|x|^2} \, dx = \frac{2\pi}{\ln \alpha}$$

and

$$\| |x| \nabla^2 f_{A,B} \|_{L^2(A<|x|<B)}^2 = \frac{2}{\ln^2 \alpha} \int_{A<|x|<B} \frac{1}{|x|^2} \, dx = \frac{4\pi}{\ln \alpha}.$$

This completes the proof of the lemma. \square

3. CUT-OFF NEAR THE RIGID BODY

We begin now the proof of Theorem 4. It suffices to prove the following statement.

Proposition 1. *For all finite times $T > 0$ there exists a subsequence v_{ε_k} which converges weak* in $L^\infty(0, T; L^2) \cap L^2(0, T; H^1)$ towards a solution $v \in L^\infty(0, T; L^2) \cap L^2(0, T; H^1)$ of the Navier-Stokes equations on $[0, T) \times \mathbb{R}^2$ with initial data v_0 .*

Indeed, let us assume that Proposition 1 is proved. We know that the Navier-Stokes equations in dimension two have a unique global solution v in the space $L^\infty(\mathbb{R}_+; L^2) \cap L^2(\mathbb{R}_+; H^1)$, see for example [12]. The solution v from Proposition 1 is necessarily the restriction to $[0, T]$ of this unique global solution. Since we have uniqueness of the limit, we deduce that the whole sequence v_ε converges weak* in $L^\infty(0, T; L^2) \cap L^2(0, T; H^1)$ towards v . Since T is arbitrary, Theorem 4 follows.

The rest of this paper is devoted to the proof of Proposition 1. Let $T > 0$ be fixed. From now on the time t is assumed to belong to the interval $[0, T]$. The constant K will denote a constant which depends only on ν and

$$\sup_{0 < \varepsilon \leq 1} \|v_\varepsilon\|_{L^\infty(0, T; L^2) \cap L^2(0, T; H^1)}$$

and whose value may change from one line to another. In particular, the constant K does not depend on ε .

By hypothesis we know that

$$(18) \quad \limsup_{\varepsilon \rightarrow 0} \varepsilon |h'_\varepsilon(t)| = 0.$$

We assume that $\varepsilon \leq 1/100$ and we choose α_ε such that

$$(19) \quad 100 \leq \alpha_\varepsilon \leq \frac{1}{\varepsilon}, \quad \lim_{\varepsilon \rightarrow 0} \alpha_\varepsilon = \infty \quad \text{and} \quad \lim_{\varepsilon \rightarrow 0} \varepsilon \alpha_\varepsilon (1 + |h'_\varepsilon(t)|) = 0$$

uniformly in $t \in [0, T]$. The existence of such an α_ε follows from (18). Indeed, we could choose for instance

$$\alpha_\varepsilon = \max\left(100, \frac{1}{\sup_{[0, T]} \sqrt{\varepsilon + \varepsilon |h'_\varepsilon(t)|}}\right).$$

We construct in the following lemma a special cut-off function f_ε near the disk $D(h_\varepsilon(t), \varepsilon)$ such that $f_\varepsilon(x) = 0$ for all $|x| \leq \varepsilon$ and $f_\varepsilon(x) = 1$ for all $|x| \geq \varepsilon \alpha_\varepsilon$.

Lemma 3. *There exists a smooth cut-off function $f_\varepsilon \in C^\infty(\mathbb{R}^2; [0, 1])$ such that*

- (i) f_ε vanishes in the neighborhood of the disk $\overline{D}(0, \varepsilon)$ and $f_\varepsilon = 1$ for $|x| \geq \varepsilon \alpha_\varepsilon$;
- (ii) there exists a universal constant C such that

$$\|f_\varepsilon\|_{L^\infty} = 1, \quad \|\nabla f_\varepsilon\|_{L^2} \leq \frac{C}{\sqrt{\ln \alpha_\varepsilon}}, \quad \| |x| \nabla^2 f_\varepsilon \|_{L^2} \leq \frac{C}{\sqrt{\ln \alpha_\varepsilon}} \quad \text{and} \quad \|f_\varepsilon - 1\|_{L^2} \leq C \frac{\varepsilon \alpha_\varepsilon}{\ln \alpha_\varepsilon}.$$

Proof. From Lemma 2 we observe that the function

$$\tilde{f}_\varepsilon = f_{\varepsilon, \varepsilon \alpha_\varepsilon} = \begin{cases} 0 & \text{if } |x| < \varepsilon \\ \frac{\ln(|x|/\varepsilon)}{\ln \alpha_\varepsilon} & \text{if } \varepsilon < |x| < \varepsilon \alpha_\varepsilon \\ 1 & \text{if } |x| > \varepsilon \alpha_\varepsilon \end{cases}$$

satisfies

$$\|\tilde{f}_\varepsilon\|_{L^\infty} = 1, \quad \|\nabla \tilde{f}_\varepsilon\|_{L^2} \leq \frac{C}{\sqrt{\ln \alpha_\varepsilon}}, \quad \| |x| \nabla^2 \tilde{f}_\varepsilon \|_{L^2(\varepsilon < |x| < \varepsilon \alpha_\varepsilon)} \leq \frac{C}{\sqrt{\ln \alpha_\varepsilon}}$$

and

$$\|\tilde{f}_\varepsilon - 1\|_{L^2} \leq C \frac{\varepsilon \alpha_\varepsilon}{\ln \alpha_\varepsilon}$$

so it has all the required properties except smoothness. More precisely, \tilde{f}_ε is not smooth across $|x| = \varepsilon$ and $|x| = \varepsilon \alpha_\varepsilon$. To obtain a smooth function f_ε from \tilde{f}_ε we need to cut-off in the neighborhood of these two circles.

Let $g \in C_0^\infty(\mathbb{R}^2; [0, 1])$ be such that $g(x) = 0$ for $|x| < 2$ and $g(x) = 1$ for $|x| > 4$. We define

$$g_\varepsilon^1(x) = g\left(\frac{x}{\varepsilon}\right) = \begin{cases} 0, & |x| < 2\varepsilon \\ 1, & |x| > 4\varepsilon \end{cases}$$

and

$$g_\varepsilon^2(x) = 1 - g\left(\frac{8x}{\varepsilon \alpha_\varepsilon}\right) = \begin{cases} 1, & |x| < \frac{\varepsilon \alpha_\varepsilon}{4} \\ 0, & |x| > \frac{\varepsilon \alpha_\varepsilon}{2}. \end{cases}$$

With the help of all the auxiliary functions above, we define a new function

$$f_\varepsilon = 1 + g_\varepsilon^2(g_\varepsilon^1 \tilde{f}_\varepsilon - 1) = \begin{cases} 1, & |x| > \frac{\varepsilon \alpha_\varepsilon}{2} \\ 1 + g_\varepsilon^2(\tilde{f}_\varepsilon - 1), & \frac{\varepsilon \alpha_\varepsilon}{4} < |x| < \frac{\varepsilon \alpha_\varepsilon}{2} \\ \tilde{f}_\varepsilon, & 4\varepsilon < |x| < \frac{\varepsilon \alpha_\varepsilon}{4} \\ g_\varepsilon^1 \tilde{f}_\varepsilon, & 2\varepsilon < |x| < 4\varepsilon \\ 0, & |x| < 2\varepsilon. \end{cases}$$

Clearly f_ε satisfies (i) and is smooth across $|x| = \varepsilon$ and $|x| = \varepsilon\alpha_\varepsilon$, so it remains to prove (ii). From the definition of f_ε , we immediately see that $\|f_\varepsilon\|_{L^\infty} = 1$. To simplify the write-up, we use the notation $L^p(a, b) = L^p(a < |x| < b)$. Clearly g_ε^1 and g_ε^2 are uniformly bounded in L^∞ and ∇g_ε^1 and ∇g_ε^2 are uniformly bounded in L^2 . Using these observations we estimate

$$\begin{aligned} \|\nabla f_\varepsilon\|_{L^2} &\leq \|\nabla(g_\varepsilon^1 \tilde{f}_\varepsilon)\|_{L^2(2\varepsilon, 4\varepsilon)} + \|\nabla \tilde{f}_\varepsilon\|_{L^2(4\varepsilon, \frac{\varepsilon\alpha_\varepsilon}{4})} + \|\nabla(g_\varepsilon^2(\tilde{f}_\varepsilon - 1))\|_{L^2(\frac{\varepsilon\alpha_\varepsilon}{4}, \frac{\varepsilon\alpha_\varepsilon}{2})} \\ &\leq C(\|\nabla \tilde{f}_\varepsilon\|_{L^2} + \|\tilde{f}_\varepsilon\|_{L^\infty(2\varepsilon, 4\varepsilon)} + \|\tilde{f}_\varepsilon - 1\|_{L^\infty(\frac{\varepsilon\alpha_\varepsilon}{4}, \frac{\varepsilon\alpha_\varepsilon}{2})}) \\ &\leq \frac{C}{\sqrt{\ln \alpha_\varepsilon}} + \frac{C}{\ln \alpha_\varepsilon} \\ &\leq \frac{C}{\sqrt{\ln \alpha_\varepsilon}} \end{aligned}$$

where we used the bounds

$$\|\tilde{f}_\varepsilon\|_{L^\infty(2\varepsilon, 4\varepsilon)} = \left\| \frac{\ln(|x|/\varepsilon)}{\ln \alpha_\varepsilon} \right\|_{L^\infty(2\varepsilon, 4\varepsilon)} \leq \frac{C}{\ln \alpha_\varepsilon}$$

and

$$\|\tilde{f}_\varepsilon - 1\|_{L^\infty(\frac{\varepsilon\alpha_\varepsilon}{4}, \frac{\varepsilon\alpha_\varepsilon}{2})} = \left\| \frac{\ln(|x|/(\varepsilon\alpha_\varepsilon))}{\ln \alpha_\varepsilon} \right\|_{L^\infty(\frac{\varepsilon\alpha_\varepsilon}{4}, \frac{\varepsilon\alpha_\varepsilon}{2})} \leq \frac{C}{\ln \alpha_\varepsilon}.$$

Similarly, using in addition that $\| |x| \nabla g_\varepsilon^i \|_{L^\infty}$ and $\| |x| \nabla^2 g_\varepsilon^i \|_{L^2}$ are bounded independently of ε for $i = 1, 2$, we can estimate

$$\begin{aligned} \| |x| \nabla^2 f_\varepsilon \|_{L^2} &\leq \| |x| \nabla^2 (g_\varepsilon^1 \tilde{f}_\varepsilon) \|_{L^2(2\varepsilon, 4\varepsilon)} + \| |x| \nabla^2 \tilde{f}_\varepsilon \|_{L^2(4\varepsilon, \frac{\varepsilon\alpha_\varepsilon}{4})} \\ &\quad + \| |x| \nabla^2 (g_\varepsilon^2 (\tilde{f}_\varepsilon - 1)) \|_{L^2(\frac{\varepsilon\alpha_\varepsilon}{4}, \frac{\varepsilon\alpha_\varepsilon}{2})} \\ &\leq C(\| |x| \nabla^2 \tilde{f}_\varepsilon \|_{L^2} + \|\nabla \tilde{f}_\varepsilon\|_{L^2} + \|\tilde{f}_\varepsilon\|_{L^\infty(2\varepsilon, 4\varepsilon)} + \|\tilde{f}_\varepsilon - 1\|_{L^\infty(\frac{\varepsilon\alpha_\varepsilon}{4}, \frac{\varepsilon\alpha_\varepsilon}{2})}) \\ &\leq \frac{C}{\sqrt{\ln \alpha_\varepsilon}} + \frac{C}{\ln \alpha_\varepsilon} \\ &\leq \frac{C}{\sqrt{\ln \alpha_\varepsilon}}. \end{aligned}$$

Finally,

$$\begin{aligned} \|f_\varepsilon - 1\|_{L^2} &\leq \|g_\varepsilon^1 \tilde{f}_\varepsilon - 1\|_{L^2(2\varepsilon, 4\varepsilon)} + \|\tilde{f}_\varepsilon - 1\|_{L^2(4\varepsilon, \frac{\varepsilon\alpha_\varepsilon}{4})} + \|g_\varepsilon^2(\tilde{f}_\varepsilon - 1)\|_{L^2(\frac{\varepsilon\alpha_\varepsilon}{4}, \frac{\varepsilon\alpha_\varepsilon}{2})} + \|1\|_{L^2(|x| < 2\varepsilon)} \\ &\leq C(\|\tilde{f}_\varepsilon - 1\|_{L^2} + \|1\|_{L^2(|x| < 4\varepsilon)}) \\ &\leq C\left(\frac{\varepsilon\alpha_\varepsilon}{\ln \alpha_\varepsilon} + \varepsilon\right) \\ &\leq C\frac{\varepsilon\alpha_\varepsilon}{\ln \alpha_\varepsilon}. \end{aligned}$$

This completes the proof of the lemma. \square

The function f_ε is a cut-off in the neighborhood of the disk $D(0, \varepsilon)$. We define now a cut-off in the neighborhood of the disk $D(h_\varepsilon(t), \varepsilon)$ by setting

$$\eta_\varepsilon(t, x) = f_\varepsilon(x - h_\varepsilon(t)).$$

Lemma 3 immediately implies that η_ε has the following properties:

Lemma 4. *We have that*

- (i) $\eta_\varepsilon \in W^{1,\infty}(\mathbb{R}_+; C_{0,\sigma}^\infty)$;
- (ii) η_ε vanishes in the neighborhood of the disk $\overline{D}(h_\varepsilon(t), \varepsilon)$ and $\eta_\varepsilon = 1$ for $|x - h_\varepsilon(t)| \geq \varepsilon\alpha_\varepsilon$;
- (iii) there exists a universal constant C such that

$$(20) \quad \|\eta_\varepsilon\|_{L^\infty} = 1, \quad \|\nabla\eta_\varepsilon\|_{L^2} \leq \frac{C}{\sqrt{\ln \alpha_\varepsilon}}, \quad \||x - h_\varepsilon(t)|\nabla^2\eta_\varepsilon\|_{L^2} \leq \frac{C}{\sqrt{\ln \alpha_\varepsilon}}$$

and

$$(21) \quad \|\eta_\varepsilon - 1\|_{L^2} \leq C \frac{\varepsilon\alpha_\varepsilon}{\ln \alpha_\varepsilon}.$$

Given a test function $\varphi \in C_b^1(\mathbb{R}_+; C_{0,\sigma}^\infty)$ we construct a test function φ_ε on the set $|x - h_\varepsilon(t)| > \varepsilon$ by setting

$$(22) \quad \varphi_\varepsilon = \nabla^\perp(\eta_\varepsilon\psi_\varepsilon)$$

where ψ_ε was defined in Section 2 (see relation (14)). We state some properties of φ_ε in the following lemma:

Lemma 5. *The test function φ_ε has the following properties:*

- (i) $\varphi_\varepsilon \in W^{1,\infty}(\mathbb{R}_+; C_{0,\sigma}^\infty)$ and is supported in the set $|x - h_\varepsilon(t)| > \varepsilon$;
- (ii) $\varphi_\varepsilon \rightarrow \varphi$ strongly in $L^\infty(0, T; H^1)$ as $\varepsilon \rightarrow 0$;
- (iii) there exists a universal constant C such that

$$(23) \quad \|\varphi_\varepsilon\|_{L^\infty(0, T; H^1)} \leq C\|\varphi\|_{L^\infty(0, T; H^3)}.$$

Proof. Since η_ε and ψ_ε are $W^{1,\infty}$ in time and smooth in space, so is φ_ε . The compact support in x of φ_ε in the set $|x - h_\varepsilon(t)| > \varepsilon$ follows from the compact support of φ and the localization properties of η_ε . Obviously φ_ε is also divergence free so claim (i) follows.

Recalling that $\nabla^\perp\psi_\varepsilon = \varphi$ we write

$$\varphi_\varepsilon - \varphi = \nabla^\perp(\eta_\varepsilon\psi_\varepsilon) - \varphi = \nabla^\perp\eta_\varepsilon\psi_\varepsilon + \eta_\varepsilon\nabla^\perp\psi_\varepsilon - \varphi = \nabla^\perp\eta_\varepsilon\psi_\varepsilon + (\eta_\varepsilon - 1)\varphi.$$

Using the bound (15) and recalling that $\nabla\eta_\varepsilon$ is supported in $D(h_\varepsilon(t), \varepsilon\alpha_\varepsilon)$ we can estimate

$$\begin{aligned} \|\varphi_\varepsilon - \varphi\|_{L^2} &\leq \|(\eta_\varepsilon - 1)\varphi\|_{L^2} + \|\nabla\eta_\varepsilon\psi_\varepsilon\|_{L^2} \\ &\leq \|\eta_\varepsilon - 1\|_{L^2}\|\varphi\|_{L^\infty} + \|\nabla\eta_\varepsilon\|_{L^2}\|\psi_\varepsilon\|_{L^\infty(D(h_\varepsilon(t), \varepsilon\alpha_\varepsilon))} \\ &\leq \|\eta_\varepsilon - 1\|_{L^2}\|\varphi\|_{L^\infty} + \varepsilon\alpha_\varepsilon\|\nabla\eta_\varepsilon\|_{L^2}\|\varphi\|_{L^\infty} \\ &= \|\varphi\|_{L^\infty}(\|\eta_\varepsilon - 1\|_{L^2} + \varepsilon\alpha_\varepsilon\|\nabla\eta_\varepsilon\|_{L^2}). \end{aligned}$$

Taking the supremum on $[0, T]$ and using (19), (20) and (21) we deduce that

$$(24) \quad \|\varphi_\varepsilon - \varphi\|_{L^\infty(0, T; L^2)} \leq C \frac{\varepsilon\alpha_\varepsilon}{\sqrt{\ln \alpha_\varepsilon}} \|\varphi\|_{L^\infty([0, T] \times \mathbb{R}^2)} \xrightarrow{\varepsilon \rightarrow 0} 0.$$

Next,

$$\begin{aligned} \|\nabla(\varphi_\varepsilon - \varphi)\|_{L^2} &= \|\nabla\nabla^\perp((\eta_\varepsilon - 1)\psi_\varepsilon)\|_{L^2} \\ &\leq \|\nabla\nabla^\perp\eta_\varepsilon\psi_\varepsilon\|_{L^2} + C\|\nabla\eta_\varepsilon\|_{L^2}\|\nabla\psi_\varepsilon\|_{L^\infty} + \|\eta_\varepsilon - 1\|_{L^2}\|\nabla^2\psi_\varepsilon\|_{L^\infty}. \end{aligned}$$

We bound the first term on the right-hand side using (17) and (20):

$$\|\nabla\nabla^\perp\eta_\varepsilon\psi_\varepsilon\|_{L^2} \leq C\|\varphi\|_{L^\infty}\||x - h_\varepsilon(t)|\nabla^2\eta_\varepsilon\|_{L^2} \leq \frac{C}{\sqrt{\ln \alpha_\varepsilon}}\|\varphi\|_{L^\infty}.$$

Recalling that $\nabla^\perp \psi_\varepsilon = \varphi$ and using again Lemma 4 we infer that

$$\begin{aligned} \|\nabla(\varphi_\varepsilon - \varphi)\|_{L^2} &\leq \frac{C}{\sqrt{\ln \alpha_\varepsilon}} \|\varphi\|_{L^\infty} + C \|\nabla \eta_\varepsilon\|_{L^2} \|\varphi\|_{L^\infty} + \|\eta_\varepsilon - 1\|_{L^2} \|\nabla \varphi\|_{L^\infty} \\ &\leq \frac{C}{\sqrt{\ln \alpha_\varepsilon}} \|\varphi\|_{W^{1,\infty}}. \end{aligned}$$

Combining this bound with (24) implies that

$$\|\varphi_\varepsilon - \varphi\|_{L^\infty(0,T;H^1)} \leq \frac{C}{\sqrt{\ln \alpha_\varepsilon}} \|\varphi\|_{L^\infty(0,T;W^{1,\infty})} \xrightarrow{\varepsilon \rightarrow 0} 0.$$

In addition, we obtain that there exists a universal constant $C > 0$ such that

$$\|\varphi_\varepsilon\|_{L^\infty(0,T;H^1)} \leq C \|\varphi\|_{L^\infty(0,T;H^1 \cap W^{1,\infty})}.$$

Using the Sobolev embedding $H^3 \hookrightarrow W^{1,\infty}$ completes the proof of the lemma. \square

We end this section with an estimate on the H^{-1} norm of the time-derivative of φ_ε .

Lemma 6. *Let w be an H^1 vector field. There exists a universal constant $C > 0$ such that for all times $t \geq 0$ where h_ε is differentiable we have that*

$$\begin{aligned} \left| \int_{\mathbb{R}^2} w(x) \cdot (\partial_t \varphi_\varepsilon(t, x) - \partial_t \varphi(t, x)) dx \right| &\leq C \|\operatorname{curl} w\|_{L^2} \left(\frac{\varepsilon^2 \alpha_\varepsilon^2}{\ln \alpha_\varepsilon} \|\partial_t \varphi(t, \cdot)\|_{L^\infty} \right. \\ &\quad \left. + \frac{\varepsilon \alpha_\varepsilon}{\sqrt{\ln \alpha_\varepsilon}} |h'_\varepsilon(t)| \|\varphi(t, \cdot)\|_{L^\infty} \right). \end{aligned}$$

Proof. Let t be a time where h_ε is differentiable. We use (22) to write

$$\begin{aligned} \int_{\mathbb{R}^2} w(x) \cdot (\partial_t \varphi_\varepsilon(t, x) - \partial_t \varphi(t, x)) dx &= \int_{\mathbb{R}^2} w \cdot \partial_t \nabla^\perp((\eta_\varepsilon - 1)\psi_\varepsilon) \\ &= - \int_{\mathbb{R}^2} \operatorname{curl} w \partial_t((\eta_\varepsilon - 1)\psi_\varepsilon) \\ &= - \int_{\mathbb{R}^2} \operatorname{curl} w \partial_t \eta_\varepsilon \psi_\varepsilon - \int_{\mathbb{R}^2} \operatorname{curl} w (\eta_\varepsilon - 1) \partial_t \psi_\varepsilon. \end{aligned}$$

Clearly

$$\partial_t \eta_\varepsilon = \partial_t (f_\varepsilon(x - h_\varepsilon(t))) = -h'_\varepsilon(t) \cdot \nabla f_\varepsilon(x - h_\varepsilon(t))$$

is supported in the set $\{|x - h_\varepsilon(t)| \leq \varepsilon \alpha_\varepsilon\}$. We can therefore bound

$$\begin{aligned} \left| \int_{\mathbb{R}^2} \operatorname{curl} w \partial_t \eta_\varepsilon \psi_\varepsilon \right| &\leq C |h'_\varepsilon(t)| \int_{|x - h_\varepsilon(t)| \leq \varepsilon \alpha_\varepsilon} |\operatorname{curl} w| |\nabla f_\varepsilon(x - h_\varepsilon(t))| |\psi_\varepsilon| \\ &\leq C |h'_\varepsilon(t)| \|\operatorname{curl} w\|_{L^2} \|\nabla f_\varepsilon\|_{L^2} \|\psi_\varepsilon\|_{L^\infty(D(h_\varepsilon(t), \varepsilon \alpha_\varepsilon))} \\ &\leq C \frac{\varepsilon \alpha_\varepsilon}{\sqrt{\ln \alpha_\varepsilon}} |h'_\varepsilon(t)| \|\operatorname{curl} w\|_{L^2} \|\varphi\|_{L^\infty} \end{aligned}$$

where we used (15) and Lemma 3.

Similarly, $\eta_\varepsilon - 1$ is supported in the set $\{|x - h_\varepsilon(t)| \leq \varepsilon \alpha_\varepsilon\}$ so we can use (16) and (21) to deduce that

$$\begin{aligned} \left| \int_{\mathbb{R}^2} \operatorname{curl} w (\eta_\varepsilon - 1) \partial_t \psi_\varepsilon \right| &\leq \|\operatorname{curl} w\|_{L^2} \|\eta_\varepsilon - 1\|_{L^2} \|\partial_t \psi_\varepsilon\|_{L^\infty(D(h_\varepsilon(t), \varepsilon \alpha_\varepsilon))} \\ &\leq C \frac{\varepsilon \alpha_\varepsilon}{\ln \alpha_\varepsilon} \|\operatorname{curl} w\|_{L^2} (\varepsilon \alpha_\varepsilon \|\partial_t \varphi\|_{L^\infty} + |h'_\varepsilon(t)| \|\varphi\|_{L^\infty}). \end{aligned}$$

The conclusion follows putting together the above relations. \square

4. TEMPORAL ESTIMATE AND STRONG CONVERGENCE

The aim of this section is to prove the strong convergence of some sub-sequence of v_ε . More precisely, we will prove the following result.

Lemma 7. *There exists a sub-sequence v_{ε_k} of v_ε which converges strongly in $L^2(0, T; L^2_{loc})$.*

To prove this lemma we first show some time-derivative estimates and then use the Ascoli theorem.

Let $\varphi \in C_{0,\sigma}^\infty(\mathbb{R}^2)$ be a test function which does not depend on the time. Even though φ does not depend on t , we can still perform the construction of the cut-off φ_ε as in Section 3 (see relation (22)) and all the results of that section remain valid. Observe that even though φ does not depend on the time, the modified test function φ_ε is time-dependent.

Let us denote by H_σ^s the space of H^s divergence free vector fields on \mathbb{R}^2 . We endow H_σ^s with the H^s norm. The dual space of H_σ^s is H_σ^{-s} . We have that $C_{0,\sigma}^\infty$ is dense in H_σ^s for all $s \in \mathbb{R}$.

Let $t \in [0, T]$ be fixed. We use Lemma 4 and relation (15) to bound

$$\begin{aligned} \left| \int_{\mathbb{R}^2} v_\varepsilon(t, x) \cdot \varphi_\varepsilon(t, x) dx \right| &= \left| \int_{\mathbb{R}^2} v_\varepsilon \cdot \nabla^\perp(\eta_\varepsilon \psi_\varepsilon) dx \right| \\ &= \left| \int_{\mathbb{R}^2} v_\varepsilon \cdot (\nabla^\perp \eta_\varepsilon \psi_\varepsilon + \eta_\varepsilon \varphi) dx \right| \\ &\leq \|v_\varepsilon\|_{L^2} \|\nabla \eta_\varepsilon\|_{L^2} \|\psi_\varepsilon\|_{L^\infty(D(h_\varepsilon(t), \varepsilon \alpha_\varepsilon))} + \|v_\varepsilon\|_{L^2} \|\eta_\varepsilon\|_{L^\infty} \|\varphi\|_{L^2} \\ &\leq C \frac{\varepsilon \alpha_\varepsilon}{\sqrt{\ln \alpha_\varepsilon}} \|v_\varepsilon\|_{L^2} \|\varphi\|_{L^\infty} + \|v_\varepsilon\|_{L^2} \|\varphi\|_{L^2} \\ &\leq K_1 \|\varphi\|_{H^2} \end{aligned}$$

for some constant K_1 independent of ε and t . We used above the Sobolev embedding $H^2 \hookrightarrow L^\infty$, the boundedness of v_ε in $L^\infty(0, T; L^2)$ and relations (19) and (20). We infer that, for fixed t , the map

$$C_{0,\sigma}^\infty \ni \varphi \mapsto \int_{\mathbb{R}^2} v_\varepsilon(t, x) \cdot \varphi_\varepsilon(t, x) dx \in \mathbb{R}$$

is linear and continuous for the H^2 norm. Since the closure of $C_{0,\sigma}^\infty$ for the H^2 norm is H_σ^2 , the above map can be uniquely extended to a continuous linear mapping from H_σ^2 to \mathbb{R} . Therefore it can be identified to an element of the dual of H_σ^2 which is H_σ^{-2} . We conclude that there exists some $\Xi_\varepsilon(t) \in H_\sigma^{-2}$ such that

$$\langle \Xi_\varepsilon(t), \varphi \rangle = \int_{\mathbb{R}^2} v_\varepsilon(t, x) \cdot \varphi_\varepsilon(t, x) dx \quad \forall \varphi \in H_\sigma^2.$$

Above $\langle \cdot, \cdot \rangle$ denotes the duality bracket between H_σ^{-2} and H_σ^2 which is the extension of the usual L^2 scalar product. In addition, we have that $\|\Xi_\varepsilon(t)\|_{H^{-2}} \leq K_1$, so Ξ_ε belongs to the space $L^\infty(0, T; H_\sigma^{-2})$ and is bounded independently of ε in this space.

Because φ_ε is compactly supported in $\{|x - h_\varepsilon(t)| > \varepsilon\}$ it can be used as test function in (13). Multiplying (13) by φ_ε and integrating in space and time from s to t yields

$$\int_s^t \int_{\mathbb{R}^2} \partial_\tau v_\varepsilon \cdot \varphi_\varepsilon + \nu \int_s^t \int_{\mathbb{R}^2} \nabla v_\varepsilon : \nabla \varphi_\varepsilon + \int_s^t \int_{\mathbb{R}^2} v_\varepsilon \cdot \nabla v_\varepsilon \cdot \varphi_\varepsilon = 0.$$

We integrate by parts in time the first term above:

$$\begin{aligned} \int_s^t \int_{\mathbb{R}^2} \partial_\tau v_\varepsilon \cdot \varphi_\varepsilon &= \int_{\mathbb{R}^2} v_\varepsilon(t, x) \cdot \varphi_\varepsilon(t, x) dx - \int_{\mathbb{R}^2} v_\varepsilon(s, x) \cdot \varphi_\varepsilon(s, x) dx - \int_s^t \int_{\mathbb{R}^2} v_\varepsilon \cdot \partial_\tau \varphi_\varepsilon \\ &= \langle \Xi_\varepsilon(t) - \Xi_\varepsilon(s), \varphi \rangle - \int_s^t \int_{\mathbb{R}^2} v_\varepsilon \cdot \partial_\tau \varphi_\varepsilon. \end{aligned}$$

We deduce that

$$(25) \quad \langle \Xi_\varepsilon(t) - \Xi_\varepsilon(s), \varphi \rangle = \int_s^t \int_{\mathbb{R}^2} v_\varepsilon \cdot \partial_\tau \varphi_\varepsilon - \nu \int_s^t \int_{\mathbb{R}^2} \nabla v_\varepsilon : \nabla \varphi_\varepsilon - \int_s^t \int_{\mathbb{R}^2} v_\varepsilon \cdot \nabla v_\varepsilon \cdot \varphi_\varepsilon.$$

We bound first

$$\begin{aligned} \left| \nu \int_s^t \int_{\mathbb{R}^2} \nabla v_\varepsilon : \nabla \varphi_\varepsilon \right| &\leq \nu \int_s^t \|\nabla v_\varepsilon\|_{L^2} \|\nabla \varphi_\varepsilon\|_{L^2} \\ &\leq C\nu(t-s)^{\frac{1}{2}} \|\varphi\|_{H^3} \|\nabla v_\varepsilon\|_{L^2([0, T] \times \mathbb{R}^2)} \\ &\leq K(t-s)^{\frac{1}{2}} \|\varphi\|_{H^3} \end{aligned}$$

where we used (23) and the hypothesis that v_ε is bounded in $L^2(0, T; H^1)$.

To estimate the last term in (25) we use the Gagliardo-Nirenberg inequality $\|f\|_{L^4} \leq C\|f\|_{L^2}^{\frac{1}{2}}\|\nabla f\|_{L^2}^{\frac{1}{2}}$, the boundedness of v_ε in the space displayed in (12) and relation (23):

$$\begin{aligned} \left| \int_s^t \int_{\mathbb{R}^2} v_\varepsilon \cdot \nabla v_\varepsilon \cdot \varphi_\varepsilon \right| &= \left| \int_s^t \int_{\mathbb{R}^2} v_\varepsilon \cdot \nabla \varphi_\varepsilon \cdot v_\varepsilon \right| \\ &\leq \int_s^t \|v_\varepsilon\|_{L^4}^2 \|\nabla \varphi_\varepsilon\|_{L^2} \\ &\leq \int_s^t \|v_\varepsilon\|_{L^2} \|\nabla v_\varepsilon\|_{L^2} \|\varphi_\varepsilon\|_{H^1} \\ &\leq C(t-s)^{\frac{1}{2}} \|v_\varepsilon\|_{L^\infty(0, T; L^2)} \|\nabla v_\varepsilon\|_{L^2([0, T] \times \mathbb{R}^2)} \|\varphi\|_{H^3} \\ &\leq K(t-s)^{\frac{1}{2}} \|\varphi\|_{H^3}. \end{aligned}$$

It remains to estimate the first term on the right-hand side of (25). To do that, we use Lemma 6. Recalling that φ does not depend on the time, we can write

$$\begin{aligned} \left| \int_s^t \int_{\mathbb{R}^2} v_\varepsilon \cdot \partial_\tau \varphi_\varepsilon \right| &\leq C \frac{\varepsilon \alpha_\varepsilon}{\sqrt{\ln \alpha_\varepsilon}} \int_s^t \|\operatorname{curl} v_\varepsilon\|_{L^2} |h'_\varepsilon| \|\varphi\|_{L^\infty} \\ &\leq C \frac{\varepsilon \alpha_\varepsilon}{\sqrt{\ln \alpha_\varepsilon}} \|\varphi\|_{H^2} \int_s^t \|\operatorname{curl} v_\varepsilon\|_{L^2} |h'_\varepsilon| \\ &\leq C(t-s)^{\frac{1}{2}} \|\varphi\|_{H^2} \frac{\varepsilon \alpha_\varepsilon}{\sqrt{\ln \alpha_\varepsilon}} \sup_{[0, T]} |h'_\varepsilon| \|\operatorname{curl} v_\varepsilon\|_{L^2([0, T] \times \mathbb{R}^2)}. \end{aligned}$$

Due to the hypothesis imposed on α_ε , see (19), we know that $\frac{\varepsilon \alpha_\varepsilon}{\sqrt{\ln \alpha_\varepsilon}} \sup_{[0, T]} |h'_\varepsilon|$ goes to 0 as $\varepsilon \rightarrow 0$. In particular it is bounded uniformly in ε .

Recalling again the boundedness of v_ε in the space $L^2(0, T; H^1)$, we infer from the above relations that

$$\left| \langle \Xi_\varepsilon(t) - \Xi_\varepsilon(s), \varphi \rangle \right| \leq K(t-s)^{\frac{1}{2}} \|\varphi\|_{H^3}$$

where the constant K does not depend on ε and φ . By density of $C_{0, \sigma}^\infty$ in H_σ^3 we infer that $\|\Xi_\varepsilon(t) - \Xi_\varepsilon(s)\|_{H^{-3}} \leq K(t-s)^{\frac{1}{2}}$. The functions $\Xi_\varepsilon(t)$ are therefore equicontinuous in time with values in H_σ^{-3} . They are also bounded in H_σ^{-3} because we already know

that they are bounded in H_σ^{-2} . Since the embedding $H^{-3} \hookrightarrow H_{\text{loc}}^{-4}$ is compact, the Ascoli theorem implies that there exists a subsequence Ξ_{ε_k} of Ξ_ε which converges strongly in $C^0([0, T]; H_{\text{loc}}^{-4})$.

Recalling the definition of Ξ_ε and using Lemma 4 we can write

$$\begin{aligned} |\langle \Xi_\varepsilon(t) - v_\varepsilon(t), \varphi \rangle| &= \left| \int_{\mathbb{R}^2} v_\varepsilon \cdot (\nabla^\perp \eta_\varepsilon \psi_\varepsilon + \eta_\varepsilon \varphi) dx - \int_{\mathbb{R}^2} v_\varepsilon \cdot \varphi \right| \\ &= \left| \int_{\mathbb{R}^2} v_\varepsilon \cdot (\nabla^\perp \eta_\varepsilon \psi_\varepsilon + (\eta_\varepsilon - 1)\varphi) dx \right| \\ &\leq C \|v_\varepsilon\|_{L^2} \|\nabla \eta_\varepsilon\|_{L^2} \|\psi_\varepsilon\|_{L^\infty(D(h_\varepsilon(t), \varepsilon \alpha_\varepsilon))} + C \|v_\varepsilon\|_{L^2} \|\eta_\varepsilon - 1\|_{L^2} \|\varphi\|_{L^\infty} \\ &\leq C \|v_\varepsilon\|_{L^2} \|\varphi\|_{L^\infty} \left(\frac{\varepsilon \alpha_\varepsilon}{\sqrt{\ln \alpha_\varepsilon}} + \frac{\varepsilon \alpha_\varepsilon}{\ln \alpha_\varepsilon} \right) \\ &\leq C \|v_\varepsilon\|_{L^2} \|\varphi\|_{H^2} \frac{\varepsilon \alpha_\varepsilon}{\sqrt{\ln \alpha_\varepsilon}}. \end{aligned}$$

Hence

$$\|\Xi_\varepsilon(t) - v_\varepsilon(t)\|_{H^{-2}} \leq C \|v_\varepsilon\|_{L^2} \frac{\varepsilon \alpha_\varepsilon}{\sqrt{\ln \alpha_\varepsilon}} \xrightarrow{\varepsilon \rightarrow 0} 0$$

uniformly in time. Recalling that Ξ_{ε_k} converges strongly in H_{loc}^{-4} uniformly in time we infer that v_{ε_k} also converges strongly in $L^\infty(0, T; H_{\text{loc}}^{-4})$. The interpolation inequality $\|\cdot\|_{L^2} \leq \|\cdot\|_{H^{-4}}^{\frac{1}{5}} \|\cdot\|_{H^1}^{\frac{4}{5}}$ and the boundedness of v_ε in $L^2(0, T; H^1)$ finally imply that v_{ε_k} converges strongly in $L^{\frac{5}{2}}(0, T; L_{\text{loc}}^2) \hookrightarrow L^2(0, T; L_{\text{loc}}^2)$. This completes the proof of Lemma 7.

5. PASSING TO THE LIMIT

In this section we complete the proof of Theorem 4. It is now only a matter of putting together the results proved in the previous sections.

Given the boundedness of v_ε in $L^\infty(0, T; L^2) \cap L^2(0, T; H^1)$ and Lemma 7, we know that there exists some $v \in L^\infty(0, T; L^2) \cap L^2(0, T; H^1)$ and some sub-sequence v_{ε_k} such that

$$(26) \quad \begin{aligned} v_{\varepsilon_k} &\rightharpoonup v \quad \text{weak* in } L^\infty(0, T; L^2) \\ v_{\varepsilon_k} &\rightharpoonup v \quad \text{weakly in } L^2(0, T; H^1) \end{aligned}$$

and

$$(27) \quad v_{\varepsilon_k} \rightarrow v \quad \text{strongly in } L^2(0, T; L_{\text{loc}}^2).$$

Let $\varphi \in C_0^\infty([0, T] \times \mathbb{R}^2)$ be a divergence-free vector field. We construct φ_{ε_k} as in Section 3, see relation (22). Since φ_{ε_k} is compactly supported in the set $\{|x - h_{\varepsilon_k}(t)| > \varepsilon_k\}$, we can use it as test function in (13) written for ε_k . We multiply (13) by φ_{ε_k} and integrate by parts in time and space to obtain that

$$(28) \quad \begin{aligned} - \int_0^T \int_{\mathbb{R}^2} v_{\varepsilon_k} \cdot \partial_t \varphi_{\varepsilon_k} + \nu \int_0^T \int_{\mathbb{R}^2} \nabla v_{\varepsilon_k} : \nabla \varphi_{\varepsilon_k} + \int_0^T \int_{\mathbb{R}^2} v_{\varepsilon_k} \cdot \nabla v_{\varepsilon_k} \cdot \varphi_{\varepsilon_k} \\ = \int_{\mathbb{R}^2} v_{\varepsilon_k}(0) \cdot \varphi_{\varepsilon_k}(0). \end{aligned}$$

We will pass to the limit $\varepsilon_k \rightarrow 0$ in each of the terms above.

First, we know by hypothesis that $v_{\varepsilon_k}(0) \rightharpoonup v_0$ weakly in L^2 . From Lemma 5 we also have that $\varphi_{\varepsilon_k}(0) \rightarrow \varphi(0)$ strongly in L^2 , so

$$(29) \quad \int_{\mathbb{R}^2} v_{\varepsilon_k}(0) \cdot \varphi_{\varepsilon_k}(0) \xrightarrow{\varepsilon_k \rightarrow 0} \int_{\mathbb{R}^2} v(0) \cdot \varphi(0).$$

Next, we also know from Lemma 5 that $\nabla \varphi_{\varepsilon_k} \rightarrow \nabla \varphi$ strongly in $L^2([0, T] \times \mathbb{R}^2)$. Given that $\nabla v_{\varepsilon_k} \rightharpoonup \nabla v$ weakly in $L^2([0, T] \times \mathbb{R}^2)$, see relation (26), we infer that

$$(30) \quad \int_0^T \int_{\mathbb{R}^2} \nabla v_{\varepsilon_k} : \nabla \varphi_{\varepsilon_k} \xrightarrow{\varepsilon_k \rightarrow 0} \int_0^T \int_{\mathbb{R}^2} \nabla v : \nabla \varphi.$$

The nonlinear term also passes to the limit quite easily. We decompose

$$\int_0^T \int_{\mathbb{R}^2} v_{\varepsilon_k} \cdot \nabla v_{\varepsilon_k} \cdot \varphi_{\varepsilon_k} = \int_0^T \int_{\mathbb{R}^2} v_{\varepsilon_k} \cdot \nabla v_{\varepsilon_k} \cdot \varphi + \int_0^T \int_{\mathbb{R}^2} v_{\varepsilon_k} \cdot \nabla v_{\varepsilon_k} \cdot (\varphi_{\varepsilon_k} - \varphi).$$

We know from (27) that $v_{\varepsilon_k} \rightarrow v$ strongly in $L^2(0, T; L^2_{\text{loc}})$, from (26) that $\nabla v_{\varepsilon_k} \rightharpoonup \nabla v$ weakly in $L^2(0, T; L^2)$. Recalling that φ is compactly supported and since we obviously have that φ is uniformly bounded in space and time we can pass to the limit in the first term on the right-hand side:

$$\int_0^T \int_{\mathbb{R}^2} v_{\varepsilon_k} \cdot \nabla v_{\varepsilon_k} \cdot \varphi \xrightarrow{\varepsilon_k \rightarrow 0} \int_0^T \int_{\mathbb{R}^2} v \cdot \nabla v \cdot \varphi.$$

To pass to the limit in the second term we make an integration by parts and use the Hölder inequality, the Gagliardo-Nirenberg inequality $\|f\|_{L^4} \leq C\|f\|_{L^2}^{\frac{1}{2}}\|\nabla f\|_{L^2}^{\frac{1}{2}}$ and Lemma 5

$$\begin{aligned} \left| \int_0^T \int_{\mathbb{R}^2} v_{\varepsilon_k} \cdot \nabla v_{\varepsilon_k} \cdot (\varphi_{\varepsilon_k} - \varphi) \right| &= \left| \int_0^T \int_{\mathbb{R}^2} v_{\varepsilon_k} \otimes v_{\varepsilon_k} : \nabla(\varphi_{\varepsilon_k} - \varphi) \right| \\ &\leq \int_0^T \|v_{\varepsilon_k}\|_{L^4}^2 \|\nabla(\varphi_{\varepsilon_k} - \varphi)\|_{L^2} \\ &\leq C \int_0^T \|v_{\varepsilon_k}\|_{L^2} \|\nabla v_{\varepsilon_k}\|_{L^2} \|\nabla(\varphi_{\varepsilon_k} - \varphi)\|_{L^2} \\ &\leq CT^{\frac{1}{2}} \|v_{\varepsilon_k}\|_{L^\infty(0, T; L^2)} \|v_{\varepsilon_k}\|_{L^2(0, T; H^1)} \|\varphi_{\varepsilon_k} - \varphi\|_{L^\infty(0, T; H^1)} \\ &\xrightarrow{\varepsilon_k \rightarrow 0} 0. \end{aligned}$$

We infer that

$$(31) \quad \int_0^T \int_{\mathbb{R}^2} v_{\varepsilon_k} \cdot \nabla v_{\varepsilon_k} \cdot \varphi_{\varepsilon_k} \xrightarrow{\varepsilon_k \rightarrow 0} \int_0^T \int_{\mathbb{R}^2} v \cdot \nabla v \cdot \varphi.$$

The last term we need to pass to the limit is the term with the time-derivative. Thanks to Lemma 6 we can bound

$$\begin{aligned} \left| \int_0^T \int_{\mathbb{R}^2} v_{\varepsilon_k} \cdot (\partial_t \varphi_{\varepsilon_k} - \partial_t \varphi) dx \right| &\leq C \int_0^T \|\text{curl } v_{\varepsilon_k}\|_{L^2} \left(\frac{\varepsilon_k^2 \alpha_{\varepsilon_k}^2}{\ln \alpha_{\varepsilon_k}} \|\partial_t \varphi\|_{L^\infty} + \frac{\varepsilon_k \alpha_{\varepsilon_k}}{\sqrt{\ln \alpha_{\varepsilon_k}}} |h'_{\varepsilon_k}(t)| \|\varphi\|_{L^\infty} \right) \\ &\leq CT^{\frac{1}{2}} \|v_{\varepsilon_k}\|_{L^2(0, T; H^1)} \|\varphi\|_{W^{1, \infty}(0, T; L^\infty)} \max \left(\frac{\varepsilon_k^2 \alpha_{\varepsilon_k}^2}{\ln \alpha_{\varepsilon_k}}, \frac{\varepsilon_k \alpha_{\varepsilon_k}}{\sqrt{\ln \alpha_{\varepsilon_k}}} |h'_{\varepsilon_k}(t)| \right) \\ &\xrightarrow{\varepsilon_k \rightarrow 0} 0 \end{aligned}$$

where we used (19). But we also have that

$$\int_0^T \int_{\mathbb{R}^2} v_{\varepsilon_k} \cdot \partial_t \varphi \xrightarrow{\varepsilon_k \rightarrow 0} \int_0^T \int_{\mathbb{R}^2} v \cdot \partial_t \varphi$$

so we can conclude that

$$(32) \quad \int_0^T \int_{\mathbb{R}^2} v_{\varepsilon_k} \partial_t \varphi_{\varepsilon_k} \xrightarrow{\varepsilon_k \rightarrow 0} \int_0^T \int_{\mathbb{R}^2} v \cdot \partial_t \varphi.$$

Gathering (28), (29), (30), (31) and (32), we conclude that

$$- \int_0^T \int_{\mathbb{R}^2} v \cdot \partial_t \varphi + \nu \int_0^T \int_{\mathbb{R}^2} \nabla v : \nabla \varphi + \int_0^T \int_{\mathbb{R}^2} v \cdot \nabla v \cdot \varphi = \int_{\mathbb{R}^2} v(0) \cdot \varphi(0)$$

which is the weak formulation of Navier-Stokes equations in \mathbb{R}^2 . This completes the proof of Proposition 1.

ACKNOWLEDGMENTS

J.H. and D.I. have been partially funded by the ANR project Dyficolti ANR-13-BS01-0003-01. D.I. has been partially funded by the LABEX MILYON (ANR-10-LABX-0070) of Université de Lyon, within the program “Investissements d’Avenir” (ANR-11-IDEX-0007) operated by the French National Research Agency (ANR).

REFERENCES

- [1] M. Chipot, G. Planas, J. C. Robinson, and W. Xue. Limits of the Stokes and Navier-Stokes equations in a punctured periodic domain. *arXiv:1407.6942 [math]*, 2014.
- [2] B. Desjardins and M. J. Esteban. Existence of weak solutions for the motion of rigid bodies in a viscous fluid. *Archive for Rational Mechanics and Analysis*, 146(1):59–71, 1999.
- [3] S. Ervedoza, M. Hillairet, and C. Lacave. Long-Time Behavior for the Two-Dimensional Motion of a Disk in a Viscous Fluid. *Communications in Mathematical Physics*, 329(1):325–382, 2014.
- [4] O. Glass, C. Lacave, and F. Sueur. On the motion of a small body immersed in a two dimensional incompressible perfect fluid. *Bulletin de la Société Mathématique de France*, 142(3):489–536, 2014.
- [5] O. Glass, C. Lacave, and F. Sueur. On the Motion of a Small Light Body Immersed in a Two Dimensional Incompressible Perfect Fluid with Vorticity. *Communications in Mathematical Physics*, 341(3):1015–1065, 2016.
- [6] O. Glass, A. Munnier, and F. Sueur. Point vortex dynamics as zero-radius limit of the motion of a rigid body in an irrotational fluid. *Inventiones mathematicae*, 214(1):171–287, 2018.
- [7] K.-H. Hoffmann and V. N. Starovoitov. On a motion of a solid body in a viscous fluid. Two-dimensional case. *Advances in Mathematical Sciences and Applications*, 9(2):633–648, 1999.
- [8] D. Iftimie, M. C. Lopes Filho, and H. J. Nussenzweig Lopes. Two Dimensional Incompressible Ideal Flow Around a Small Obstacle. *Communications in Partial Differential Equations*, 28(1-2):349–379, 2003.
- [9] D. Iftimie, M. C. Lopes Filho, and H. J. Nussenzweig Lopes. Two-dimensional incompressible viscous flow around a small obstacle. *Mathematische Annalen*, 336(2):449–489, 2006.
- [10] C. Lacave. Two-dimensional incompressible viscous flow around a thin obstacle tending to a curve. *Proceedings of the Royal Society of Edinburgh Section A: Mathematics*, 139(6):1237–1254, 2009.
- [11] C. Lacave and T. Takahashi. Small Moving Rigid Body into a Viscous Incompressible Fluid. *Archive for Rational Mechanics and Analysis*, 223(3):1307–1335, 2017.
- [12] J.-L. Lions. *Quelques méthodes de résolution des problèmes aux limites non linéaires*. Dunod, 1969.
- [13] J. San Martín, V. Starovoitov, and M. Tucsnak. Global Weak Solutions for the Two-Dimensional Motion of Several Rigid Bodies in an Incompressible Viscous Fluid. *Archive for Rational Mechanics and Analysis*, 161(2):113–147, 2002.
- [14] D. Serre. Chute Libre d’un Solide dans un Fluide Visqueux Incompressible. Existence. *Japan Journal of Industrial and Applied Mathematics*, 4(1):99–110, 1987.
- [15] T. Takahashi and M. Tucsnak. Global strong solutions for the two-dimensional motion of an infinite cylinder in a viscous fluid. *Journal of Mathematical Fluid Mechanics*, 6(1):53–77, 2004.

- [16] N. V. Yudakov. The solvability of the problem of the motion of a rigid body in a viscous incompressible fluid. *Dinamika Sploshnoi Sredy*, 18:249–253, 1974.

J. He: Université de Lyon, Université Lyon 1 – CNRS UMR 5208 Institut Camille Jordan – 43 bd. du 11 Novembre 1918 – Villeurbanne Cedex F-69622, France.

Email: `jiao.he@math.univ-lyon1.fr`

D. Iftimie: Université de Lyon, Université Lyon 1 – CNRS UMR 5208 Institut Camille Jordan – 43 bd. du 11 Novembre 1918 – Villeurbanne Cedex F-69622, France.

Email: `iftimie@math.univ-lyon1.fr`