Homogeneity and algebraic closure in free groups

Abderezak Ould Houcine

Université de Mons, Université Lyon 1

Anglo-Belgian Workshop in Model Theory & Applications[2]

14 september 2010

Contents

1 Homogeneity & prime models
- Definitions
- Existential homogeneity & prime models
- Homogeneity

2 Algebraic & definable closure
- Definitions
- Constructibility over the algebraic closure
- A counterexample

Contents

1 Homogeneity & prime models
- Definitions
- Existential homogeneity & prime models
- Homogeneity

2 Algebraic & definable closure
- Definitions
- Constructibility over the algebraic closure
- A counterexample

Homogeneity & existential homogeneity

Homogeneity & existential homogeneity

Let $\mathcal{M}$ be a countable model.

Homogeneity & existential homogeneity

Let $\mathcal{M}$ be a countable model.

- Let $\bar{a} \in \mathcal{M}^n$.

Homogeneity & existential homogeneity

Let $\mathcal{M}$ be a countable model.

- Let $\bar{a} \in \mathcal{M}^n$. The type of $\bar{a}$ is defined by

Homogeneity & existential homogeneity

Let $\mathcal{M}$ be a countable model.

- Let $\bar{a} \in \mathcal{M}^n$. The type of $\bar{a}$ is defined by

$$tp^{\mathcal{M}}(\bar{a}) = \{\psi(\bar{x}) | \mathcal{M} \models \psi(\bar{a})\}.$$

Homogeneity & existential homogeneity

Let $\mathcal{M}$ be a countable model.

- Let $\bar{a} \in \mathcal{M}^n$. The type of $\bar{a}$ is defined by

$$tp^{\mathcal{M}}(\bar{a}) = \{\psi(\bar{x}) | \mathcal{M} \models \psi(\bar{a})\}.$$

- $\mathcal{M}$ is *homogeneous*

Homogeneity & existential homogeneity

Let $\mathcal{M}$ be a countable model.

- Let $\bar{a} \in \mathcal{M}^n$. The type of $\bar{a}$ is defined by

$$tp^{\mathcal{M}}(\bar{a}) = \{\psi(\bar{x}) | \mathcal{M} \models \psi(\bar{a})\}.$$

- $\mathcal{M}$ is *homogeneous* $\Leftrightarrow$ for any $n \geq 1$, for any $\bar{a}$, $\bar{b} \in \mathcal{M}^n$,

Homogeneity & existential homogeneity

Let $\mathcal{M}$ be a countable model.

- Let $\bar{a} \in \mathcal{M}^n$. The type of $\bar{a}$ is defined by

$$tp^{\mathcal{M}}(\bar{a}) = \{\psi(\bar{x}) | \mathcal{M} \models \psi(\bar{a})\}.$$

- $\mathcal{M}$ is *homogeneous* $\Leftrightarrow$ for any $n \geq 1$, for any $\bar{a}$, $\bar{b} \in \mathcal{M}^n$, $tp^{\mathcal{M}}(\bar{a}) = tp^{\mathcal{M}}(\bar{b}) \implies \exists f \in Aut(\mathcal{M})$ s.t. $f(\bar{a}) = f(\bar{b})$.

Homogeneity & existential homogeneity

Let $\mathcal{M}$ be a countable model.

- Let $\bar{a} \in \mathcal{M}^n$. The type of $\bar{a}$ is defined by

$$tp^{\mathcal{M}}(\bar{a}) = \{\psi(\bar{x}) | \mathcal{M} \models \psi(\bar{a})\}.$$

- $\mathcal{M}$ is *homogeneous* $\Leftrightarrow$ for any $n \geq 1$, for any $\bar{a}$, $\bar{b} \in \mathcal{M}^n$, $tp^{\mathcal{M}}(\bar{a}) = tp^{\mathcal{M}}(\bar{b}) \implies \exists f \in Aut(\mathcal{M})$ s.t. $f(\bar{a}) = f(\bar{b})$.
- $\mathcal{M}$ is $\exists$-*homogeneous*

Homogeneity & existential homogeneity

Let $\mathcal{M}$ be a countable model.

- Let $\bar{a} \in \mathcal{M}^n$. The type of $\bar{a}$ is defined by

$$tp^{\mathcal{M}}(\bar{a}) = \{\psi(\bar{x}) | \mathcal{M} \models \psi(\bar{a})\}.$$

- $\mathcal{M}$ is *homogeneous* $\Leftrightarrow$ for any $n \geq 1$, for any $\bar{a}$, $\bar{b} \in \mathcal{M}^n$, $tp^{\mathcal{M}}(\bar{a}) = tp^{\mathcal{M}}(\bar{b}) \implies \exists f \in Aut(\mathcal{M})$ s.t. $f(\bar{a}) = f(\bar{b})$.
- $\mathcal{M}$ is $\exists$-*homogeneous* $\Leftrightarrow$ for any $n \geq 1$, for any $\bar{a}$, $\bar{b} \in \mathcal{M}^n$,

Homogeneity & existential homogeneity

Let $\mathcal{M}$ be a countable model.

- Let $\bar{a} \in \mathcal{M}^n$. The type of $\bar{a}$ is defined by

$$tp^{\mathcal{M}}(\bar{a}) = \{\psi(\bar{x}) | \mathcal{M} \models \psi(\bar{a})\}.$$

- $\mathcal{M}$ is *homogeneous* $\Leftrightarrow$ for any $n \geq 1$, for any $\bar{a}$, $\bar{b} \in \mathcal{M}^n$, $tp^{\mathcal{M}}(\bar{a}) = tp^{\mathcal{M}}(\bar{b}) \implies \exists f \in Aut(\mathcal{M})$ s.t. $f(\bar{a}) = f(\bar{b})$.
- $\mathcal{M}$ is $\exists$*-homogeneous* $\Leftrightarrow$ for any $n \geq 1$, for any $\bar{a}$, $\bar{b} \in \mathcal{M}^n$, $tp^{\mathcal{M}}_{\exists}(\bar{a}) = tp^{\mathcal{M}}_{\exists}(\bar{b}) \implies \exists f \in Aut(\mathcal{M})$ s.t. $f(\bar{a}) = f(\bar{b})$.

Homogeneity & existential homogeneity

Let $\mathcal{M}$ be a countable model.

- Let $\bar{a} \in \mathcal{M}^n$. The type of $\bar{a}$ is defined by

$$tp^{\mathcal{M}}(\bar{a}) = \{\psi(\bar{x}) | \mathcal{M} \models \psi(\bar{a})\}.$$

- $\mathcal{M}$ is *homogeneous* $\Leftrightarrow$ for any $n \geq 1$, for any $\bar{a}$, $\bar{b} \in \mathcal{M}^n$, $tp^{\mathcal{M}}(\bar{a}) = tp^{\mathcal{M}}(\bar{b}) \implies \exists f \in Aut(\mathcal{M})$ s.t. $f(\bar{a}) = f(\bar{b})$.
- $\mathcal{M}$ is $\exists$*-homogeneous* $\Leftrightarrow$ for any $n \geq 1$, for any $\bar{a}$, $\bar{b} \in \mathcal{M}^n$, $tp^{\mathcal{M}}_{\exists}(\bar{a}) = tp^{\mathcal{M}}_{\exists}(\bar{b}) \implies \exists f \in Aut(\mathcal{M})$ s.t. $f(\bar{a}) = f(\bar{b})$.

Remark.

Homogeneity & existential homogeneity

Let $\mathcal{M}$ be a countable model.

- Let $\bar{a} \in \mathcal{M}^n$. The type of $\bar{a}$ is defined by

$$tp^{\mathcal{M}}(\bar{a}) = \{\psi(\bar{x}) | \mathcal{M} \models \psi(\bar{a})\}.$$

- $\mathcal{M}$ is *homogeneous* $\Leftrightarrow$ for any $n \geq 1$, for any $\bar{a}$, $\bar{b} \in \mathcal{M}^n$, $tp^{\mathcal{M}}(\bar{a}) = tp^{\mathcal{M}}(\bar{b}) \implies \exists f \in Aut(\mathcal{M})$ s.t. $f(\bar{a}) = f(\bar{b})$.
- $\mathcal{M}$ is $\exists$*-homogeneous* $\Leftrightarrow$ for any $n \geq 1$, for any $\bar{a}$, $\bar{b} \in \mathcal{M}^n$, $tp^{\mathcal{M}}_{\exists}(\bar{a}) = tp^{\mathcal{M}}_{\exists}(\bar{b}) \implies \exists f \in Aut(\mathcal{M})$ s.t. $f(\bar{a}) = f(\bar{b})$.

Remark. $\exists$-homogeneity $\implies$ homogeneity.

Prime Models

Prime Models

- $\mathcal{M}$ is called *prime*

Prime Models

- $\mathcal{M}$ is called *prime* if $\mathcal{M}$ is elementary embeddable in every model of $Th(\mathcal{M})$.

The free group of rank 2

The free group of rank 2

Theorem 1 (A. Nies, 2003)

The free group of rank 2

Theorem 1 (A. Nies, 2003)

A free group F_2 of rank 2 is $\exists$-homogeneous and **not** *prime.*

The free group of rank 2

Theorem 1 (A. Nies, 2003)

A free group F_2 of rank 2 is $\exists$-homogeneous and **not** *prime. In particular $Th(F_2)$ has no prime model.*

The free group of rank 2

Theorem 1 (A. Nies, 2003)

A free group F_2 of rank 2 is $\exists$-homogeneous and **not** *prime. In particular $Th(F_2)$ has no prime model.*

The proof uses the following strong property of the free group F_2 with basis $\{a, b\}$:

The free group of rank 2

Theorem 1 (A. Nies, 2003)

A free group F_2 of rank 2 is $\exists$-homogeneous and **not** *prime. In particular $Th(F_2)$ has no prime model.*

The proof uses the following strong property of the free group F_2 with basis $\{a, b\}$: there exists a quantifier-free formula $\varphi(x, y)$,

The free group of rank 2

Theorem 1 (A. Nies, 2003)

A free group F_2 of rank 2 is $\exists$-homogeneous and **not** *prime. In particular $Th(F_2)$ has no prime model.*

The proof uses the following strong property of the free group F_2 with basis $\{a, b\}$: there exists a quantifier-free formula $\varphi(x, y)$, such that for any endomorphism f of F_2 if $F_2 \models \varphi(f(a), f(b))$ then f is an embedding.

The free group of rank 2

Theorem 1 (A. Nies, 2003)

A free group F_2 of rank 2 is $\exists$-homogeneous and **not** *prime. In particular $Th(F_2)$ has no prime model.*

The proof uses the following strong property of the free group F_2 with basis $\{a, b\}$: there exists a quantifier-free formula $\varphi(x, y)$, such that for any endomorphism f of F_2 if $F_2 \models \varphi(f(a), f(b))$ then f is an embedding.

Indeed, we can take $\varphi(x, y) := [x, y] \neq 1$

Two-generated torsion-free hyperbolic groups

Two-generated torsion-free hyperbolic groups

Question:

Two-generated torsion-free hyperbolic groups

Question: What can be said about the $\exists$-homogeneity and "primeness" of two-generated torsion-free hyperbolic groups?

Two-generated torsion-free hyperbolic groups

Question: What can be said about the $\exists$-homogeneity and "primeness" of two-generated torsion-free hyperbolic groups?

Definition

A group G is said to be co-hopfian, if any injective endomorphism of G is an automorphism.

Two-generated torsion-free hyperbolic groups

Question: What can be said about the $\exists$-homogeneity and "primeness" of two-generated torsion-free hyperbolic groups?

Definition

A group G is said to be co-hopfian, if any injective endomorphism of G is an automorphism.

That is a group is co-hopfian if it does not contain a subgroup isomorphic to itself.

Two-generated torsion-free hyperbolic groups

Two-generated torsion-free hyperbolic groups

Examples:

Two-generated torsion-free hyperbolic groups

Examples:

- Finite groups, the group of the rationals $\mathbb{Q}$.

Two-generated torsion-free hyperbolic groups

Examples:

- Finite groups, the group of the rationals $\mathbb{Q}$.
- $SL_n(\mathbb{Z})$ with $n \geq 3$ (G. Prasad, 1976).

Two-generated torsion-free hyperbolic groups

Examples:

- Finite groups, the group of the rationals $\mathbb{Q}$.
- $SL_n(\mathbb{Z})$ with $n \geq 3$ (G. Prasad, 1976).
- Nonabelian freely indecomposable trosion-free hyperbolic groups (Z. Sela, 1997).

Two-generated torsion-free hyperbolic groups

Examples:

- Finite groups, the group of the rationals $\mathbb{Q}$.
- $SL_n(\mathbb{Z})$ with $n \geq 3$ (G. Prasad, 1976).
- Nonabelian freely indecomposable trosion-free hyperbolic groups (Z. Sela, 1997).
- Mapping class groups of closed surfaces (N.V. Ivanov, J.D. McCarthy, 1999).

Two-generated torsion-free hyperbolic groups

Examples:

- Finite groups, the group of the rationals $\mathbb{Q}$.
- $SL_n(\mathbb{Z})$ with $n \geq 3$ (G. Prasad, 1976).
- Nonabelian freely indecomposable trosion-free hyperbolic groups (Z. Sela, 1997).
- Mapping class groups of closed surfaces (N.V. Ivanov, J.D. McCarthy, 1999).
- $B_n/Z(B_n)$, where B_n is the Braid group on $n \geq 4$ strands (R.W. Bell, D. Margalit, 2005).

Two-generated torsion-free hyperbolic groups

Examples:

- Finite groups, the group of the rationals $\mathbb{Q}$.
- $SL_n(\mathbb{Z})$ with $n \geq 3$ (G. Prasad, 1976).
- Nonabelian freely indecomposable trosion-free hyperbolic groups (Z. Sela, 1997).
- Mapping class groups of closed surfaces (N.V. Ivanov, J.D. McCarthy, 1999).
- $B_n/Z(B_n)$, where B_n is the Braid group on $n \geq 4$ strands (R.W. Bell, D. Margalit, 2005).
- $Out(F_n)$, where F_n is a free group of rank n (B. Farb, M. Handel, 2007).

Two-generated torsion-free hyperbolic groups

Two-generated torsion-free hyperbolic groups

We introduce a strong form of the co-hopf property.

Two-generated torsion-free hyperbolic groups

We introduce a strong form of the co-hopf property.

Definition

Two-generated torsion-free hyperbolic groups

We introduce a strong form of the co-hopf property.

Definition

A group G is said to be **strongly co-hopfian**,

Two-generated torsion-free hyperbolic groups

We introduce a strong form of the co-hopf property.

Definition

A group G is said to be **strongly co-hopfian**, *if there exists a finite subset $S \subseteq G \setminus \{1\}$*

Two-generated torsion-free hyperbolic groups

We introduce a strong form of the co-hopf property.

Definition

A group G is said to be **strongly co-hopfian**, *if there exists a finite subset $S \subseteq G \setminus \{1\}$ such that for any endomorphism φ of G, if $1 \notin \varphi(S)$ then φ is an automorphism.*

Two-generated torsion-free hyperbolic groups

We introduce a strong form of the co-hopf property.

Definition

A group G is said to be **strongly co-hopfian**, *if there exists a finite subset $S \subseteq G \setminus \{1\}$ such that for any endomorphism φ of G, if $1 \notin \varphi(S)$ then φ is an automorphism.*

Examples:

Two-generated torsion-free hyperbolic groups

We introduce a strong form of the co-hopf property.

Definition

A group G is said to be **strongly co-hopfian**, *if there exists a finite subset $S \subseteq G \setminus \{1\}$ such that for any endomorphism φ of G, if $1 \notin \varphi(S)$ then φ is an automorphism.*

Examples:

- Finite groups, the group of the rationals $\mathbb{Q}$.

Two-generated torsion-free hyperbolic groups

We introduce a strong form of the co-hopf property.

Definition

A group G is said to be **strongly co-hopfian**, *if there exists a finite subset $S \subseteq G \setminus \{1\}$ such that for any endomorphism φ of G, if $1 \notin \varphi(S)$ then φ is an automorphism.*

Examples:

- Finite groups, the group of the rationals $\mathbb{Q}$.
- Tarski Monster groups.

The previous notion is very interesting from the view point of model theory:

The previous notion is very interesting from the view point of model theory:

Lemma

The previous notion is very interesting from the view point of model theory:

Lemma

A finitely presented strongly co-hopfian group is $\exists$-homogeneous and prime.

The previous notion is very interesting from the view point of model theory:

Lemma

A finitely presented strongly co-hopfian group is $\exists$-homogeneous and prime.

Let $G = \langle \bar{a} | r_1(\bar{a}), r_2(\bar{a}), \ldots, r_n(\bar{a}) \rangle$ be finitely presented and strongly co-hopfian.

The previous notion is very interesting from the view point of model theory:

Lemma

A finitely presented strongly co-hopfian group is $\exists$-homogeneous and prime.

Let $G = \langle \bar{a} | r_1(\bar{a}), r_2(\bar{a}), \ldots, r_n(\bar{a}) \rangle$ be finitely presented and strongly co-hopfian. Then there exists a quantifier-free formula $\varphi(\bar{x})$ such that for any endomorphism f of G, if $G \models \varphi(f(\bar{a}))$ then f is an automorphism.

The previous notion is very interesting from the view point of model theory:

Lemma

A finitely presented strongly co-hopfian group is $\exists$-homogeneous and prime.

Let $G = \langle \bar{a} | r_1(\bar{a}), r_2(\bar{a}), \ldots, r_n(\bar{a}) \rangle$ be finitely presented and strongly co-hopfian. Then there exists a quantifier-free formula $\varphi(\bar{x})$ such that for any endomorphism f of G, if $G \models \varphi(f(\bar{a}))$ then f is an automorphism. A property analogous to that of the free group of rank 2.

Two-generated torsion-free hyperbolic groups

Two-generated torsion-free hyperbolic groups

Theorem 2

Two-generated torsion-free hyperbolic groups

Theorem 2

A non-free two-generated torsion-free hyperbolic group is strongly co-hopfian.

Two-generated torsion-free hyperbolic groups

Two-generated torsion-free hyperbolic groups

Since torsion-free hyperbolic groups are finitely presented, we conclude by the previous Lemma:

Two-generated torsion-free hyperbolic groups

Since torsion-free hyperbolic groups are finitely presented, we conclude by the previous Lemma:

Theorem 3

Two-generated torsion-free hyperbolic groups

Since torsion-free hyperbolic groups are finitely presented, we conclude by the previous Lemma:

Theorem 3

A non-free two-generated torsion-free hyperbolic group is $\exists$-homogeneous and prime.

Sketch of proof of Theorem 2

Sketch of proof of Theorem 2

Let Γ be a non-free two-generated trosion-free hyperbolic group.

Sketch of proof of Theorem 2

Let Γ be a non-free two-generated trosion-free hyperbolic group.

- There exists a sequence of subroups $\Gamma = \Gamma_1 \geq \Gamma_2 \geq \cdots \geq \Gamma_n$ satisfying the following properties:

Sketch of proof of Theorem 2

Let Γ be a non-free two-generated trosion-free hyperbolic group.

- There exists a sequence of subroups $\Gamma = \Gamma_1 \geq \Gamma_2 \geq \cdots \geq \Gamma_n$ satisfying the following properties:

  (i) Each Γ_i is two-generated, hyperbolic and quasiconvex;

Sketch of proof of Theorem 2

Let Γ be a non-free two-generated trosion-free hyperbolic group.

- There exists a sequence of subroups $\Gamma = \Gamma_1 \geq \Gamma_2 \geq \cdots \geq \Gamma_n$ satisfying the following properties:

  (i) Each Γ_i is two-generated, hyperbolic and quasiconvex;
  (ii) $\Gamma_i = \langle \Gamma_{i+1}, t | A^t = B \rangle$, where A and B are a nontrivial malnormal cyclic subgroups of Γ_{i+1};

Sketch of proof of Theorem 2

Let Γ be a non-free two-generated trosion-free hyperbolic group.

- There exists a sequence of subroups $\Gamma = \Gamma_1 \geq \Gamma_2 \geq \cdots \geq \Gamma_n$ satisfying the following properties:

  (i) Each Γ_i is two-generated, hyperbolic and quasiconvex;
  (ii) $\Gamma_i = \langle \Gamma_{i+1}, t | A^t = B \rangle$, where A and B are a nontrivial malnormal cyclic subgroups of Γ_{i+1};
  (iii) Γ_n is a rigid subgroup of Γ.

Sketch of proof of Theorem 2

Sketch of proof of Theorem 2

- Γ_n is Γ-determined;

Sketch of proof of Theorem 2

- Γ_n is Γ-determined; that is, there exists a finite subset $S \subseteq G \setminus \{1\}$ such that for any homomorphism $\varphi : \Gamma_n \to \Gamma$, if $1 \notin \varphi(S)$ then φ is an embedding.

Sketch of proof of Theorem 2

- Γ_n is Γ-determined; that is, there exists a finite subset $S \subseteq G \setminus \{1\}$ such that for any homomorphism $\varphi : \Gamma_n \to \Gamma$, if $1 \notin \varphi(S)$ then φ is an embedding.
- Let φ be an endomorphism of Γ such that $1 \notin \varphi(S)$. Then the restriction of φ to every Γ_i is an automorphism of Γ_i.

Sketch of proof of Theorem 2

- Γ_n is Γ-determined; that is, there exists a finite subset $S \subseteq G \setminus \{1\}$ such that for any homomorphism $\varphi : \Gamma_n \to \Gamma$, if $1 \notin \varphi(S)$ then φ is an embedding.
- Let φ be an endomorphism of Γ such that $1 \notin \varphi(S)$. Then the restriction of φ to every Γ_i is an automorphism of Γ_i. In particular φ is an automorphism of Γ. $\square$

Quasi-axiomatizable groups

Quasi-axiomatizable groups

Definition (A. Nies)

Quasi-axiomatizable groups

Definition (A. Nies)

A finitely generated group G is said to be Quasi-Axiomatizable

Quasi-axiomatizable groups

Definition (A. Nies)

A finitely generated group G is said to be Quasi-Axiomatizable if any finitely generated group which is elementary equivalent to G is isomorphic to G.

Quasi-axiomatizable groups

Definition (A. Nies)

A finitely generated group G is said to be Quasi-Axiomatizable if any finitely generated group which is elementary equivalent to G is isomorphic to G.

Question (A. Nies):

Quasi-axiomatizable groups

Definition (A. Nies)

A finitely generated group G is said to be Quasi-Axiomatizable if any finitely generated group which is elementary equivalent to G is isomorphic to G.

Question (A. Nies): Is there a f.g. group which is prime but not QA?

Quasi-axiomatizable groups

Quasi-axiomatizable groups

Using Sela's work on the elementary theory of torsion-free hyperbolic groups, we have

Quasi-axiomatizable groups

Using Sela's work on the elementary theory of torsion-free hyperbolic groups, we have

Theorem 4

Quasi-axiomatizable groups

Using Sela's work on the elementary theory of torsion-free hyperbolic groups, we have

Theorem 4

Let Γ be a two-generated trosion-free hyperbolic group.

Quasi-axiomatizable groups

Using Sela's work on the elementary theory of torsion-free hyperbolic groups, we have

Theorem 4

Let Γ be a two-generated trosion-free hyperbolic group. Then Γ is elementary equivalent to $\Gamma * \mathbb{Z}$.

Quasi-axiomatizable groups

Using Sela's work on the elementary theory of torsion-free hyperbolic groups, we have

Theorem 4

Let Γ be a two-generated trosion-free hyperbolic group. Then Γ is elementary equivalent to $\Gamma * \mathbb{Z}$.

Hence

Quasi-axiomatizable groups

Using Sela's work on the elementary theory of torsion-free hyperbolic groups, we have

Theorem 4

Let Γ be a two-generated trosion-free hyperbolic group. Then Γ is elementary equivalent to $\Gamma * \mathbb{Z}$.

Hence if Γ is non-free then it is prime but not QA.

Existential homogeneity in free groups

Existential homogeneity in free groups

Question : What can be said about existential homogeneity in free groups?

Existential homogeneity in free groups

Question : What can be said about existential homogeneity in free groups?
Recall that:

Existential homogeneity in free groups

Question : What can be said about existential homogeneity in free groups?
Recall that:

Definition

Existential homogeneity in free groups

Question : What can be said about existential homogeneity in free groups?
Recall that:

Definition

Let $\mathcal{M}$ be a model and $\mathcal{N}$ a submodel of $\mathcal{M}$.

Existential homogeneity in free groups

Question : What can be said about existential homogeneity in free groups?
Recall that:

Definition

Let $\mathcal{M}$ be a model and $\mathcal{N}$ a submodel of $\mathcal{M}$. The model $\mathcal{N}$ is said to be existentially closed (abbreviated e.c.) in $\mathcal{M}$,

Existential homogeneity in free groups

Question : What can be said about existential homogeneity in free groups?
Recall that:

Definition

Let $\mathcal{M}$ be a model and $\mathcal{N}$ a submodel of $\mathcal{M}$. The model $\mathcal{N}$ is said to be existentially closed (abbreviated e.c.) in $\mathcal{M}$, if for any existential formula φ with parameters from $\mathcal{N}$, if $\mathcal{M} \models \varphi$, then $\mathcal{N} \models \varphi$.

Existential homogeneity in free groups

Question : What can be said about existential homogeneity in free groups?
Recall that:

Definition

Let $\mathcal{M}$ be a model and $\mathcal{N}$ a submodel of $\mathcal{M}$. The model $\mathcal{N}$ is said to be existentially closed (abbreviated e.c.) in $\mathcal{M}$, if for any existential formula φ with parameters from $\mathcal{N}$, if $\mathcal{M} \models \varphi$, then $\mathcal{N} \models \varphi$.

Definition

Existential homogeneity in free groups

Question : What can be said about existential homogeneity in free groups?
Recall that:

Definition

Let $\mathcal{M}$ be a model and $\mathcal{N}$ a submodel of $\mathcal{M}$. The model $\mathcal{N}$ is said to be existentially closed (abbreviated e.c.) in $\mathcal{M}$, if for any existential formula φ with parameters from $\mathcal{N}$, if $\mathcal{M} \models \varphi$, then $\mathcal{N} \models \varphi$.

Definition

Let F be a free group and let $\bar{a} = (a_1, \ldots, a_m)$ be a tuple from F.

Existential homogeneity in free groups

Question : What can be said about existential homogeneity in free groups?
Recall that:

Definition

Let $\mathcal{M}$ be a model and $\mathcal{N}$ a submodel of $\mathcal{M}$. The model $\mathcal{N}$ is said to be existentially closed (abbreviated e.c.) in $\mathcal{M}$, if for any existential formula φ with parameters from $\mathcal{N}$, if $\mathcal{M} \models \varphi$, then $\mathcal{N} \models \varphi$.

Definition

Let F be a free group and let $\bar{a} = (a_1, \ldots, a_m)$ be a tuple from F.We say that $\bar{a}$ is a power of a primitive element if there exist integers $p_1, \ldots, p_m$ and a primitive element u such that $a_i = u^{p_i}$ for all i.

Theorem 3

Theorem 3

Let F be a nonabelian free group of finite rank.

Theorem 3

Let F be a nonabelian free group of finite rank. Let $\bar{a}, \bar{b} \in F^n$ and $P \subseteq F$ such that $tp^F_\exists(\bar{a}|P) = tp^F_\exists(\bar{b}|P)$.

Theorem 3

Let F be a nonabelian free group of finite rank. Let $\bar{a}, \bar{b} \in F^n$ and $P \subseteq F$ such that $tp^F_\exists(\bar{a}|P) = tp^F_\exists(\bar{b}|P)$. Then one of the following cases holds:

Theorem 3

Let F be a nonabelian free group of finite rank. Let $\bar{a}, \bar{b} \in F^n$ and $P \subseteq F$ such that $tp^F_\exists(\bar{a}|P) = tp^F_\exists(\bar{b}|P)$. Then one of the following cases holds:

- *the tuple $\bar{a}$ has the same existential type as a power of a primitive element;*

Theorem 3

Let F be a nonabelian free group of finite rank. Let $\bar{a}, \bar{b} \in F^n$ and $P \subseteq F$ such that $tp^F_\exists(\bar{a}|P) = tp^F_\exists(\bar{b}|P)$. Then one of the following cases holds:

- *the tuple $\bar{a}$ has the same existential type as a power of a primitive element;*
- *there exists an e.c. subgroup $E(\bar{a})$ (resp. $E(\bar{b})$) containing P and $\bar{a}$ (resp. $\bar{b}$) and an isomorphism $\sigma : E(\bar{a}) \to E(\bar{b})$ fixing pointwise P and sending $\bar{a}$ to $\bar{b}$.*

Sketch of proof of Theorem 3

Sketch of proof of Theorem 3

We eliminate first parameters.

Sketch of proof of Theorem 3

We eliminate first parameters. Recall that a free group is *equationnally noetherian*;

Sketch of proof of Theorem 3

We eliminate first parameters. Recall that a free group is *equationnally noetherian*; that is any system of equations in finitely many variable is equivalent to a finite subsystem.

Sketch of proof of Theorem 3

We eliminate first parameters. Recall that a free group is *equationnally noetherian*; that is any system of equations in finitely many variable is equivalent to a finite subsystem.

Lemma

Sketch of proof of Theorem 3

We eliminate first parameters. Recall that a free group is *equationnally noetherian*; that is any system of equations in finitely many variable is equivalent to a finite subsystem.

Lemma

Let G be a finitely generated equationally noetherian group.

Sketch of proof of Theorem 3

We eliminate first parameters. Recall that a free group is *equationnally noetherian*; that is any system of equations in finitely many variable is equivalent to a finite subsystem.

Lemma

Let G be a finitely generated equationally noetherian group. Let P be a subset of G.

Sketch of proof of Theorem 3

We eliminate first parameters. Recall that a free group is *equationnally noetherian*; that is any system of equations in finitely many variable is equivalent to a finite subsystem.

Lemma

Let G be a finitely generated equationally noetherian group. Let P be a subset of G. Then there exists a finite subset $P_0 \subseteq P$ such that for any endomorphism f of G, if f fixes pointwise P_0 then f fixes pointwise P.

Sketch of proof of Theorem 3

We eliminate first parameters. Recall that a free group is *equationnally noetherian*; that is any system of equations in finitely many variable is equivalent to a finite subsystem.

Lemma

Let G be a finitely generated equationally noetherian group. Let P be a subset of G. Then there exists a finite subset $P_0 \subseteq P$ such that for any endomorphism f of G, if f fixes pointwise P_0 then f fixes pointwise P.

Hence:

Sketch of proof of Theorem 3

We eliminate first parameters. Recall that a free group is *equationnally noetherian*; that is any system of equations in finitely many variable is equivalent to a finite subsystem.

Lemma

Let G be a finitely generated equationally noetherian group. Let P be a subset of G. Then there exists a finite subset $P_0 \subseteq P$ such that for any endomorphism f of G, if f fixes pointwise P_0 then f fixes pointwise P.

Hence: $\exists f \in Aut(F|P)$ s.t. $f(\bar{a}) = f(\bar{b}) \Leftrightarrow \exists f \in Aut(F)$ s.t. $f(\bar{a}P_0) = f(\bar{b}P_0)$.

Sketch of proof of Theorem 3

We eliminate first parameters. Recall that a free group is *equationnally noetherian*; that is any system of equations in finitely many variable is equivalent to a finite subsystem.

Lemma

Let G be a finitely generated equationally noetherian group. Let P be a subset of G. Then there exists a finite subset $P_0 \subseteq P$ such that for any endomorphism f of G, if f fixes pointwise P_0 then f fixes pointwise P.

Hence: $\exists f \in Aut(F|P)$ s.t. $f(\bar{a}) = f(\bar{b}) \Leftrightarrow \exists f \in Aut(F)$ s.t. $f(\bar{a}P_0) = f(\bar{b}P_0)$.
Note that : If $tp_\exists(\bar{a}|P) = tp_\exists(\bar{b}|P)$ then $tp_\exists(\bar{a}|P_0) = tp_\exists(\bar{b}|P_0)$ and $tp_\exists(\bar{a}P_0) = tp_\exists(\bar{b}P_0)$.

Sketch of proof of Theorem 3

Sketch of proof of Theorem 3

Definition

Sketch of proof of Theorem 3

Definition

Let F_1 and F_2 be nonabelian free groups of finite rank and let $\bar{a}$ (resp. $\bar{b}$) be a tuple from F_1 (resp. F_2).

Sketch of proof of Theorem 3

Definition

Let F_1 and F_2 be nonabelian free groups of finite rank and let $\bar{a}$ (resp. $\bar{b}$) be a tuple from F_1 (resp. F_2). We say that $(\bar{a}, \bar{b})$ is existentially rigid, if there is no nontrivial free decomposition $F_1 = A * B$ such that A contains a tuple $\bar{c}$ with $tp_{\exists}^{F_1}(\bar{a}) \subseteq tp_{\exists}^{A}(\bar{c}) \subseteq tp_{\exists}^{F_2}(\bar{b})$.

Sketch of proof of Theorem 3

Sketch of proof of Theorem 3

Proposition (1)

Sketch of proof of Theorem 3

Proposition (1)

Let F_1 and F_2 be nonabelian free groups of finite rank and let $\bar{a}$ (resp. $\bar{b}$) be a tuple from F_1 (resp. F_2).

Sketch of proof of Theorem 3

Proposition (1)

Let F_1 and F_2 be nonabelian free groups of finite rank and let $\bar{a}$ (resp. $\bar{b}$) be a tuple from F_1 (resp. F_2). Suppose that $(\bar{a}, \bar{b})$ is existentially rigid and let $\bar{s}$ be a basis of F_1.

Sketch of proof of Theorem 3

Proposition (1)

Let F_1 and F_2 be nonabelian free groups of finite rank and let $\bar{a}$ (resp. $\bar{b}$) be a tuple from F_1 (resp. F_2). Suppose that $(\bar{a}, \bar{b})$ is existentially rigid and let $\bar{s}$ be a basis of F_1.
Then there exists a quantifier-free formula $\varphi(\bar{x}, \bar{y})$, such that $F_1 \models \varphi(\bar{a}, \bar{s})$ and such that for any $f \in Hom(F_1|\bar{a}, F_2|\bar{b})$, if $F_2 \models \varphi(\bar{b}, f(\bar{s}))$ then f is an embedding.

Sketch of proof of Theorem 3

Sketch of proof of Theorem 3

Proposition (2)

Sketch of proof of Theorem 3

Proposition (2)

Let F_1 and F_2 be nonabelian free groups of finite rank and $\bar{a}$ (resp. $\bar{b}$) a tuple from F_1 (resp. F_2) such that $tp_{\exists}^{F_1}(\bar{a}) = tp_{\exists}^{F_2}(\bar{b})$.

Sketch of proof of Theorem 3

Proposition (2)

Let F_1 and F_2 be nonabelian free groups of finite rank and $\bar{a}$ (resp. $\bar{b}$) a tuple from F_1 (resp. F_2) such that $tp_{\exists}^{F_1}(\bar{a}) = tp_{\exists}^{F_2}(\bar{b})$. Suppose that $(\bar{a}, \bar{b})$ is existentially rigid.

Sketch of proof of Theorem 3

Proposition (2)

Let F_1 and F_2 be nonabelian free groups of finite rank and $\bar{a}$ (resp. $\bar{b}$) a tuple from F_1 (resp. F_2) such that $tp_{\exists}^{F_1}(\bar{a}) = tp_{\exists}^{F_2}(\bar{b})$.
Suppose that $(\bar{a}, \bar{b})$ is existentially rigid.
Then either $rk(F_1) = 2$ and $\bar{a}$ is a power of a primitive element, or there exists an embedding $h \in Hom(F_1|\bar{a}, F_2|\bar{b})$ such that $h(F_1)$ is an e.c. subgroup of F_2.

Sketch of proof of Theorem 3

Sketch of proof of Theorem 3

Let F be a free group of finite rank and let $\bar{a}, \bar{b} \in F^n$ s.t. $tp^F_\exists(\bar{a}) = tp^F_\exists(\bar{b})$.

Sketch of proof of Theorem 3

Let F be a free group of finite rank and let $\bar{a}, \bar{b} \in F^n$ s.t. $tp^F_\exists(\bar{a}) = tp^F_\exists(\bar{b})$.
Let C be the smallest free factor of F such that C contains $\bar{c}$ with $tp^F_\exists(\bar{a}) = tp^C_\exists(\bar{c})$.

Sketch of proof of Theorem 3

Let F be a free group of finite rank and let $\bar{a}, \bar{b} \in F^n$ s.t. $tp^F_\exists(\bar{a}) = tp^F_\exists(\bar{b})$.
Let C be the smallest free factor of F such that C contains $\bar{c}$ with $tp^F_\exists(\bar{a}) = tp^C_\exists(\bar{c})$.
Then $(\bar{c}, \bar{a})$ and $(\bar{c}, \bar{b})$ are existentially rigid.

Sketch of proof of Theorem 3

Let F be a free group of finite rank and let $\bar{a}, \bar{b} \in F^n$ s.t. $tp^F_\exists(\bar{a}) = tp^F_\exists(\bar{b})$.
Let C be the smallest free factor of F such that C contains $\bar{c}$ with $tp^F_\exists(\bar{a}) = tp^C_\exists(\bar{c})$.
Then $(\bar{c}, \bar{a})$ and $(\bar{c}, \bar{b})$ are existentially rigid.
By Propsition (2), either $rk(C) = 2$ and $\bar{c}$ is a power of a primitive element or there exists an embedding $h_1 \in Hom(C|\bar{c}, F|\bar{a})$ (resp. $h_2 \in Hom(C|\bar{c}, F|\bar{b})$) such that $h_1(C)$ (resp. $h_2(C)$) is an e.c. subgroup of F.

Sketch of proof of Theorem 3

Let F be a free group of finite rank and let $\bar{a}, \bar{b} \in F^n$ s.t. $tp^F_\exists(\bar{a}) = tp^F_\exists(\bar{b})$.
Let C be the smallest free factor of F such that C contains $\bar{c}$ with $tp^F_\exists(\bar{a}) = tp^C_\exists(\bar{c})$.
Then $(\bar{c}, \bar{a})$ and $(\bar{c}, \bar{b})$ are existentially rigid.
By Propsition (2), either $rk(C) = 2$ and $\bar{c}$ is a power of a primitive element or there exists an embedding $h_1 \in Hom(C|\bar{c}, F|\bar{a})$ (resp. $h_2 \in Hom(C|\bar{c}, F|\bar{b})$) such that $h_1(C)$ (resp. $h_2(C)$) is an e.c. subgroup of F.
Suppose that $\bar{c}$ is not a power of a primitive element.

Sketch of proof of Theorem 3

Let F be a free group of finite rank and let $\bar{a}, \bar{b} \in F^n$ s.t. $tp^F_\exists(\bar{a}) = tp^F_\exists(\bar{b})$.
Let C be the smallest free factor of F such that C contains $\bar{c}$ with $tp^F_\exists(\bar{a}) = tp^C_\exists(\bar{c})$.
Then $(\bar{c}, \bar{a})$ and $(\bar{c}, \bar{b})$ are existentially rigid.
By Propsition (2), either $rk(C) = 2$ and $\bar{c}$ is a power of a primitive element or there exists an embedding $h_1 \in Hom(C|\bar{c}, F|\bar{a})$ (resp. $h_2 \in Hom(C|\bar{c}, F|\bar{b})$) such that $h_1(C)$ (resp. $h_2(C)$) is an e.c. subgroup of F.
Suppose that $\bar{c}$ is not a power of a primitive element. By setting $E(\bar{a}) = h_1(C)$ and $E(\bar{b}) = h_2(C)$, we have $h_2 \circ h_1^{-1} : E(\bar{a}) \to E(\bar{b})$ is an isomorphism with $h_2 \circ h_1^{-1}(\bar{a}) = \bar{b}$. □

Homogeneity in free groups

Homogeneity in free groups

Theorem 4

Homogeneity in free groups

Theorem 4

Let F be a nonabelian free group of finite rank.

Homogeneity in free groups

Theorem 4

Let F be a nonabelian free group of finite rank. For any tuples $\bar{a}, \bar{b} \in F^n$ and for any subset $P \subseteq F$, if $tp^F(\bar{a}|P) = tp^F(\bar{b}|P)$ then there exists an automorphism of F fixing pointwise P and sending $\bar{a}$ to $\bar{b}$.

Homogeneity in free groups

Theorem 4

Let F be a nonabelian free group of finite rank. For any tuples $\bar{a}, \bar{b} \in F^n$ and for any subset $P \subseteq F$, if $tp^F(\bar{a}|P) = tp^F(\bar{b}|P)$ then there exists an automorphism of F fixing pointwise P and sending $\bar{a}$ to $\bar{b}$.

The above theorem is also proved by Perin and Sklinos.

Sketch of proof of Theorem 4

Sketch of proof of Theorem 4

Z. Sela and O. Kharlampovich and A. Myasnikov show that nonabelian free groups have the same elementary theory, and in fact the following more explicit description.

Sketch of proof of Theorem 4

Z. Sela and O. Kharlampovich and A. Myasnikov show that nonabelian free groups have the same elementary theory, and in fact the following more explicit description.

Theorem 5

Sketch of proof of Theorem 4

Z. Sela and O. Kharlampovich and A. Myasnikov show that nonabelian free groups have the same elementary theory, and in fact the following more explicit description.

Theorem 5

A nonabelian free factor of a free group of finite rank is an elementary subgroup. □

Sketch of proof of Theorem 4

Sketch of proof of Theorem 4

They show also the following quantifier-elimination result.

Sketch of proof of Theorem 4

They show also the following quantifier-elimination result.

Theorem 6

Sketch of proof of Theorem 4

They show also the following quantifier-elimination result.

Theorem 6

Let $\varphi(\bar{x})$ be a formula.

Sketch of proof of Theorem 4

They show also the following quantifier-elimination result.

Theorem 6

Let $\varphi(\bar{x})$ be a formula. Then there exists a boolean combination of $\exists\forall$-forumulas $\phi(\bar{x})$, such that for any nonabelian free group F of finite rank, one has $F \models \forall\bar{x}(\varphi(\bar{x}) \Leftrightarrow \phi(\bar{x}))$. □

Sketch of proof of Theorem 4

They show also the following quantifier-elimination result.

Theorem 6

Let $\varphi(\bar{x})$ be a formula. Then there exists a boolean combination of $\exists\forall$-forumulas $\phi(\bar{x})$, such that for any nonabelian free group F of finite rank, one has $F \models \forall\bar{x}(\varphi(\bar{x}) \Leftrightarrow \phi(\bar{x}))$. □

We notice, in particular, that if $\bar{a}, \bar{b} \in F^n$ such that $tp^F_{\exists\forall}(\bar{a}) = tp^F_{\exists\forall}(\bar{b})$, then $tp^F(\bar{a}) = tp^F(\bar{b})$.

Sketch of proof of Theorem 4

Sketch of proof of Theorem 4

Theorem 7 (C. Perin, 2008)

Sketch of proof of Theorem 4

Theorem 7 (C. Perin, 2008)

An elementary subgroup of a free group of finite rank is a free factor. □

Sketch of proof of Theorem 4

Theorem 7 (C. Perin, 2008)

An elementary subgroup of a free group of finite rank is a free factor. □

Theorem 8 (A. Pillay, 2009)

Sketch of proof of Theorem 4

Theorem 7 (C. Perin, 2008)

An elementary subgroup of a free group of finite rank is a free factor. □

Theorem 8 (A. Pillay, 2009)

Let F be a nonabelian free group of finite rank and $u, v \in F$ such that $tp^F(u) = tp^F(v)$.

Sketch of proof of Theorem 4

Theorem 7 (C. Perin, 2008)

An elementary subgroup of a free group of finite rank is a free factor. □

Theorem 8 (A. Pillay, 2009)

Let F be a nonabelian free group of finite rank and $u, v \in F$ such that $tp^F(u) = tp^F(v)$. If u is primitive, then v is primitive. □

Sketch of proof of Theorem 4

Sketch of proof of Theorem 4

Definition

Sketch of proof of Theorem 4

Definition

Let F be nonabelian free group of finite rank and let $\bar{a}$ be a tuple of F.

Sketch of proof of Theorem 4

Definition

Let F be nonabelian free group of finite rank and let $\bar{a}$ be a tuple of F. We say that $\bar{a}$ is rigid if there is no nontrivial free decomposition $F = A * B$ such that A contains a tuple $\bar{c}$ with $tp^{F_1}_{\exists\forall}(\bar{a}) = tp^{A}_{\exists\forall}(\bar{c})$.

Sketch of proof of Theorem 4

Sketch of proof of Theorem 4

Proposition (3)

Sketch of proof of Theorem 4

Proposition (3)

Let F_1 and F_2 be nonabelian free groups of finite rank and let $\bar{a}$ (resp. $\bar{b}$) be a tuple from F_1 (resp. F_2) such that $tp^{F_1}_{\exists\forall}(\bar{a}) = tp^{F_2}_{\exists\forall}(\bar{b})$.

Sketch of proof of Theorem 4

Proposition (3)

Let F_1 and F_2 be nonabelian free groups of finite rank and let $\bar{a}$ (resp. $\bar{b}$) be a tuple from F_1 (resp. F_2) such that $tp^{F_1}_{\exists\forall}(\bar{a}) = tp^{F_2}_{\exists\forall}(\bar{b})$. Suppose that $\bar{a}$ is rigid.

Sketch of proof of Theorem 4

Proposition (3)

Let F_1 and F_2 be nonabelian free groups of finite rank and let $\bar{a}$ (resp. $\bar{b}$) be a tuple from F_1 (resp. F_2) such that $tp^{F_1}_{\exists\forall}(\bar{a}) = tp^{F_2}_{\exists\forall}(\bar{b})$. Suppose that $\bar{a}$ is rigid.
Then either $rk(F_1) = 2$ and $\bar{a}$ is a power of a primitive element, or there exists an embedding $h \in Hom(F_1|\bar{a}, F_2|\bar{b})$ such that $h(F_1) \preceq_{\exists\forall} F_2$.

Sketch of proof of Theorem 4

Sketch of proof of Theorem 4

Let F be a free group of finite rank and let $\bar{a}, \bar{b} \in F^n$ s.t. $tp^F(\bar{a}) = tp^F(\bar{b})$.

Sketch of proof of Theorem 4

Let F be a free group of finite rank and let $\bar{a}, \bar{b} \in F^n$ s.t. $tp^F(\bar{a}) = tp^F(\bar{b})$.
Let C be the smallest free factor of F such that C contains $\bar{c}$ with $tp^F(\bar{a}) = tp^C(\bar{c})$.

Sketch of proof of Theorem 4

Let F be a free group of finite rank and let $\bar{a}, \bar{b} \in F^n$ s.t. $tp^F(\bar{a}) = tp^F(\bar{b})$.
Let C be the smallest free factor of F such that C contains $\bar{c}$ with $tp^F(\bar{a}) = tp^C(\bar{c})$.
Then $\bar{c}$ is rigid.

Sketch of proof of Theorem 4

Let F be a free group of finite rank and let $\bar{a}, \bar{b} \in F^n$ s.t. $tp^F(\bar{a}) = tp^F(\bar{b})$.
Let C be the smallest free factor of F such that C contains $\bar{c}$ with $tp^F(\bar{a}) = tp^C(\bar{c})$.
Then $\bar{c}$ is rigid.
By Propsition (3), either $rk(C) = 2$ and $\bar{c}$ is a power of a primitive element or there exists an embedding $h_1 \in Hom(C|\bar{a}, F|\bar{b})$ (resp. $h_2 \in Hom(C|\bar{a}, F|\bar{b})$) such that $h_1(C) \preceq_{\exists\forall} F$ (resp. $h_2(C) \preceq_{\exists\forall} F$).

Sketch of proof of Theorem 4

Let F be a free group of finite rank and let $\bar{a}, \bar{b} \in F^n$ s.t. $tp^F(\bar{a}) = tp^F(\bar{b})$.
Let C be the smallest free factor of F such that C contains $\bar{c}$ with $tp^F(\bar{a}) = tp^C(\bar{c})$.
Then $\bar{c}$ is rigid.
By Propsition (3), either $rk(C) = 2$ and $\bar{c}$ is a power of a primitive element or there exists an embedding $h_1 \in Hom(C|\bar{a}, F|\bar{b})$ (resp. $h_2 \in Hom(C|\bar{a}, F|\bar{b})$) such that $h_1(C) \preceq_{\exists\forall} F$ (resp. $h_2(C) \preceq_{\exists\forall} F$).

If $\bar{c}$ is a power of a primitive element then the result follows from Theorem 8.

Sketch of proof of Theorem 4

Sketch of proof of Theorem 4

Suppose that $\bar{c}$ is not a power of a primitive element.

Sketch of proof of Theorem 4

Suppose that $\bar{c}$ is not a power of a primitive element. Set $E(\bar{a}) = h_1(C)$ and $E(\bar{b}) = h_2(C)$.

Sketch of proof of Theorem 4

Suppose that $\bar{c}$ is not a power of a primitive element. Set $E(\bar{a}) = h_1(C)$ and $E(\bar{b}) = h_2(C)$.
We have $h_2 \circ h_1^{-1} : E(\bar{a}) \to E(\bar{b})$ is an isomorphism with $h_2 \circ h_1^{-1}(\bar{a}) = \bar{b}$.

Sketch of proof of Theorem 4

Suppose that $\bar{c}$ is not a power of a primitive element. Set $E(\bar{a}) = h_1(C)$ and $E(\bar{b}) = h_2(C)$.
We have $h_2 \circ h_1^{-1} : E(\bar{a}) \to E(\bar{b})$ is an isomorphism with $h_2 \circ h_1^{-1}(\bar{a}) = \bar{b}$.
Since $E(\bar{a}) \preceq_{\exists\forall} F$ (resp. $E(\bar{b}) \preceq_{\exists\forall} F$) we get by Theorem 7 that $E(\bar{a}) \preceq F$ (resp. $E(\bar{b}) \preceq F$).

Sketch of proof of Theorem 4

Suppose that $\bar{c}$ is not a power of a primitive element. Set $E(\bar{a}) = h_1(C)$ and $E(\bar{b}) = h_2(C)$.
We have $h_2 \circ h_1^{-1} : E(\bar{a}) \to E(\bar{b})$ is an isomorphism with $h_2 \circ h_1^{-1}(\bar{a}) = \bar{b}$.
Since $E(\bar{a}) \preceq_{\exists\forall} F$ (resp. $E(\bar{b}) \preceq_{\exists\forall} F$) we get by Theorem 7 that $E(\bar{a}) \preceq F$ (resp. $E(\bar{b}) \preceq F$).
By Theorem 7, $E(\bar{a})$ (resp. $E(\bar{b})$) is a free factor of F.

Sketch of proof of Theorem 4

Suppose that $\bar{c}$ is not a power of a primitive element. Set $E(\bar{a}) = h_1(C)$ and $E(\bar{b}) = h_2(C)$.
We have $h_2 \circ h_1^{-1} : E(\bar{a}) \to E(\bar{b})$ is an isomorphism with $h_2 \circ h_1^{-1}(\bar{a}) = \bar{b}$.
Since $E(\bar{a}) \preceq_{\exists\forall} F$ (resp. $E(\bar{b}) \preceq_{\exists\forall} F$) we get by Theorem 7 that $E(\bar{a}) \preceq F$ (resp. $E(\bar{b}) \preceq F$).
By Theorem 7, $E(\bar{a})$ (resp. $E(\bar{b})$) is a free factor of F.
Therefore $h_2 \circ h_1^{-1}$ can be extended to an isomorphism of F. $\square$

Contents

1 Homogeneity & prime models
- Definitions
- Existential homogeneity & prime models
- Homogeneity

2 Algebraic & definable closure
- Definitions
- Constructibility over the algebraic closure
- A counterexample

Definitions

Definitions

This part is a joint work with D. Vallino.

Definitions

This part is a joint work with D. Vallino.
Recall that:

Definitions

This part is a joint work with D. Vallino.
Recall that:

Definition

Let G be a group and A a subset of G.

Definitions

This part is a joint work with D. Vallino.
Recall that:

Definition

Let G be a group and A a subset of G.

- *The algebraic closure of A, denoted $acl_G(A)$, is the set of elements $g \in G$ such that there exists a formula $\phi(x)$ with parameters from A such that $G \models \phi(g)$ and $\phi(G)$ is finite.*

Definitions

This part is a joint work with D. Vallino.
Recall that:

Definition

Let G be a group and A a subset of G.

- *The algebraic closure of A, denoted $acl_G(A)$, is the set of elements $g \in G$ such that there exists a formula $\phi(x)$ with parameters from A such that $G \models \phi(g)$ and $\phi(G)$ is finite.*
- *The definable closure of A, denoted $dcl_G(A)$, is the set of elements $g \in G$ such that there exists a formula $\phi(x)$ with parameters from A such that $G \models \phi(g)$ and $\phi(G)$ is a singleton.*

Question (Z. Sela, 2008):

Question (Z. Sela, 2008): Is it true that $acl(A) = dcl(A)$ in free groups?

Question (Z. Sela, 2008): Is it true that $acl(A) = dcl(A)$ in free groups?

Remarks.

Question (Z. Sela, 2008): Is it true that $acl(A) = dcl(A)$ in free groups?

Remarks. Let Γ be a torsion-free hyperbolic group and $A \subseteq \Gamma$.

Question (Z. Sela, 2008): Is it true that $acl(A) = dcl(A)$ in free groups?

Remarks. Let Γ be a torsion-free hyperbolic group and $A \subseteq \Gamma$.

- $acl(A) = acl(\langle A \rangle)$ and $dcl(A) = dcl(\langle A \rangle)$.

Question (Z. Sela, 2008): Is it true that $acl(A) = dcl(A)$ in free groups?

Remarks. Let Γ be a torsion-free hyperbolic group and $A \subseteq \Gamma$.

- $acl(A) = acl(\langle A \rangle)$ and $dcl(A) = dcl(\langle A \rangle)$. Hence, we may assume that A is a subgroup.

Question (Z. Sela, 2008): Is it true that $acl(A) = dcl(A)$ in free groups?

Remarks. Let Γ be a torsion-free hyperbolic group and $A \subseteq \Gamma$.

- $acl(A) = acl(\langle A \rangle)$ and $dcl(A) = dcl(\langle A \rangle)$. Hence, we may assume that A is a subgroup.
- If $\Gamma = \Gamma_1 * \Gamma_2$ and $A \leq \Gamma_1$ then $acl(A) \leq \Gamma_1$.

Question (Z. Sela, 2008): Is it true that $acl(A) = dcl(A)$ in free groups?

Remarks. Let Γ be a torsion-free hyperbolic group and $A \subseteq \Gamma$.

- $acl(A) = acl(\langle A \rangle)$ and $dcl(A) = dcl(\langle A \rangle)$. Hence, we may assume that A is a subgroup.
- If $\Gamma = \Gamma_1 * \Gamma_2$ and $A \leq \Gamma_1$ then $acl(A) \leq \Gamma_1$. Similarly for $dcl(A)$.

Question (Z. Sela, 2008): Is it true that $acl(A) = dcl(A)$ in free groups?

Remarks. Let Γ be a torsion-free hyperbolic group and $A \subseteq \Gamma$.

- $acl(A) = acl(\langle A \rangle)$ and $dcl(A) = dcl(\langle A \rangle)$. Hence, we may assume that A is a subgroup.
- If $\Gamma = \Gamma_1 * \Gamma_2$ and $A \leq \Gamma_1$ then $acl(A) \leq \Gamma_1$. Similarly for $dcl(A)$. Hence, we may assume that Γ is freely A-indecomposable.

Question (Z. Sela, 2008): Is it true that $acl(A) = dcl(A)$ in free groups?

Remarks. Let Γ be a torsion-free hyperbolic group and $A \subseteq \Gamma$.

- $acl(A) = acl(\langle A \rangle)$ and $dcl(A) = dcl(\langle A \rangle)$. Hence, we may assume that A is a subgroup.
- If $\Gamma = \Gamma_1 * \Gamma_2$ and $A \leq \Gamma_1$ then $acl(A) \leq \Gamma_1$. Similarly for $dcl(A)$. Hence, we may assume that Γ is freely A-indecomposable.
- If A is abelian then $acl(A) = dcl(A) = C_\Gamma(A)$.

Question (Z. Sela, 2008): Is it true that $acl(A) = dcl(A)$ in free groups?

Remarks. Let Γ be a torsion-free hyperbolic group and $A \subseteq \Gamma$.

- $acl(A) = acl(\langle A \rangle)$ and $dcl(A) = dcl(\langle A \rangle)$. Hence, we may assume that A is a subgroup.
- If $\Gamma = \Gamma_1 * \Gamma_2$ and $A \leq \Gamma_1$ then $acl(A) \leq \Gamma_1$. Similarly for $dcl(A)$. Hence, we may assume that Γ is freely A-indecomposable.
- If A is abelian then $acl(A) = dcl(A) = C_\Gamma(A)$. Hence , we may assume that A is nonabelian.

Theorem 9

Theorem 9

Let Γ be a torsion-free hyperbolic group and $A \leq \Gamma$ where A is nonabelian.

Theorem 9

Let Γ be a torsion-free hyperbolic group and $A \leq \Gamma$ where A is nonabelian.
Then Γ can be constructed from $acl(A)$ by a finite sequence of free products and HNN-extensions along cyclic subgroups.

Theorem 9

Let Γ be a torsion-free hyperbolic group and $A \leq \Gamma$ where A is nonabelian.
Then Γ can be constructed from $acl(A)$ by a finite sequence of free products and HNN-extensions along cyclic subgroups.
In particular, $acl(A)$ is finitely generated, quasiconvex and hyperbolic. □

Acl and the JSJ-decomposition

Acl and the JSJ-decomposition

Theorem 10

Acl and the JSJ-decomposition

Theorem 10

Let F be a nonabelian free group of finite rank and let $A \leq F$ be a nonabelian subgroup.

Acl and the JSJ-decomposition

Theorem 10

Let F be a nonabelian free group of finite rank and let $A \leq F$ be a nonabelian subgroup. Then:

Acl and the JSJ-decomposition

Theorem 10

Let F be a nonabelian free group of finite rank and let $A \leq F$ be a nonabelian subgroup. Then:

- *$dcl(A)$ is a free factor of $acl(A)$.*

Acl and the JSJ-decomposition

Theorem 10

Let F be a nonabelian free group of finite rank and let $A \leq F$ be a nonabelian subgroup. Then:

- *$dcl(A)$ is a free factor of $acl(A)$. In particular, if $rk(F) = 2$ then $acl(A) = dcl(A)$.*

Acl and the JSJ-decomposition

Theorem 10

Let F be a nonabelian free group of finite rank and let $A \leq F$ be a nonabelian subgroup. Then:

- *$dcl(A)$ is a free factor of $acl(A)$. In particular, if $rk(F) = 2$ then $acl(A) = dcl(A)$.*
- *$acl(A)$ is exactly the vertex group containing A in the cyclic JSJ-decomposition of F relative to A.*

A counterexample

A counterexample

Theorem 11

A counterexample

Theorem 11

Let A_0 be a finite set (possibly empty)

A counterexample

Theorem 11

Let A_0 be a finite set (possibly empty) and

$$A = \langle A_0, a, b, u | \rangle, \ H = A * \langle y | \rangle,$$

A counterexample

Theorem 11

Let A_0 be a finite set (possibly empty) and

$$A = \langle A_0, a, b, u | \rangle, \ H = A * \langle y | \rangle,$$

$$v = aybyay^{-1}by^{-1},$$

A counterexample

Theorem 11

Let A_0 be a finite set (possibly empty) and

$$A = \langle A_0, a, b, u | \rangle, \ H = A * \langle y | \rangle,$$

$$v = aybyay^{-1}by^{-1}, \ F = \langle H, t | u^t = v \rangle.$$

A counterexample

Theorem 11

Let A_0 be a finite set (possibly empty) and

$$A = \langle A_0, a, b, u | \rangle, \ H = A * \langle y | \rangle,$$

$$v = aybyay^{-1}by^{-1}, \ F = \langle H, t | u^t = v \rangle.$$

Then F is a free group of rank $|A_0| + 4$ and:

A counterexample

Theorem 11

Let A_0 be a finite set (possibly empty) and

$$A = \langle A_0, a, b, u | \rangle,\ H = A * \langle y | \rangle,$$

$$v = aybyay^{-1}by^{-1},\ F = \langle H, t | u^t = v \rangle.$$

Then F is a free group of rank $|A_0| + 4$ and:

- *If $f \in Hom(F|A)$ then $f \in Aut(F|A)$, and if $f_{|H} \neq 1$ then $f(y) = y^{-1}$.*

A counterexample

Theorem 11

Let A_0 be a finite set (possibly empty) and

$$A = \langle A_0, a, b, u | \rangle,\ H = A * \langle y | \rangle,$$

$$v = aybyay^{-1}by^{-1},\ F = \langle H, t | u^t = v \rangle.$$

Then F is a free group of rank $|A_0| + 4$ and:

- *If $f \in Hom(F|A)$ then $f \in Aut(F|A)$, and if $f_{|H} \neq 1$ then $f(y) = y^{-1}$.*
- *$acl(A) = acl^{\exists}(A) = H$.*

A counterexample

Theorem 11

Let A_0 be a finite set (possibly empty) and

$$A = \langle A_0, a, b, u | \rangle,\ H = A * \langle y | \rangle,$$

$$v = aybyay^{-1}by^{-1},\ F = \langle H, t | u^t = v \rangle.$$

Then F is a free group of rank $|A_0| + 4$ and:

- *If $f \in Hom(F|A)$ then $f \in Aut(F|A)$, and if $f_{|H} \neq 1$ then $f(y) = y^{-1}$.*
- *$acl(A) = acl^{\exists}(A) = H$.*
- *$dcl(A) = dcl^{\exists}(A) = A$.*