

ON THE COMPLEXITY OF TOPOLOGICAL CONJUGACY OF TOEPLITZ SUBSHIFTS

BY

MARCIN SABOK*

*Department of Mathematics and Statistics, McGill University
805, Sherbrooke Street West Montreal, Quebec, Canada H3A 2K6
and*

*Institute of Mathematics, Polish Academy of Sciences
Śniadeckich 8, 00-655 Warszawa, Poland
e-mail: marcin.sabok@mcgill.ca*

AND

TODOR TSANKOV**

*Institut de Mathématiques de Jussieu–PRG, Université Paris Diderot
75205 Paris cedex 13, France
e-mail: todor@math.univ-paris-diderot.fr*

ABSTRACT

In this paper, we study the descriptive set theoretic complexity of the equivalence relation of conjugacy of Toeplitz subshifts of a residually finite group G . On the one hand, we show that if $G = \mathbb{Z}$, then topological conjugacy on Toeplitz subshifts with separated holes is amenable. In contrast, if G is non-amenable, then conjugacy of Toeplitz G -subshifts is a non-amenable equivalence relation. The results were motivated by a general question, asked by Gao, Jackson and Seward, about the complexity of conjugacy for minimal, free subshifts of countable groups.

* The first author was partially supported by NSERC, by the Polish Ministry of Science and Higher Education (MNiSW) through the program Mobilność Plus and by the NCN through the grant 2012/05/D/ST1/03206.

** The second author was partially supported by the ANR contracts GrupoLoco (ANR-11-JS01-008) and GAMME (ANR-14-CE25-0004).

Received July 3, 2015 and in revised form April 18, 2016

1. Introduction

The theory of definable equivalence relations offers tools for classifying the complexity of equivalence relations arising from various isomorphism problems. An important class of Borel equivalence relations is given by the **countable** ones, i.e., those with countable classes. It is a classical result of Feldman and Moore that every such equivalence relation arises as the orbit equivalence relation of a Borel action of a countable group; and thus, from the beginning, the theory is intimately connected to that of dynamical systems from where it has borrowed most of its tools and techniques. The descriptive set theoretic approach to those equivalence relations has been developed over the last twenty years by Dougherty, Jackson, Kechris, Louveau, Hjorth, Thomas and others (see, e.g., [13, 5, 12]).

The natural comparison of equivalence relations is given by **Borel reducibility** and the simplest ones are those which are **smooth**, i.e., admit real numbers as complete invariants (or, equivalently, admit a Borel transversal). The next level of the complexity hierarchy is formed by the **hyperfinite** ones, equivalently, those induced by Borel actions of the group of integers \mathbb{Z} . There is also a **universal** countable Borel equivalence relation, which is maximal in the quasi-order of Borel reducibility; an example is the orbit equivalence relation $F_2 \curvearrowright 2^{F_2}$.

In topological dynamics, one of the most commonly considered types of dynamical systems are the subshifts. For a countable group G , the **Bernoulli shift** is the action of G on 2^G defined by the formula:

$$(g \cdot x)(h) = x(g^{-1}h)$$

for all $g, h \in G$, $x \in 2^G$. A **G -subshift** is a closed nonempty subset $S \subseteq 2^G$ which is invariant under the action of G . The natural isomorphism relation of subshifts is topological conjugacy: two subshifts S and T are **conjugate** if there is a homeomorphism $f: S \rightarrow T$ which commutes with the action of G . This is the equivalence relation that we study in this paper and usually we will refer to it as the **isomorphism relation** on G -subshifts. Often, special classes of subshifts are of interest in dynamics. A subshift S is **minimal** if it does not contain any proper subshift, or equivalently, if every orbit is dense. It is **free** if for any $x \in S$ and any $g \in G$ different from 1_G , we have $g \cdot x \neq x$.

It is worth noting that while many countable equivalence relations arise naturally from group actions, some do not, and their study is usually more difficult

because most of the available tools are dynamical in nature and require the presence of a (natural) group action. Some notable examples where there is no natural group action that gives the equivalence relation are Turing equivalence, isomorphism of (various classes of) finitely generated groups, and isomorphism of subshifts. (However, for subshifts, there is still a group action present that can be exploited: see Section 4.)

The recent monograph of Gao, Jackson and Seward [9] studies the complexity of isomorphism of free, minimal subshifts. (In fact, before their construction, it was an open problem whether such subshifts necessarily exist for every countable G .) It follows essentially from a classical result of Curtis, Hedlund and Lyndon (see [20] or [9, Lemma 9.2.1]) that for any countable group G , isomorphism of G -subshifts is a countable Borel equivalence relation. Gao, Jackson and Seward showed [9, Corollary 1.5.4] that if G is infinite, this equivalence relation is not smooth, and that if G is locally finite, then it is hyperfinite [9, Theorem 1.5.6]. They also pose the general question [9, Problem 9.4.11] to determine the complexity of this equivalence relation for an arbitrary countable group G . This was an important motivating question for our work and, in Theorem 1.2, we provide a partial answer.

Clemens [1] proved that isomorphism of \mathbb{Z} -subshifts is a universal countable Borel equivalence relation. However, his construction produces subshifts that are far from minimal and it remains an open question whether isomorphism of minimal \mathbb{Z} -subshifts is universal. This question is connected with the conjecture of Thomas [24, Conjecture 1.2] that isomorphism of finitely generated, amenable, simple groups is universal. It follows from the results of Matui [22], Giordano, Putnam and Skau [10] and a recent result of Juschenko and Monod [14] that the computation of the topological full group of a minimal \mathbb{Z} -subshift provides a reduction from flip-conjugacy of minimal \mathbb{Z} -subshifts to isomorphism of finitely generated, simple, amenable groups. Another related result was recently proved by J. Williams [25], who showed that isomorphism of finitely generated, solvable groups is weakly universal.

The focus of this paper is studying the complexity of the isomorphism relation for the class of Toeplitz subshifts of residually finite groups. This class of subshifts is well-known and appears in many contexts. For example, Downarowicz [7] showed that any Choquet simplex can be realized as the simplex of invariant measures of a Toeplitz subshift. (This result was recently generalized to arbitrary amenable, residually finite groups by Cortez and Petite [4].) An

important feature of Toeplitz words is that they can be constructed in stages, which allows a fair amount of control. We briefly recall the classical definition for $G = \mathbb{Z}$ here and postpone the general one for residually finite groups to Section 2.

A word $x \in 2^{\mathbb{Z}}$ is Toeplitz if every symbol occurs periodically, i.e., for every $n \in \mathbb{N}$, there exists k such that $x(n+ki) = x(n)$ for all $i \in \mathbb{Z}$. A subshift $S \subseteq 2^{\mathbb{Z}}$ is Toeplitz if it is equal to the closure of the orbit of some Toeplitz word. It is easy to check that every Toeplitz subshift is minimal.

The isomorphism relation for Toeplitz \mathbb{Z} -subshifts has been studied by Do-wnarowicz, Kwiatkowski and Lacroix [6] and it essentially follows from their results that isomorphism of **pointed** Toeplitz flows is a hyperfinite equivalence relation (cf. Proposition 5.1).

The above seems to indicate that the isomorphism relation for Toeplitz \mathbb{Z} -subshifts should not be too complicated. However, we have only been able to treat the case of Toeplitz subshifts with separated holes, a special but important class. The first of our main results is the following.

THEOREM 1.1: *For $G = \mathbb{Z}$, the isomorphism relation on Toeplitz subshifts with separated holes is amenable and therefore hyperfinite μ -a.e. for every Borel probability measure μ on the set of subshifts.*

We postpone the definitions to Section 3 and just recall that an amenable equivalence relation is hyperfinite a.e. with respect to any probability measure but it is an open problem whether it must be hyperfinite everywhere. We do not know whether the above equivalence relation is hyperfinite.

The second part of the paper deals with Toeplitz subshifts of residually finite, non-amenable groups. For those, we prove that the isomorphism relation is somewhat complicated.

THEOREM 1.2: *If G is a non-amenable, residually finite group, then the isomorphism relation on the set of free, Toeplitz G -subshifts is not hyperfinite.*

The proof of this theorem proceeds by constructing a probability measure μ on the set of Toeplitz subshifts which is invariant under a suitable action of the group G , included in the equivalence relation; then we show that the stabilizers of points are amenable and conclude that the equivalence relation is not μ -amenable and thus not hyperfinite.

In the last section of the paper, reinterpreting a result of Downarowicz, Kwiatkowski and Lacroix, we indicate how the isomorphism of Toeplitz \mathbb{Z} -subshifts is naturally generated by an action of a groupoid which is “hyperfinite-by-compact” (cf. Corollary 5.2). This seems to indicate that the equivalence relation is somewhat simple; however, we are not even able to prove that it is not universal. The following question remains open:

Question 1.3: Is the isomorphism relation of Toeplitz \mathbb{Z} -subshifts hyperfinite? Is this true for an arbitrary residually finite, amenable group?

ADDENDUM. In a very recent work, Kaya [15] has improved our Theorem 1.1 and shown that the equivalence relation of isomorphism on the set of Toeplitz subshifts with separated holes is actually hyperfinite.

ACKNOWLEDGEMENTS. This work was initiated during the stay of the first author at the Université Paris 7. He would like to thank the logic group in Paris for their hospitality. Both authors would like to thank Boban Veličković for many valuable discussions. The first author is indebted to Tomasz Downarowicz for valuable comments. We are also grateful to the anonymous referee for a careful reading of the paper and useful suggestions.

2. Toeplitz subshifts

In this section, we recall the definition and collect some basic properties of Toeplitz subshifts for residually finite groups. We also establish some basic definability properties that will be needed later.

Let G be a countable group. Recall that the **profinite topology** on G is the one with the basis of neighborhoods at 1_G consisting of all finite index subgroups (or, equivalently, all finite index normal subgroups); G is called **residually finite** if $\{1_G\}$ is closed in the profinite topology, or, equivalently, if the profinite topology is Hausdorff. For the rest of the paper, G will always be a residually finite group.

The **profinite completion** of G , denoted by \widehat{G} , is the completion of the group uniformly defined by this topology; equivalently $\widehat{G} = \varprojlim G/H$, where the limit is taken over all finite index normal subgroups of G . This profinite completion is metrizable if G has only countably many subgroups of finite index (for example, if it is finitely generated); even though it is not strictly necessary

for what we are doing (as we can always pass to suitable metrizable quotients), we will sometimes assume this for convenience.

Recall that the **left shift action** $G \curvearrowright 2^G$ is defined by $(g \cdot x)(h) = x(g^{-1}h)$. A closed subset $S \subseteq 2^G$ is called a **subshift** if it is invariant under this action.

Definition 2.1 (Krieger [18]): A word $x \in 2^G$ is called **Toeplitz** if $x: G \rightarrow 2$ is a continuous map for the profinite topology on G (where $2 = \{0, 1\}$ is taken to be discrete). A subshift $S \subseteq 2^G$ is **Toeplitz** if there exists a Toeplitz word x such that $S = \overline{G \cdot x}$.

Generalizing a well-known fact for $G = \mathbb{Z}$, Krieger showed that every Toeplitz subshift is minimal (see [18, Corollaire 2.5]).

Let

$$S(G) = \{S \subseteq 2^G : S \text{ is closed and } G\text{-invariant}\}$$

be the set of all G -subshifts. This is naturally a compact space with the Vietoris topology. It is easy to check that the family of minimal subshifts as well as that of free subshifts form Borel subsets of $S(G)$. In what follows, we see that being Toeplitz is also a Borel condition and establish a simple selection lemma that will be used later.

We need the following well-known definability property of Baire category notions for which we have not been able to find a suitable reference. It is a slight generalization of [16, 16.1] with the same proof. If X is a Polish space, $F(X)$ denotes the Effros Borel space of closed subsets of X . Recall that $\exists^* x \in F$ means “for non-meagerly many x in F .”

PROPOSITION 2.2: (i) *Let (X, \mathcal{S}) be a measurable space, Y a Polish space and let $\Phi: X \rightarrow F(Y)$ be a measurable function, where $F(Y)$ denotes the Effros Borel space of closed subsets of Y . Let $A \subseteq X \times Y$ be a measurable set (where Y is equipped with its Borel σ -algebra) and $U \subseteq Y$ an open set. Then the set*

(1) $\{x \in X : U \cap \Phi(x) = \emptyset \text{ or } A_x \cap \Phi(x) \text{ is non-meager in } U \cap \Phi(x)\}$
is measurable.

(ii) *Let X be a Polish space and $A \subseteq X$ be Borel. Then the set*

$$\{F \in F(X) : \exists^* x \in F \ x \in A\}$$

is Borel.

Proof. (i) The proof goes exactly as in [16, 16.1]. If we let A_U denote the set defined in (1), one checks that

- for all $S \in \mathcal{S}$, $V \subseteq Y$ open

$$(S \times V)_U = \{x \in X : U \cap \Phi(x) = \emptyset \text{ or } (x \in S \text{ and } U \cap V \cap \Phi(x) \neq \emptyset)\};$$

- $(\bigcup_n A_n)_U = \bigcup_n (A_n)_U$;
- $(\sim A)_U = \sim \bigcap_{U_n \subseteq U} A_{U_n}$, where the intersection is over all $U_n \subseteq U$ from a fixed countable base of Y .

(ii) This follows from (i). \blacksquare

If x is a Toeplitz word and $H \leq G$ is a finite index subgroup, we let

$$\text{Per}_H(x) = \{g \in G : x|_{Hg} \text{ is constant}\}.$$

The fact that x is Toeplitz translates to $\bigcup_H \text{Per}_H(x) = G$; H is called an **essential group of periods** for x if for all $g \notin H$,

$$\text{Per}_H(x) \not\subseteq \text{Per}_H(g \cdot x).$$

The **maximal equicontinuous factor** (**m.e.f.** for short) of a topological dynamical system $G \curvearrowright X$ is the factor generated by all equicontinuous factors of X . For a Toeplitz subshift S , this is always of the form $G \curvearrowright \varprojlim G/H_n$, where $\{H_n\}$ is a decreasing sequence of essential groups of periods (see [3, Proposition 7]). Note that every such system can also be written as $G \curvearrowright \widehat{G}/K$, where $K = \bigcap_n \overline{H_n}$ with the closures taken in \widehat{G} . We have the following folklore lemma.

LEMMA 2.3: *Let S be a Toeplitz subshift. Then the set of Toeplitz words in S is dense G_δ .*

Proof. Let Y be the m.e.f. of S and $\pi: S \rightarrow Y$ be the factor map. By [3, Theorem 2], the set of Toeplitz words in S can be written as

$$\{x \in S : \pi^{-1}(\{\pi(x)\}) = \{x\}\}.$$

This can be written as $\pi^{-1}(A)$, where

$$A = \bigcap_{\epsilon > 0} \bigcup \{U \subseteq Y \text{ open} : \text{diam}(\pi^{-1}(U)) < \epsilon\},$$

and this is clearly G_δ . It is also dense by the definition of a Toeplitz subshift. \blacksquare

Denote

$$\text{Töp}(G) = \{S \in S(G) : S \text{ is a Toeplitz subshift}\}.$$

Lemma 2.3 allows us to build the following selector map that will be used throughout the paper.

PROPOSITION 2.4: *Let G be a residually finite group. Then there exists a Borel map $\tau: \text{Töp}(G) \rightarrow 2^G$ such that for all $S \in \text{Töp}(G)$, $\tau(S) \in S$ and $\tau(S)$ is a Toeplitz word.*

Proof. This follows from Lemma 2.3, Proposition 2.2, and [16, 18.6]. ■

Now we can easily deduce the following.

LEMMA 2.5: *Let G be a residually finite group such that \widehat{G} is metrizable. The map $\text{Töp}(G) \rightarrow F(\widehat{G})$, which associates to a Toeplitz subshift S (the conjugacy class) of the subgroup $K \leq \widehat{G}$ such that the m.e.f. of S is isomorphic to \widehat{G}/K , is Borel.*

Proof. By Proposition 2.4, for a given Toeplitz subshift S , we can choose in a Borel way a Toeplitz word $x \in S$. By [3], we can choose in a Borel way a sequence $L_n(x)$ of finite index subgroups of G , which are essential periods of x and such that, posing $K = \bigcap_n \overline{L_n}$, we have that the m.e.f. of S is isomorphic to \widehat{G}/K . Now it is easy to check that the map that associates to the sequence $\{L_n\}_n$ the intersection $\bigcap_n \overline{L_n}$ is Borel. ■

PROPOSITION 2.6: *Let G be a residually finite group, \widehat{G} the profinite completion of G and $K \leq \widehat{G}$ a closed subgroup. Then the set $\text{Töp}(G)$ is Borel and, if \widehat{G} is metrizable,*

$$\text{Töp}(G, K) = \{S \in K(2^G) : S \text{ is a Toeplitz subshift with m.e.f. } \widehat{G}/K\}$$

is also Borel.

Proof. We have

$$S \in \text{Töp}(G) \iff \exists^* x \in S \text{ } x \text{ is Toeplitz}$$

and applying Proposition 2.2 yields that $\text{Töp}(G)$ is a Borel set.

The second statement follows from Lemma 2.5. ■

3. \mathbb{Z} -subshifts

In this section, we consider \mathbb{Z} -subshifts and prove Theorem 1.1.

Let $x \in 2^\mathbb{Z}$ be a Toeplitz word and $p \in \mathbb{N}$. Denote by $\text{Per}_p(x)$ the subset of $\mathbb{Z}/p\mathbb{Z}$ defined as follows:

$$\text{Per}_p(x) = \{i + p\mathbb{Z} : x(i + kp) = x(i) \text{ for all } k \in \mathbb{Z}\}.$$

If x is a Toeplitz word, p is an **essential period** for x if $\text{Per}_q(x) \subsetneq \text{Per}_p(x)$ for all $q|p$. Note that if x and y are Toeplitz words that belong to the same subshift, then they have the same essential periods. If x is Toeplitz, we write

$$\mathcal{P}(x) = \mathcal{P}(\overline{\mathbb{Z} \cdot x}) = \{p \in \mathbb{N} : \exists q \ p|q \text{ and } q \text{ is an essential period for } x\}.$$

As for a Toeplitz subshift S , $\mathcal{P}(S)$ encodes the m.e.f. of S , we have that if S and T are isomorphic, then $\mathcal{P}(S) = \mathcal{P}(T)$ (see, e.g., [26]). Also, $(\mathcal{P}(x), |)$ is a directed partial order (this is because if p and q are essential periods for x , then their least common multiple is one too).

Let $H_p(x) = (\mathbb{Z}/p\mathbb{Z}) \setminus \text{Per}_p(x)$. The elements of $H_p(x)$ are called **p -holes** for x . We say that x has **separated holes** [8] if

$$\lim_{p \rightarrow \infty} \min\{|i - j| : i, j \in H_p(x), i \neq j\} = \infty,$$

where the limit is taken over any sequence cofinal in $(\mathcal{P}(x), |)$.

Having separated holes is a property of the subshift that does not depend on the choice of the word x . Thus, by the arguments in Section 2, the condition of having separated holes defines a Borel subset of the set of all subshifts. All subshifts with separated holes are regular and have topological entropy 0.

Next we recall the definition of an **amenable** equivalence relation. Let E be a countable equivalence relation on the standard Borel space X . If $x \in X$ and $f: E \rightarrow \mathbb{R}$ is a function, denote by f_x the function $[x]_E \rightarrow \mathbb{R}$ defined by $f_x(y) = f(x, y)$; E is called **(1-)amenable** if there exist positive Borel functions $\lambda^n: E \rightarrow \mathbb{R}$ such that

- $\lambda_x^n \in \ell^1([x]_E)$, $\|\lambda_x^n\|_1 = 1$,
- $\lim_{n \rightarrow \infty} \|\lambda_x^n - \lambda_y^n\|_1 = 0$ for all $(x, y) \in E$.

By a theorem of Connes, Feldman and Weiss [2] (see also [17]), if μ is any probability measure on X and E is amenable, then it is hyperfinite μ -almost everywhere. It is an open question whether every amenable equivalence relation is hyperfinite. See [13] for more details on amenable equivalence relations in the Borel setting.

The goal of this section is to prove the following theorem.

THEOREM 3.1: *The equivalence relation of isomorphism of Toeplitz \mathbb{Z} -subshifts with separated holes is amenable.*

Proof. Let $p \in \mathbb{N}$. Denote by $\text{Sym}(2^p)$ the set of all bijections $2^p \rightarrow 2^p$. For $\pi \in \text{Sym}(2^p)$, let $\hat{\pi}: 2^{\mathbb{Z}} \rightarrow 2^{\mathbb{Z}}$ be defined by

$$\hat{\pi}(x)|_{[kp, (k+1)p)} = \pi(x|_{[kp, (k+1)p)})$$

for all $k \in \mathbb{Z}$. Then if $S \subseteq 2^{\mathbb{Z}}$ is a closed $p\mathbb{Z}$ -invariant set, $\hat{\pi}(S)$ is also a closed $p\mathbb{Z}$ -invariant set and $\hat{\pi}$ defines an isomorphism between them (as $p\mathbb{Z}$ -systems).

If $x \in 2^{\mathbb{Z}}$ and $i \in \mathbb{Z}$, denote by $x + i$ the shift of x by i . If x is a Toeplitz word and $p \in \mathcal{P}(x)$, define $A_{p,i}(x) \subseteq \text{Töp}(\mathbb{Z})$ by

$$A_{p,i}(x) = \{\overline{\mathbb{Z} \cdot \hat{\pi}(x + i)} : \pi \in \text{Sym}(2^p)\}$$

and note that $A_{p,i}(x)$ is a finite set. If $p|q$, then $A_{p,i}(x) \subseteq A_{q,i}(x)$. Denote by E the equivalence relation of isomorphism on $\text{Töp}(\mathbb{Z})$. By [6, Theorem 1], we have that

$$\bigcup_{\substack{p \in \mathcal{P}(x) \\ i < p}} A_{p,i}(x) = \{T \in \text{Töp}(\mathbb{Z}) : T \in E S\}$$

where $S = \overline{\mathbb{Z} \cdot x}$. Next, for $(S, T) \in E$ and $p \in \mathcal{P}(S) = \mathcal{P}(T)$ define

$$B^p(S, T) = \sum_{i < p} \chi_{A_{p,i}(x)}(T),$$

where χ_A denotes the characteristic function of A , and note that the value of $B^p(S, T)$ does not depend on the choice of a Toeplitz $x \in S$. Indeed, let $x_1, x_2 \in S$ be Toeplitz. We have that

$$S = \bigcup_{j < p} \overline{p\mathbb{Z} \cdot (x_1 + j)}$$

and there exists j_0 such that $x_2 \in \overline{p\mathbb{Z} \cdot (x_1 + j_0)}$ and therefore (as $\overline{p\mathbb{Z} \cdot (x_1 + j_0)}$ is Toeplitz and thus minimal) $\overline{p\mathbb{Z} \cdot x_2} = \overline{p\mathbb{Z} \cdot (x_1 + j_0)}$. Then $\overline{p\mathbb{Z} \cdot \hat{\pi}(x_2 + i)} = \overline{p\mathbb{Z} \cdot \hat{\pi}(x_1 + j_0 + i)}$, whence, by minimality,

$$\overline{\mathbb{Z} \cdot \hat{\pi}(x_2 + i)} = \overline{\mathbb{Z} \cdot \hat{\pi}(x_1 + j_0 + i)}$$

for every i and every π . Note also that, as we can choose $x = \tau(S)$ as per Proposition 2.4, the function B^p is Borel (and is defined on a Borel subset of E).

We have (for fixed $S \in \text{Töp}(\mathbb{Z})$, Toeplitz word $x \in S$, and $p \in \mathcal{P}(S)$)

$$\begin{aligned}\|B_S^p\|_1 &= \sum_{T \in S} B^p(S, T) \\ &= \sum_{i < p} \sum_{T \in S} \chi_{A_{p,i}(x)}(T) \\ &= \sum_{i < p} |A_{p,i}(x)|.\end{aligned}$$

In particular, $\|B_S^p\|_1$ is finite because all sets $A_{p,i}$ are finite. The functions B^p , appropriately normalized, will serve as our witnesses for amenability.

Recall that if S and T are subshifts and $f: S \rightarrow T$ is an isomorphism, then it is given by a **block code**, i.e., there exists $r \in \mathbb{N}$ and a function $\phi: 2^{2r+1} \rightarrow 2$ such that for all $x \in S$,

$$f(x)(i) = \phi(x|_{[i-r, i+r]}).$$

The number r is called the **radius** of the block code.

Let S and T be two Toeplitz subshifts with separated holes and let $f: S \rightarrow T$ be an isomorphism such that f and f^{-1} are given by block codes of radius r . Let $\epsilon > 0$ be given.

Let $x \in S$ be a Toeplitz word and let $y = f(x)$. Let $p \in \mathcal{P}(S)$ be so big that the distance between two consecutive holes in $H_p(x)$ and in $H_p(y)$ is larger than $M = (2r + 1)/\epsilon$. If $C \subseteq \mathbb{Z}$, denote

$$(C)_r = \{c + j \in \mathbb{Z}/p\mathbb{Z} : c \in C, |j| \leq r\}.$$

(Here and below, we suppress the natural quotient map $\mathbb{Z} \rightarrow \mathbb{Z}/p\mathbb{Z}$ from the notation.) Note that $|(C)_r| \leq (2r + 1)|C|$.

CLAIM 1: *For all $i \notin (H_p(x))_r$, we have $A_{p,i}(x) = A_{p,i}(y)$.*

Proof. This follows from the fact that for $i \notin (H_p(x))_r$, there exists $\pi \in \text{Sym}(2^p)$ such that $\hat{\pi}(x+i) = y+i$. Indeed, as $y = f(x)$ and f is given by a block code of radius r , $y|_{[i+kp, i+(k+1)p]}$ only depends on $x|_{[i+kp-r, i+(k+1)p+r]}$ and

$$\begin{aligned}x|_{[i+kp-r, i+kp]} &= x|_{[i+(k+1)p-r, i+(k+1)p]}, \\ x|_{[i+(k+1)p, i+(k+1)p+r]} &= x|_{[i+kp, i+kp+r]},\end{aligned}$$

because $(i-r, i+r) \subseteq \text{Per}_p(x)$. Thus $y|_{[i+kp, i+(k+1)p]}$ can be calculated from $x|_{[i+kp, i+(k+1)p]}$ in a way independent of k which gives the claim. ■

CLAIM 2: *For all $i \notin H_p(x)$, we have $A_{p,i}(x) = A_{p,i+1}(x)$.*

Proof. Indeed, let $\sigma: 2^{\mathbb{Z}/p\mathbb{Z}} \rightarrow 2^{\mathbb{Z}/p\mathbb{Z}}$ be defined by $\sigma(z)(j) = z(j-1)$, and observe that as $i \in \text{Per}_p(x)$, $\hat{\sigma}(x+i) = x+i+1$. Then

$$\begin{aligned} A_{p,i}(x) &= \overline{\{\mathbb{Z} \cdot \hat{\pi}(x+i) : \pi \in \text{Sym}(2^p)\}} \\ &= \overline{\{\mathbb{Z} \cdot (\widehat{\pi\sigma})(x+i) : \pi \in \text{Sym}(2^p)\}} \\ &= A_{p,i+1}(x). \quad \blacksquare \end{aligned}$$

Let $h_0, h_1, h_2, \dots, h_J$ enumerate the holes in $H_p(x)$ in circular order (so that $J+1 = 0$). By the choice of p , $|h_j - h_{j+1}| \geq M$ for all j . If f is a real-valued function, denote by f^+ the function $\max(f, 0)$ and note that $\|f\|_1 = \|f^+\|_1 + \|(-f)^+\|_1$. We calculate

$$\begin{aligned} \|(B_S^p - B_T^p)^+\|_1 &= \left\| \left(\sum_{i < p} \chi_{A_{p,i}(x)} - \sum_{i < p} \chi_{A_{p,i}(y)} \right)^+ \right\|_1 \\ &\leq \left\| \sum_{j < J} \sum_{i=h_j-r}^{h_j+r} \chi_{A_{p,i}(x)} \right\|_1 \\ &\leq \left\| \sum_{j < J} \left(\sum_{i=h_j+1}^{h_j+r} \chi_{A_{p,i}(x)} + \sum_{i=h_{j+1}-r}^{h_{j+1}} \chi_{A_{p,i}(x)} \right) \right\|_1 \\ &\leq \sum_{j < J} ((2r+1)/M) \sum_{i=h_j+1}^{h_{j+1}} |A_{p,i}(x)| \\ &= ((2r+1)/M) \|B_S^p\|_1 \\ &\leq \epsilon \|B_S^p\|_1. \end{aligned}$$

For the inequality on the second line, we use Claim 1, and for the one on the fourth line, we use Claim 2. Similarly, $\|(B_T^p - B_S^p)^+\|_1 \leq \epsilon \|B_T^p\|_1$, whence

$$(2) \quad \|B_S^p - B_T^p\|_1 \leq \epsilon (\|B_S^p\|_1 + \|B_T^p\|_1).$$

We are now in position to define the functions $\lambda^n: E \rightarrow \mathbb{R}$ that will witness the amenability of E . Let $Z \subseteq 2^{\mathbb{N}}$ be the Borel set of subsets of \mathbb{N} directed under divisibility. Let $N: Z \rightarrow \mathbb{N}^{\mathbb{N}}$ be a Borel function that for every $A \in Z$ chooses a sequence \lvert -cofinal in A . Finally, define $\lambda^n: E \rightarrow \mathbb{R}$ by

$$\lambda^n(S, T) = \frac{B^{p_n}(S, T)}{\|B_S^{p_n}\|_1},$$

where $p_n = N(\mathcal{P}(S))(n)$. The functions λ^n clearly satisfy the first condition in the definition of amenability; next, we check that they satisfy the second. Fix $S, T \in \text{Töp}(\mathbb{Z})$ with separated holes such that $(S, T) \in E$ and $\epsilon > 0$. It follows from our previous argument that for n sufficiently large, (2) is satisfied with p replaced by p_n . For all those n , we have

$$\|\lambda_S^n - \lambda_T^n\|_1 = \sum_{R \in S} \left| \frac{B_S^{p_n}(S, R)}{\|B_S^{p_n}\|_1} - \frac{B_T^{p_n}(T, R)}{\|B_T^{p_n}\|_1} \right|.$$

Put

$$s_R = B_S^{p_n}(S, R), \quad t_R = B_T^{p_n}(T, R), \quad u = \|B_S^{p_n}\|_1 = \sum_R s_R, \quad \text{and} \quad v = \|B_T^{p_n}\|_1 = \sum_R t_R$$

and note that (2) implies that $|u - v| \leq \epsilon(u + v)$. We calculate

$$\begin{aligned} \sum_R \left| \frac{s_R}{u} - \frac{t_R}{v} \right| &\leq \sum_R \left| \frac{s_R}{u} - \frac{s_R}{(u + v)/2} \right| + \sum_R \left| \frac{s_R}{(u + v)/2} - \frac{t_R}{(u + v)/2} \right| \\ &\quad + \sum_R \left| \frac{t_R}{(u + v)/2} - \frac{t_R}{v} \right| \\ &\leq \frac{|v - u|}{v + u} + \frac{2}{u + v} \|B_S^{p_n} - B_T^{p_n}\|_1 + \frac{|u - v|}{v + u} \\ &\leq 4\epsilon, \end{aligned}$$

thus showing that $\lim_{n \rightarrow \infty} \|\lambda_S^n - \lambda_T^n\|_1 = 0$. \blacksquare

4. The non-amenable case

Now we return to the general situation of Section 2, where G is a residually finite group. We also fix a decreasing sequence $\{H_n\}$ of finite index, normal subgroups with trivial intersection, to be determined later. We also denote by \hat{G} the inverse limit $\varprojlim G/H_n$ (which is a group quotient of the profinite completion \widehat{G}) and by $\pi_n: \hat{G} \rightarrow G/H_n$ the natural projections.

The goal of this section is to prove Theorem 1.2. The proof proceeds by constructing a probability measure on the set of Toeplitz subshifts which is invariant under an appropriate action of G contained in the isomorphism relation. We then check that the point stabilizers for this action are amenable and conclude that if G is non-amenable, then the isomorphism equivalence relation is not amenable either.

4.1. THE LEFT AND RIGHT ACTIONS. First we describe a method to build Toeplitz subshifts that will help us to construct the measure. Let

$$A_n = (H_{n-1}/H_n) \setminus \{H_n\}$$

and let

$$Z = 2^{\bigsqcup_n A_n}.$$

Let

$$Y = \{y \in \hat{G} : y \notin G\}.$$

Consider the maps

$$Y \times Z \xrightarrow{\sigma} 2^G \xrightarrow{\rho} S(G)$$

defined as follows. If $(y, z) \in Y \times Z$, define $\sigma(y, z)$ by

$$\sigma(y, z)(h) = z(\pi_{n_0}(y^{-1}h)), \quad \text{where } n_0 = \min\{n : \pi_n(y) \neq \pi_n(h)\}.$$

Define $\rho : 2^G \rightarrow S(G)$ by $\rho(x) = \overline{G \cdot x}$. Let $\theta = \rho \circ \sigma$.

Now G acts on 2^G on both sides: on the left by

$$(g \cdot x)(h) = x(g^{-1}h),$$

and on the right by

$$(x \cdot g)(h) = x(hg^{-1}),$$

and the two actions commute. Note that if $S \subseteq 2^G$ is a subshift and $g \in G$, then $S \cdot g$ is a subshift too, so we have a right action $S(G) \curvearrowright G$. Moreover, for every fixed $g \in G$, the map

$$S \rightarrow S \cdot g, \quad x \mapsto x \cdot g$$

is an isomorphism of subshifts.

The space $Y \times Z$ is also equipped with commuting left and right actions of G , defined as follows. For $y \in Y$ and $g \in G$, define $g \cdot y = gy$ and $y \cdot g = yg$ (recall that $G \subseteq \hat{G}$). For $z \in Z$, define $g \cdot z = z$ and $(z \cdot g)(a) = z(gag^{-1})$ for all $a \in \bigsqcup_n A_n$. Equip $Y \times Z$ with the diagonal left and right actions.

LEMMA 4.1: *The maps σ and ρ commute with both the left and right actions of G on the respective spaces.*

Proof. We only check that σ commutes with the right actions. Suppose that $(y_1, z_1), (y_2, z_2)$ are elements of $Y \times Z$ such that $(y_1, z_1) \cdot g = (y_2, z_2)$. Let $x_1 = \sigma(y_1, z_1)$ and $x_2 = \sigma(y_2, z_2)$. Note that for every $h \in G$ we have

$$\min\{n : \pi_n(y_1) \neq \pi_n(hg^{-1})\} = \min\{n : \pi_n(y_1g) \neq \pi_n(h)\}.$$

Write $n_0(h)$ for this common value. Thus

$$(x_1 \cdot g)(h) = x_1(hg^{-1}) = z_1(\pi_{n_0(h)}(y_1^{-1}hg^{-1})),$$

and

$$x_2(h) = z_2(\pi_{n_0(h)}(y_2^{-1}h)) = z_2(\pi_{n_0(h)}(g^{-1}y_1^{-1}h)).$$

But

$$\begin{aligned} z_2(\pi_{n_0(h)}(g^{-1}y_1^{-1}h)) &= z_1(g\pi_{n_0(h)}(g^{-1}y_1^{-1}h)g^{-1}) \\ &= z_1(\pi_{n_0(h)}(y_1^{-1}hg^{-1})), \end{aligned}$$

so $x_1 \cdot g = x_2$, as needed. \blacksquare

We call an element $z \in Z$ **proper** if it takes both values 0 and 1 infinitely many times. Note that z is proper if and only if for some (equivalently, every) $y \in Y$, $G \cdot \sigma(y, z)$ is infinite. For the if direction, note that for z which is constant on A_n for $n > n_0$, $\sigma(y, z)$ is stabilized by H_{n_0} for every $y \in Y$. The only if implication follows from the following lemma.

LEMMA 4.2: *For any $y \in Y$ and proper $z \in Z$, $\sigma(y, z)$ is a Toeplitz word and the m.e.f. of $\overline{G \cdot \sigma(y, z)}$ is isomorphic to \hat{G} . In particular, the subshift $\overline{G \cdot \sigma(y, z)}$ is free.*

Proof. Write $x = \sigma(y, z)$. First note that for any $g \in G$, the value $x(g)$ is assumed on the whole H_{n_0} -coset $\pi_{n_0}(y^{-1}g)$, where

$$n_0 = \min\{n : \pi_n(y) \neq \pi_n(g)\},$$

showing that x is Toeplitz.

To calculate the m.e.f., by [3], it suffices to observe that for all n , H_n is an essential group of periods for x . Indeed, using the fact that z is proper, we have that $\text{Per}_{H_n}(x) = G \setminus \pi_n(y)$ which is not contained in $\text{Per}_{H_n}(g \cdot x) = G \setminus \pi_n(g \cdot y)$ for any $g \notin H_n$.

For the final claim, just note that as $\bigcap_n H_n = \{1_G\}$ and the H_n are normal, the translation action $G \curvearrowright \hat{G}$ is free. \blacksquare

The main ingredient for the proof of the theorem is the following lemma.

LEMMA 4.3: *For all proper $z_1, z_2 \in Z$ and all $y_1, y_2 \in Y$,*

$$z_1 \neq z_2 \implies \theta(y_1, z_1) \neq \theta(y_2, z_2).$$

Proof. Let $x_i = \sigma(y_i, z_i)$. To show that $\overline{G \cdot x_1} \neq \overline{G \cdot x_2}$, it suffices to find a “subword” of x_2 that does not occur in x_1 , i.e., a finite set $F \subseteq G$ such that

$$\forall g \in G \exists f \in F \quad x_1(gf) \neq x_2(f).$$

Let n and $a_0 \in A_n$ be such that $z_1(a_0) \neq z_2(a_0)$. By replacing (y_1, z_1) and (y_2, z_2) with $g_1 \cdot (y_1, z_1)$ and $g_2 \cdot (y_2, z_2)$ (which does not change either z_i or $\theta(y_i, z_i)$) for suitably chosen $g_1, g_2 \in G$, we may assume that

$$\pi_n(y_1) = \pi_n(y_2) = 1_{G/H_n}.$$

Let $f_0, f_1 \in G$ be such that $\pi_n(f_0) = \pi_n(f_1)$, $x_2(f_0) = 0$, $x_2(f_1) = 1$ (such f_0, f_1 exist by the assumption that z_2 is proper). Let $f \in G$ be such that $\pi_n(f) = a_0$ and, finally, let $F = \{f, f_0, f_1\}$.

Next we show that this F works. Let $g \in G$ be arbitrary. We distinguish two cases: $g \in H_n$ and $g \notin H_n$. Let first $g \in H_n$. We check that $x_1(gf) \neq x_2(f)$. Indeed,

$$x_1(gf) = z_1(\pi_n(y_1^{-1}gf)) = z_1(\pi_n(f)) = z_1(a_0)$$

and

$$x_2(f) = z_2(\pi_n(y_2^{-1}f)) = z_2(\pi_n(f)) = z_2(a_0),$$

which are different by the choice of a_0 .

Suppose now that $g \notin H_n$. We will show that $x_1(gf_0) = x_1(gf_1)$ which will complete the proof (as $x_2(f_0) \neq x_2(f_1)$). Indeed, observe that the least k for which $\pi_k(gf_i) \neq 1_{G/H_k}$ is at most n and is the same for $i = 0, 1$ (as $\pi_n(f_0) = \pi_n(f_1)$). Recall also that $\pi_k(y_1) = \pi_k(y_2) = 1_{G/H_k}$. Now we have

$$\begin{aligned} x_1(gf_0) &= z_1(\pi_k(y_1^{-1}gf_0)) = z_1(\pi_k(gf_0)) \\ &= z_1(\pi_k(gf_1)) = x_1(gf_1), \end{aligned}$$

completing the proof. ■

4.2. A.E. AMENABLE STABILIZERS. Let λ be the Haar measure on \hat{G} and note that as $\lambda(Y) = 1$, we can consider λ as a measure on Y . Let ν be the Bernoulli $(\frac{1}{2}, \frac{1}{2})$ measure on $Z = 2\sqcup_n A_n$. Equip $Y \times Z$ with the product measure $\lambda \times \nu$. It is clear that this measure is invariant under both the left and the right action of G (but for us it will be the right one that will be important). Let $\mu = \theta_*(\lambda \times \nu)$.

LEMMA 4.4: *The measure μ concentrates on the set of free, Toeplitz G -subshifts and is invariant under the right action.*

Proof. This follows from Lemma 4.2 and the fact that ν concentrates on the set of proper elements of Z . Invariance follows directly from Lemma 4.1. ■

For $g \in G$, denote by $C(g)$ the **centralizer** of g in G , i.e., the set of all elements of G that commute with g . Let

$$Z_f(G) = \{g \in G : C(g) \text{ has finite index in } G\}.$$

LEMMA 4.5: $Z_f(G)$ is an amenable, normal subgroup of G .

Proof. It is clear that $Z_f(G)$ is a normal subgroup of G . To see that it is amenable, note that the centralizers of all elements in $Z_f(G)$ have finite index in $Z_f(G)$. Hence, $Z_f(G)$ has finite conjugacy classes and this implies amenability (Leptin [19]). ■

Next we choose a suitable sequence $\{H_n\}$ of finite index, normal subgroups of G that will allow us to prove the theorem. Note that the residual finiteness of G implies that for any finite index subgroup $H \triangleleft G$ and $g \in G \setminus Z_f(G)$ there exists a finite index subgroup $H' \leq H$ normal in G such that for some $C \in H/H'$ we have $gCg^{-1} \neq C$. Enumerate $G \setminus Z_f(G)$ as $\{g_n : n \in \mathbb{N}\}$ and construct inductively $\{H_n : n \in \mathbb{N}\}$ a decreasing sequence with trivial intersection such that

(3) for all $m \geq n$ there exists $C \in H_m/H_{m+1}$ such that $g_n^{-1}Cg_n \neq C$.

Proof of Theorem 1.2. We will show that the right action of G on $\text{Töp}(G)$ has μ -a.e amenable stabilizers. Recall that (see, e.g., [11, Lemma 3.6]) if a measure-preserving action of a group has amenable stabilizers and induces a hyperfinite equivalence relation, then the group is amenable. Thus, if G is non-amenable, we obtain that the orbit equivalence relation induced by the right action of G on the set of Toeplitz G -subshifts is not hyperfinite and, as it is contained in the isomorphism relation, the latter is not hyperfinite either.

By Lemma 4.5, it is enough to show that there is a measure 1 subset A of $\text{Töp}(G)$ such that the stabilizer of every element in A is contained in $Z_f(G)$. Since G is countable, it suffices to see that for every $g \in G \setminus Z_f(G)$, the set

$$\{(y, z) \in Y \times Z : \theta(y, z) \cdot g \neq \theta(y, z)\}$$

has measure 1. By Lemma 4.3, it is enough to show that $\{z \in Z : z \cdot g \neq z\}$ is of measure 1. As $g \notin Z_f(G)$, by (3), for almost all n , there is an element, say $C_n \in H_{n+1}/H_n$, such that $gC_ng^{-1} \neq C_n$. By the definition of the action of G

on Z , for any $z \in Z$ such that there exists n with $z(C_n) \neq z(gC_n g^{-1})$, we have $z \neq z \cdot g$. But the latter condition is clearly satisfied on a measure 1 set. This completes the proof of the theorem. ■

Remark 4.6: Note that in case G is a non-cyclic free group, the right action of G on $\text{Töp}(G)$ is free μ -a.e. because $Z_f(G)$ is trivial (centralizers of non-trivial elements of free groups are cyclic). This implies in particular that the equivalence relation of isomorphism of Toeplitz G -subshifts embeds a free, measure-preserving action of a free group.

5. The groupoid viewpoint

The equivalence relation of isomorphism of subshifts is naturally given by an action of a groupoid rather than a group. Recall that a **groupoid** is a small category where each arrow is invertible. We will only be interested in countable Borel groupoids, defined as follows (see also [21]). A **countable Borel groupoid** Γ is a tuple (X, A, s, r, \circ) , where X is a standard Borel space of objects, A is a standard Borel space of arrows, $s: A \rightarrow X$ is a Borel map specifying the **source** of each arrow and $r: A \rightarrow X$ a Borel map specifying its **range**; \circ is a partial Borel map $A \times A \rightarrow A$ which represents the composition of arrows; $f \circ g$ is defined whenever $r(g) = s(f)$. Of course, we require that (X, A, \circ) be a groupoid. Furthermore, we require that $r^{-1}(\{x\})$ and $s^{-1}(\{x\})$ be countable sets for every $x \in X$. By the Lusin–Novikov selection theorem, there exist sequences of Borel maps $P_n: X \rightarrow A$ and $Q_n: X \rightarrow A$ such that $P_n(x)$ enumerate $s^{-1}(x)$ and $Q_n(x)$ enumerate $r^{-1}(x)$ for every x . If $x, y \in X$, we will denote by $A(x, y)$ the set of arrows between x and y . We will sometimes identify a groupoid with the set of its arrows as the other information can be recovered from it.

Every countable Borel groupoid Γ gives rise to a countable Borel equivalence relation E_Γ on X defined by

$$x E_\Gamma y \iff \exists f \in A \ s(f) = x \text{ and } r(f) = y.$$

Conversely, every countable equivalence relation can be viewed as a groupoid, where there is a unique arrow between every pair of equivalent elements of X .

Now let G be a residually finite group as before, $\{H_n\}$ be a sequence of finite index, normal subgroups of G , and let X be the standard Borel space of all Toeplitz subshifts of G with m.e.f. $\hat{G} = \varprojlim G/H_n$. Let A be the set of all

isomorphisms between elements of X . Recalling that every such isomorphism is given by a block code (which is a finite object), it is easy to endow A with a standard Borel structure so that (X, A, \circ) becomes a countable Borel groupoid (\circ is just composition of maps). The equivalence relation generated by this groupoid is exactly isomorphism of subshifts. (All of this is defined for arbitrary subshifts of countable groups; however, below we will need that they be Toeplitz.)

Note that for every $S \in X$ and every $x \in S$, there exists a unique factor map $\pi_x: S \rightarrow \hat{G}$ such that $\pi(x) = 1_{\hat{G}}$. (Existence follows from the fact that if $\pi: S \rightarrow \hat{G}$ is any G -map, then post-composing π with right multiplication by $\pi(x)^{-1}$ yields a map that sends x to $1_{\hat{G}}$.) Let $\tau: X \rightarrow 2^G$ be a Borel map that selects, for every $S \in X$, a Toeplitz word $\tau(S) \in S$ as given by Proposition 2.4. Now we can define a Borel cocycle $\alpha_0: \Gamma \rightarrow \hat{G}$ by

$$\alpha_0(f) = \pi_{\tau(r(f))}(f(\tau(s(f))))^{-1}.$$

Recall that a **cocycle** is just a map $\Gamma \rightarrow \hat{G}$ that satisfies $\alpha(f \circ g) = \alpha(f)\alpha(g)$. Two cocycles α, β are **cohomologous** if there exists a Borel map $F: X \rightarrow \hat{G}$ such that $\beta(f) = F(r(f))^{-1}\alpha(f)F(s(f))$ for all $f \in \Gamma$. Note that changing the selection map τ transforms α_0 into a cohomologous cocycle. We also have a natural Borel homomorphism $\rho: \Gamma \rightarrow E$ defined by

$$\rho(f) = (s(f), r(f)).$$

Now we specialize to the case $G = \mathbb{Z}$. We have the following proposition, which basically follows from a result by Downarowicz, Kwiatkowski and Lacroix [6].

PROPOSITION 5.1: *The equivalence relation E on Toeplitz words in $2^{\mathbb{Z}}$ defined by*

$x E y \iff$ there exists an isomorphism $f: \overline{\mathbb{Z} \cdot x} \rightarrow \overline{\mathbb{Z} \cdot y}$ such that $f(x) = y$
is hyperfinite.

Proof. Define the equivalence relation E_m on Toeplitz words in $2^{\mathbb{Z}}$ by

$$x E_m y \iff x E y \text{ and } \exists \sigma \in \text{Sym}(2^m) \ \hat{\sigma}(x) = y.$$

Each E_m is a finite equivalence relation, $E_m \subseteq E_n$ if $m|n$, and by the first paragraph of the proof of [6, Theorem 1], $E = \bigcup_n E_{n!}$. This shows that E is hyperfinite. ■

COROLLARY 5.2: *Let $\Delta = \ker \alpha_0 = \{f \in \Gamma : \alpha_0(f) = 0\}$. Then the equivalence relation E_Δ is hyperfinite.*

Proof. Let E be the equivalence relation on Toeplitz words as in Proposition 5.1. Then the map $\text{T}\ddot{\text{o}}\text{p}(\mathbb{Z}) \rightarrow 2^{\mathbb{Z}}$, $S \mapsto \tau(S)$ is a reduction from E_Δ to E thus showing that E_Δ is hyperfinite. (Here we use the fact that an isomorphism sends Toeplitz words to Toeplitz words and that for a Toeplitz S , the map $\pi_{\tau(S)}$ is injective on the set of Toeplitz sequences (see [6]).) ■

This corollary suggests that the groupoid Γ is not very complicated: it is an extension of a groupoid generating a hyperfinite equivalence relation by a compact group.

If S is a subshift, the **centralizer** $C(S)$ of S is the group of automorphisms of S , or equivalently, the group of arrows from S to itself. Note that by our observations above, if $S \in X$, then $C(S)$ embeds in \hat{G} and is therefore an abelian group. So, in some sense, the groupoid Γ differs from the equivalence relation E_Γ only a little.

We finally observe that the existence of the cocycle α_0 gives some restrictions on the groupoid Γ . For example, using Popa's cocycle superrigidity results [23], it is easy to prove that Γ does not embed the groupoid of the free part of any Bernoulli action of an infinite property (T) group. However, it is not clear how to conclude anything from that about the equivalence relation E ; in particular, we do not know whether E is universal.

References

- [1] J. D. Clemens, *Isomorphism of subshifts is a universal countable Borel equivalence relation*, Israel Journal of Mathematics **170** (2009), 113–123.
- [2] A. Connes, J. Feldman and B. Weiss, *An amenable equivalence relation is generated by a single transformation*, Ergodic Theory and Dynamical Systems **1** (1981), 431–450.
- [3] M. I. Cortez and S. Petite, *G -odometers and their almost one-to-one extensions*, Journal of the London Mathematical Society **78** (2008), 1–20.
- [4] M. I. Cortez and S. Petite, *Invariant measures and orbit equivalence for generalized Toeplitz subshifts*, Groups, Geometry, and Dynamics **8** (2014), 1007–1045.
- [5] R. Dougherty, S. Jackson and A. S. Kechris, *The structure of hyperfinite Borel equivalence relations*, Transactions of the American Mathematical Society **341** (1994), 193–225.
- [6] T. Downarowicz, J. Kwiatkowski and Y. Lacroix, *A criterion for Toeplitz flows to be topologically isomorphic and applications*, Colloquium Mathematicum **68** (1995), 219–228.

- [7] T. Downarowicz, *The Choquet simplex of invariant measures for minimal flows*, Israel Journal of Mathematics **74** (1991), 241–256.
- [8] T. Downarowicz, *Survey of odometers and Toeplitz flows*, in *Algebraic and Topological Dynamics*, Contemporary Mathematics, Vol. 385, American Mathematical Society, Providence, RI, 2005, pp. 7–37.
- [9] S. Gao, S. Jackson and B. Seward, *Group colorings and Bernoulli subflows*, Memoirs of the American Mathematical Society **241** (2016).
- [10] T. Giordano, I. F. Putnam and C. F. Skau, *Full groups of Cantor minimal systems*, Israel Journal of Mathematics **111** (1999), 285–320.
- [11] G. Hjorth, *Non-treeability for product group actions*, Israel Journal of Mathematics **163** (2008), 383–409.
- [12] G. Hjorth and A. S. Kechris, *Rigidity theorems for actions of product groups and countable Borel equivalence relations*, Memoirs of the American Mathematical Society **177** (2005).
- [13] S. Jackson, A. S. Kechris and A. Louveau, *Countable Borel equivalence relations*, Journal of Mathematical Logic **2** (2002), 1–80.
- [14] K. Juschenko and N. Monod, *Cantor systems, piecewise translations and simple amenable groups*, Annals of Mathematics **178** (2013), 775–787.
- [15] B. Kaya, *The complexity of the topological conjugacy problem for Toeplitz subshifts*, Israel Journal of Mathematics **220** (2017), 871–895..
- [16] A. S. Kechris, *Classical Descriptive Set Theory*, Graduate Texts in Mathematics, Vol. 156, Springer-Verlag, New York, 1995.
- [17] A. S. Kechris and B. D. Miller, *Topics in Orbit Equivalence*, Lecture Notes in Mathematics, Vol. 1852, Springer-Verlag, Berlin, 2004.
- [18] F. Krieger, *Sous-décalages de Toeplitz sur les groupes moyennables résiduellement finis*, Journal of the London Mathematical Society **75** (2007), 447–462.
- [19] H. Leptin, *Zur harmonischen Analyse klassenkomakter Gruppen*, Inventiones Mathematicae **5** (1968), 249–254.
- [20] D. Lind and B. Marcus, *An Introduction to Symbolic Dynamics and Coding*, Cambridge University Press, Cambridge, 1995.
- [21] M. Lupini, *Polish groupoids and functorial complexity*, Transactions of the American Mathematical Society (2017), to appear.
- [22] H. Matui, *Some remarks on topological full groups of Cantor minimal systems*, International Journal of Mathematics **17** (2006), 231–251.
- [23] S. Popa, *Cocycle and orbit equivalence superrigidity for malleable actions of w -rigid groups*, Inventiones Mathematicae **170** (2007), 243–295.
- [24] S. Thomas, *Topological full groups of minimal subshifts and just-infinite groups*, in *Proceedings of the 12th Asian Logic Conference*, World Scientific, Hackensack, NJ, 2012, pp. 298–313.
- [25] J. Williams, *Isomorphism of finitely generated solvable groups is weakly universal*, Journal of Pure and Applied Algebra **219** (2015), 1639–1644 .
- [26] S. Williams, *Toeplitz minimal flows which are not uniquely ergodic*, Zeitschrift für Wahrscheinlichkeitstheorie und Verwandte Gebiete **67** (1984), 95–107.