

Generalized p -angulations in higher dimension

Luca Lionni

Paris-Sud (LPT) & Paris 13 (LIPN)

Institut Camille Jordan– 10/03/2017

1 – Motivation and main ideas

2 – Colored triangulations and edge-colored graphs

3 – Generalized p -angulations

4 – Quadrangulations in 4D

5 – Gluings of octahedra

1 – Motivation and main ideas

1 – Motivation and main ideas

Motivation : quantum gravity

Einstein-Hilbert partition function for Euclidean pure gravity in dimension D

$$Z(\lambda, N) = \int_{\mathcal{M}} D[g] e^{-\int d^D x \sqrt{|g|} (2\Lambda - \frac{1}{2\kappa} R)} = \sum_{\substack{T \\ \text{connected} \\ \text{triangulation}}} \lambda^{n_D} N^{n_{D-2} - a n_D}$$

of D simplices

of D-2 simplices

1 – Motivation and main ideas

Motivation : quantum gravity

Einstein-Hilbert partition function for Euclidean pure gravity in dimension D

$$Z(\lambda, N) = \int_{\mathcal{M}} D[g] e^{-\int d^D x \sqrt{|g|} (2\Lambda - \frac{1}{2\kappa} R)} = \sum_{\substack{T \\ \text{connected} \\ \text{triangulation}}} \lambda^{n_D} N^{n_{D-2} - a n_D}$$

Allow topology fluctuations -> non-classical 

1 – Motivation and main ideas

Motivation : quantum gravity

Einstein-Hilbert partition function for Euclidean pure gravity in dimension D

$$Z(\lambda, N) = \int_{\mathcal{M}} D[g] e^{-\int d^D x \sqrt{|g|} (2\Lambda - \frac{1}{2\kappa} R)} = \sum_{\substack{T \\ \text{connected} \\ \text{triangulation}}} \lambda^{n_D} N^{n_{D-2} - a n_D}$$

Allow topology fluctuations \rightarrow non-classical 

- **Large N limit** (physical limit of small Newton constant) :

configurations which maximize 

- $\lambda \longrightarrow \lambda_c$: Continuum limit \rightarrow quantum space-time

1 – Motivation and main ideas

D=2 : continuum limit = Brownian map

Hausdorff dimension 4, homeomorphic to S^2 ,

Quantum sphere of Liouville quantum gravity (Miller, Sheffield, 2016)





Fig : J. Bettinelli

1 – Motivation and main ideas

D>2 : Basic idea

- Glue building blocks together

“Quanta of space-time”



D>2 : Main ideas

- Identify configurations which maximize n_{D-2} at fix n_D

D>2 : Main ideas

- Identify configurations which maximize n_{D-2} at fix n_D
- Gurau's theorem :
$$n_{D-2} \leq D + \frac{D(D-1)}{4}n_D$$

D>2 : Main ideas

- Identify configurations which maximize n_{D-2} at fix n_D
- Gurau's theorem : $n_{D-2} \leq D + \frac{D(D-1)}{4}n_D$
- In known cases, max. configurations verify $n_{D-2} = D + a n_D$

with $a \leq \frac{D(D-1)}{4}$ (" < " for interesting cases)

→ find the *coefficient a* ? what is their *topology* ?

D>2 : Main ideas

- Identify configurations which maximize n_{D-2} at fix n_D
- Gurau's theorem : $n_{D-2} \leq D + \frac{D(D-1)}{4} n_D$
- In known cases, max. configurations verify $n_{D-2} = D + a n_D$

with $a \leq \frac{D(D-1)}{4}$ (" < " for interesting cases)

→ find the *coefficient a* ? what is their *topology* ?

- Count maximal configurations : generating function has a *singularity*
→ continuum limit → space-time

→ *critical exponent* ? ... *Hausdorff dimension* ? *Fractal dimension* ? Etc.


2 – Colored triangulations and edge-colored graphs

2 – Colored triangulations and edge-colored graphs

Simplicial pseudo-complexes obtained by gluing D-simplices

Colored faces (D-1 simplices) are glued in a **unique way** :


with matching colors on their sub-simplices



2 – Colored triangulations and edge-colored graphs

D-simplices are represented by
(D+1)-valent vertices

The colored faces are dual to colored
edges




2 – Colored triangulations and edge-colored graphs

D-simplices are represented by
(D+1)-valent vertices


The colored faces are dual to colored
edges

Black vertex / white vertex : opposite
ordering of colors around faces




2 – Colored triangulations and edge-colored graphs

A color- i edge encodes the gluing of two color- i “faces” (D-1 simplices) in the *unique* possible way




2 – Colored triangulations and edge-colored graphs

3D triangulation
with boundary :




An octahedron,
or bipyramid...


2 – Colored triangulations and edge-colored graphs



2 – Colored triangulations and edge-colored graphs




2 – Colored triangulations and edge-colored graphs



2 – Colored triangulations and edge-colored graphs

Dictionary :


triangulation	\leftrightarrow	dual graph
D-simplex	\leftrightarrow	vertex
(D-1) simplex	\leftrightarrow	edge
(D-2) simplex	\leftrightarrow	two-colored cycle
(D-k) simplex	\leftrightarrow	sub-graph with k colors only



2 – Colored triangulations and edge-colored graphs

Dictionary :

triangulation	\leftrightarrow	dual graph
D-simplex	\leftrightarrow	vertex
(D-1) simplex	\leftrightarrow	edge
(D-2) simplex	\leftrightarrow	two-colored cycle
(D-k) simplex	\leftrightarrow	sub-graph with k colors only




Edges of the
triangulation

2 – Colored triangulations and edge-colored graphs

Dictionary :

triangulation	\leftrightarrow	dual graph
D-simplex	\leftrightarrow	vertex
(D-1) simplex	\leftrightarrow	edge
(D-2) simplex	\leftrightarrow	two-colored cycle
(D-k) simplex	\leftrightarrow	sub-graph with k colors only



Vertex of the
triangulation

2 – Colored triangulations and edge-colored graphs

We are interested in configurations with maximal number of $(D-2)$ simplices at fixed number of D -simplices.

- $D=2$: maximal # vertices, fixed # triangles
→ minimize the genus
- $D=3$: maximal # edges, fixed # tetrahedra

Dual picture : graphs that maximize the number of two-colored cycles at fixed number of vertices.

→ « *maximal graphs* »

2 – Colored triangulations and edge-colored graphs

Colored triangulations verify $n_{D-2} \leq D + \frac{D(D-1)}{4}n_D$

2 – Colored triangulations and edge-colored graphs

Colored triangulations verify $n_{D-2} \leq D + \frac{D(D-1)}{4} n_D$

Maximal triangulations : $D=2$


$$V = 2 + \frac{F}{2} \iff g = 0$$

→ planar triangulations

$$T(n) = \frac{1}{16} \sqrt{\frac{3}{2\pi}} n^{-\frac{5}{2}} \left(\frac{256}{27}\right)^n \propto n^{\gamma-2} \lambda_c^{-n}$$

$$\rightarrow \boxed{\gamma = -\frac{1}{2}}$$

Continuum limit = brownian map




2 – Colored triangulations and edge-colored graphs

Colored triangulations verify $n_{D-2} \leq D + \frac{D(D-1)}{4} n_D$

Maximal triangulations : $D > 2$

They are called *melonc graphs*



2 – Colored triangulations and edge-colored graphs

Colored triangulations verify $n_{D-2} \leq D + \frac{D(D-1)}{4} n_D$


Maximal triangulations : $D > 2$

They are called *melon graphs*

Tree-like structure

$$\rightarrow \boxed{\gamma = \frac{1}{2}}$$

Continuum limit = branched polymers
...not a good space-time candidate...



3 – Generalized p-angulations

3 – Generalized p-angulations


p-angulation in 2D

Maximize the number of vertices at fixed number of p-gons

$$n_{\text{vertices}} \leq 2 + \frac{p-2}{2} n_{p\text{-gons}}$$

→ Selects planar p-angulations, as before for triangulations


→ **Universality** (critical exponent, continuum limit...)



hexangulation, locally

3 – Generalized p-angulations

p-angulation in higher dimension




3 – Generalized p-angulations

Gluing of **building blocks** with p external faces of color 0 in dimension D

e.g. :


8-angulation in 3D



3 – Generalized p-angulations

Gluing of building blocks with p external faces of color 0 in dimension D

Internal colored structure



3 – Generalized p -angulations




Gluing of building blocks with p external faces of color 0 in dimension D

↳ D -colored triangulations with a
connected boundary of size p , and color 0

3 – Generalized p-angulations

Gluing of building blocks with p external faces of color 0 in dimension D

↳ D -colored triangulations with a connected boundary of size p , and color 0



3 – Generalized p-angulations




Gluing of building blocks with p external faces of color 0 in dimension D

↳ D -colored triangulations with a connected boundary of size p , and color 0




e.g. :

8-gons in 3D

(dual)



...



3 – Generalized p-angulations


Dual picture

An edge of color 0 (dashed) identifies two faces (D-1 simplices) of color 0

e.g. :

8-angulation in 3D

→ 4-colored graph




3 – Generalized p-angulations

N.B : building blocks made of D-simplices $\rightarrow n_{D-2} \leq D + \frac{D(D-1)}{4}n_D$

always true but **not saturated!!** and finite # gluings per order (Gurau-Schaeffer)

\rightarrow Find smallest a such that $n_{D-2} \leq D + an_D$



and = is saturated by infinite # of gluings




4 – Generalized quadrangulations in 4D

4 – Generalized quadrangulations in 4D

Building blocks






Quadrangulation




4 – Generalized quadrangulations in 4D

Building blocks





Quadrangulation




4 – Generalized quadrangulations in 4D

Building blocks





Quadrangulation




4 – Generalized quadrangulations in 4D

Building blocks





Cycles 0-Pairs





4 – Generalized quadrangulations in 4D

Building blocks




Quadrangulation



4 – Generalized quadrangulations in 4D


Bi-colored cycles are faces around one-colored sub-map




Maximize the sum of faces of one-colored sub-maps

- Trees behave as :


$$n_{D-2}(T) = 4 + \frac{5}{2}n_D^{34} + 3n_D^4$$




- Trees behave as : $n_{D-2}(T) = 4 + \frac{5}{2}n_D^{34} + 3n_D^4$




- Deleting an edge e : $n'_{D-2} = n_{D-2}(M) + 4 - 2I_2(e)$




- Trees behave as : $n_{D-2}(T) = 4 + \frac{5}{2}n_D^{34} + 3n_D^4$




- Deleting an edge e : $n'_{D-2} = n_{D-2}(M) + 4 - 2I_2(e)$


M maximal \Rightarrow  must be **bridges**

≤ 1 for $\left| \begin{array}{c} 4 \\ \approx \end{array} \right.$ 

- Trees behave as : $n_{D-2}(T) = 4 + \frac{5}{2}n_D^{34} + 3n_D^4$



- Deleting an edge e : $n'_{D-2} = n_{D-2}(M) + 4 - 2I_2(e)$





- Once they are : $n_{D-2}(M) = n_{D-2}(T) - 4g(M)$



4 – Generalized quadrangulations in 4D

The sharp bounds are

Gluing of  :
$$n_{D-2} \leq 4 + 3n_D$$
 Maximal config. are TREES

Gluing of  :
$$n_{D-2} \leq 4 + \frac{5}{2}n_D$$
 Maximal config. are PLANAR

Maximal gluings have the topology of the 4-sphere

Gluing of **both** :
$$n_{D-2} \leq 4 + \frac{5}{2}n_D^{34} + 3n_D^4$$

$$\left(3 = \frac{D(D-1)}{4} \right)$$

And maximal Configs : Planar, and $\left| \begin{array}{c} 4 \\ \vdots \end{array} \right.$ are bridges

Generating function :
$$F(t, \lambda) = \sum_{M \text{ max.}} t^{E(M)} \lambda^{E_4(M)}$$

$\lambda > 3$: $F \sim a_1(\lambda) + b_1(\lambda) \sqrt{t_1(\lambda) - t} + \dots$ Tree regime $\gamma = \frac{1}{2}$

$\lambda < 3$: $F \sim a_2(\lambda) + b_2(\lambda)(t_2(\lambda) - t) + c_2(\lambda)(t_2(\lambda) - t)^{3/2} + \dots$ Planar regime $\gamma = -\frac{1}{2}$

$\lambda = 3$: $F \sim \frac{16}{9} + \frac{128}{3^{5/3}} \left(\frac{3}{64} - t \right)^{2/3} + \dots$ $\gamma = \frac{1}{3}$ Proliferation of baby universes


4 – Generalized quadrangulations in 4D

- The critical behavior of maximal configurations
is NOT universal (unlike $D=2$)

4 – Generalized quadrangulations in 4D


These results can be extended to blocks of any size, in any even dimension :

“Necklaces”



+

“Melonic” graphs



+

(and their
connected
sums)

$$n_{D-2} \leq 4 + 2\left(1 + \frac{1}{p}\right)n_D$$

$$n_{D-2} \leq 4 + 3n_D$$

This is rather easy (only size 4 or connected sums)

Can we do bigger building blocks with new internal structure??

This is rather easy (only size 4 or connected sums)

Can we do bigger building blocks with new internal structure??


→ Yes ! But it is more difficult. First example : gluings of octahedra

N.B. tool = same kind of bijection


5 – Gluings of octahedra

5 – Gluings of octahedra

Building blocks



Gluings of octahedra



5 – Gluings of octahedra

Maximal triangulations verify

$$n_{\text{edges}} = 3 + 5n_{\text{octahedra}}$$

And 3D gluings of octahedra verify (w.r.t. their constituting tetrahedra)

$$n_{\text{edges}} \leq 2 + \frac{11}{8}n_{\text{tetrahedra}}$$



Compare with 3D gluings of melonic 8-gons $n_{\text{edges}} \leq 3 + \frac{3}{2}n_{\text{tetrahedra}}$

5 – Gluings of octahedra



Proofs also rely on a **bijection** (with “stuffed” hyper-maps)

5 – Gluings of octahedra

Building blocks





Bicolored cycles 03





5 – Gluings of octahedra

Building blocks



Quadrangulation



5 – Gluings of octahedra

Maximal triangulations are in bijection with a family of trees.

5 – Gluings of octahedra

The generating function of maximal maps with one marked corner is s.t.




$$G(z) = 1 + 3zG(z)^4 \quad \rightarrow \quad G(z) = \frac{4}{3} - \sqrt{\frac{2048}{243} \left(\frac{9}{256} - z \right)} + \dots$$

$$\rightarrow \quad z_c = \frac{9}{256} \quad \gamma = \frac{1}{2}$$

Maximal triangulations are shown to have the [topology of the 3-sphere](#).

5 – Gluings of octahedra

These results generalize to the infinite family of bi-pyramids (and connected sums)



$$n_{D-2} \leq 3 + \left(\frac{3}{2} - \frac{1}{2p}\right)n_D$$

Compare with 3D gluings of melonic p -gons $n_{\text{edges}} \leq 3 + \frac{3}{2}n_{\text{tetrahedra}}$

Conclusions

	type of p-gon	D	size	sharp bound	critical exponent
A.	2D p-gon (∞)	2	p	$n_{\text{vertices}} \leq 2 + \frac{p-2}{2} n_{p\text{-gons}}$	-1/2
B.	"melonic" (∞)		even	$n_{D-2} \leq D + \frac{D(D-1)}{4} n_D$ (Gurau)	1/2
		3	even	$n_{\text{edges}} \leq 3 + \frac{3}{2} n_{\text{tetrahedra}}$	
		4	even	$n_{D-2} \leq 4 + 3n_D$	
C.	"necklaces" (∞)	even	even	$n_{D-2} \leq 4 + 2(1 + \frac{1}{p}) n_D$ (Bonzom, Delepoue, Rivasseau, 2015)	-1/2
D.	4-gons	even	4	$n_{D-2} \leq D + (\frac{D(D-1)}{4} - \frac{\alpha(D-1-\alpha)}{4}) n_D$	1/2, -1/2, 1/3
E.	6-gons	3	6	B. or K_{33} : $n_{\text{edges}} \leq 3 + n_{\text{tetra}}$ (Bonzom & L.L, 2015)	1/2
		4	6	Various (L.L & J. Thürigen, IP)	1/2, -1/2, 1/3
F.	Bi-pyramids (∞)	3	8	$n_{\text{edges}} \leq 3 + (\frac{3}{2} - \frac{1}{2p}) n_{\text{tetrahedra}}$ (Bonzom & L.L, 2016)	1/2



Conclusions

- Colored triangulations provide a good framework for combinatorics
- Bijection which generalizes Tutte's bijection for any D -dimensional p -angulation (Bonzom, LL, Rivasseau 2015)

It precisely represents topologies by superposed hyper-maps

Allows to identify and count maximal graphs


Relies tensor models to (complicated) multi-trace matrix models

- First non-melonic/non-quartic random tensor models solved (Bonzom, LL, Thürigen..)
- Maximal configurations exhibit different critical behaviors ($\neq 2D$)
- A lot to be explored!

What next?

1 - Are there building blocks s.t. n_{D-2} is a non linear function of n_D for maximal gluings?

(Possible candidate in D=6)



2 - Can we exhibit building blocks with more interesting maximal maps?

3 - Exact counting of gluings of a single building block (\rightarrow Unicellular maps)

(Harer-Zagier formula ? Chapuy's identity ?)

4 - Gluings of building blocks with colored faces and no internal structure

