

On Serre's conjecture II for groups of type E_7

Philippe Gille

Institut Camille Jordan, Lyon

Geometry Over Nonclosed Fields
Krün, Simons Symposium, April 21, 2016

G -torsors

Let k be a field.

- Let G be an affine algebraic group. A G -torsor is a non-empty affine (right) G -variety X such that the action map

$$X \times_k G \rightarrow X \times_k X, \quad (x, g) \mapsto (x, x.g)$$

is an isomorphism.

- The variety G itself equipped with the right translation is a G -torsor; it is called the trivial G -torsor.
- A G -torsor X is isomorphic to G if and only if $X(k) \neq \emptyset$.
- We denote by $H^1(k, G)$ the set of equivalent classes of G -torsors. It is a pointed set.

G -torsors, II

- We consider an exact sequence of affine algebraic groups
 $1 \rightarrow G_1 \xrightarrow{i} G_2 \xrightarrow{f} G_3 \rightarrow 1.$
- Then we have an exact sequence of pointed sets

$$1 \rightarrow G_1(k) \rightarrow G_2(k) \rightarrow G_3(k) \xrightarrow{\varphi} H^1(k, G_1) \rightarrow H^1(k, G_2) \rightarrow H^1(k, G_3).$$

- The map $\varphi : G_3(k) \rightarrow H^1(k, G_1)$ is called the characteristic map and is defined by $\varphi(g_3) = [f^{-1}(g_3)]$.
- A classical example is that of μ_n where we have the Kummer isomorphism $k^\times / (k^\times)^n \cong H^1(k, \mu_n)$.

Forms

- If G is the automorphism group of an affine k -variety Y (having eventually more structures), there is a way to construct G -torsors.
- Let Y' be a k -form of Y , that is a k -variety such that $Y \times \bar{k} \cong Y' \times \bar{k}$. Then the k -functor

$$R \mapsto \text{Isom}_R(Y_R, Y'_R)$$

is representable by a G -torsor denoted by $\text{Isom}(Y, Y')$.

- In this case $H^1(k, G)$ classifies isomorphism classes of k -forms of Y .
- Artificial exemple : Hilbert 90 theorem, i.e. $H^1(k, \text{GL}_n) = 1$.
- Let q be a regular quadratic form. Then $H^1(k, O(q))$ classifies isometry classes of regular quadratic forms of dimension $\dim(q)$.
- G is the split group of type G_2 . Then $H^1(k, G)$ classifies octonions k -algebras and semisimple k -groups of type G_2 .

Serre's conjecture II

- **Conjecture (1962)** : Assume that k is perfect of cohomological dimension ≤ 2 . Let G be a semisimple simply connected k -group. Then $H^1(k, G) = 1$.
- The assumption on k means that the Galois cohomology groups $H^i(k, A)$ vanish for each finite Galois module A and each $i \geq 3$.
- The assumption on G means first that G is smooth connected with trivial solvable radical and secondly there are no non trivial separable isogeny of algebraic groups $G' \rightarrow G$.
- If k embeds in \mathbb{C} , it is equivalent to say that $G(\mathbb{C})$ is simply connected.
- Examples : $SL_n, Sp_{2n}, Spin_{2n}, \dots$
- Examples of fields : $\mathbb{Q}_p, \mathbb{Q}(i), \mathbb{F}_p(x), \mathbb{C}((x, y)), \mathbb{C}(x, y)$.

Comments on Serre's conjecture II

- The conjecture has strong consequences on the classification on semisimple algebraic groups. More precisely, k -forms of G are classified by their quasi-split form and some Brauer invariants (the Tits class).
- One can dream about stronger conjectures. One is to say that G has enough cohomological invariants, that is one can detect the triviality of a given class of $H^1(k, G)$ by successive Galois cohomology classes.
- This would answer another question by Serre namely whether a G -torsor having a 0-cycle of degree one has a rational point.
- For classical groups, this is known (Bayer/Lenstra, Black) as well for quasi-split groups excepted E_8 (Chernousov, Garibaldi, G.).
- The case of other exceptional groups is widely open.

Approaches to the conjecture

- There are essentially two approaches. The first one is to fix the kind of fields and the other one the kind of groups.
- The first evidence has been the case of p -adic fields done by Kneser but also by Bruhat/Tits in an uniform way.
- For totally imaginary fields except E_8 , this is due to Kneser-Harder in the late 60's. Chernousov solved the E_8 case in 1989.
- The case of global geometric fields is due to Harder in 1975.
- The case of $\mathbb{C}((x, y))$ is due to Colliot-Thélène/Ojanguren/Parimala in 2000.

Geometric fields, index and period

- The case of field of functions of surfaces excluded E_8 was obtained by Colliot-Thélène/G/Parimala in 2003.
- The general case (including E_8) was solved in 2013 by He/de Jong/Starr using deformation techniques of algebraic geometry. This is essentially uniform !
- One common thing between those fields is that they share the “index=period” property for algebras.
- It means that for any central division k -algebra D , its index is the same that its period, that is its exponent in the Brauer group $\text{Br}(k)$.
- For number fields, it is a consequence of the Brauer/Hasse/Noether’s theorem. For fields of functions of surfaces, it is a result by de Jong (2002).

There are other fields

- Let $n \geq 1$ be an integer. Merkurjev constructed in 1991 a field k of cohomological dimension 2 with quaternion algebras Q_1, \dots, Q_n such that $A = Q_1 \otimes Q_2 \cdots \otimes Q_n$ is a division algebra.
- Then A is of period 2 and of index 2^n .
- This field is constructed by an infinite tower of fields.
- Excepted the E_8 case, conjecture II is proven for fields of cohomological dimension ≤ 2 satisfying the “period=index” property for period 2 and 3.
- Next we discuss mainly then exotic fields by the second viewpoint, that is by selecting a class of algebraic groups.

Separable cohomological dimension

- By using Kato's Galois cohomology groups, there is a way to extend the setting also in the positive characteristic. The relevant cohomological dimension is called the separable cohomological dimension.

- Serre refined his conjecture in 1994 as follows.

Refined conjecture Let G be a semisimple simply connected k -group. Assume that k satisfies $\text{scd}_l(k) \leq 2$ for each $l \in S(G)$. Then $H^1(k, G) = 1$.

- The finite set $S(G)$ is called the set of torsion primes of G .

Reduced norms

- Let A be a central simple algebra and consider its special linear group $SL_1(A)$. Then $H^1(k, SL_1(A)) = k^\times / \text{Nrd}(A^\times)$.
- **Theorem** (Suslin, G in positive characteristic) Let p be a prime number. The following are equivalent :
 - (i) $\text{scd}_p(k) \leq 2$;
 - (ii) for each finite separable field extension L/k and each central simple L -algebra B of p -primary index, we have $L^\times = \text{Nrd}(B^\times)$.
- In other words, the surjectivity of reduced norms characterizes the property $\text{scd}(k) \leq 2$.

Classical groups

- Bayer/Parimala proved Serre's conjecture II for classical groups (and G_2 , F_4) in 1995 for perfect fields.
- It was extended by Berhuy/Frings/Tignol in free characteristic in 2008.
- The simplest example is that of $\mathrm{Spin}(q)$. The underlying classification fact is that quadratic forms are classified by their discriminant and their Hasse invariant for a field satisfying $\mathrm{scd}_2(k) \leq 2$.
- The most complicated case is that of outer groups of type A.

Norm groups

- Let X be a k -variety. Its (separable) norm group $N_X(k)$ is the subgroup of k^\times generated by the $N_{L/k}(L^\times)$ for L running over the finite separable finite extensions of k satisfying $X(L) \neq \emptyset$.
- Example 1 : If X has a k -point or more generally a separable 0-cycle of degree one, then $N_X(k) = k^\times$. Note also the property $N_{L/k}(N_{X_L}(L)) \subseteq N_X(k)$ for each finite separable finite extension L/k .
- Example 2 : If $X = \text{SB}(A)$, then $N_X(k) = \text{Nrd}(A^\times)$.
- Example 3 : If $X = \{q = 0\}$ is a smooth projective quadric, then $N_X(k)$ is the image of the spinor norm $\Gamma(q)(k) \rightarrow k^\times$, that is the subgroup of k^\times generated by even products $q(v_1)q(v_2)$ of non zero values of q .

Norm groups, II

The following statement generalizes Suslin's theorem.

- **Proposition** Let G be a k -group satisfying the hypothesis of Serre's conjecture II. Let X be the Borel variety of G . Then $N_X(k) = k^\times$.
- In particular for the quadratic form q we have $N_{X_q}(k) = k^\times$, so that the map $k^\times \rightarrow H^1(k, \text{Spin}(q))$ is trivial.
- More generally, under the conditions of conjecture II, we are interested in the image of the map $f_* : H^1(k, \mu_n) \rightarrow H^1(k, G)$ for $f : \mu_n \rightarrow G$ a k -homomorphism.
- If $f(\mu_n)$ is central, then using the "normprinciple" and the proposition, we have that f_* is the trivial map.


The non-central case

Again G satisfies the hypothesis of conjecture II. We are given $f : \mu_n \rightarrow G$ and denote by $H = Z_G(f)$ its centralizer.

- By Steinberg-Springer's connectedness theorem, H is connected so that H is reductive.
- The map $f_* : H^1(k, \mu_n) \rightarrow H^1(k, G)$ factorizes then by $H^1(k, H)$ and $f(\mu_n)$ is central in H . The vanishing of f_* boils then to the vanishing of $H^1(k, \mu_n) \rightarrow H^1(k, H)$.
- The problem is that H can be semisimple not simply connected and even not semisimple. In the split case, conjugacy classes of subgroups μ_n of G are classified by means of Kac coordinates.
- It provides then a precise combinatorial classification of the centralizers H occurring there. A consequence is that the quasi-split form H^{qs} of H contains a maximal quasi-trivial torus.
- This is enough to ensure the vanishing of $H^1(k, \mu_n) \rightarrow H^1(k, H)$. To summarize the map $f_* : H^1(k, \mu_n) \rightarrow H^1(k, G)$ is trivial.

Groups of type E_7


We deal with a semisimple algebraic group G of type E_7



- Let G_0 be the split form of G . It carries a linear representation $G_0 \hookrightarrow \mathrm{GL}_{56}$.
- This representation descends to a representation $G \hookrightarrow \mathrm{GL}_7(A)$ where A is central simple algebra of degree 8 and of period 2.
- A is called the Tits algebra of G .
- The algebra A is Brauer equivalent to a tensor product of 4 quaternion algebras. If $\mathrm{scd}_2(k) \leq 2$, a result by Sivatski implies that A is a product of 3 quaternion algebras.

The main result

Again G is a semisimple simply connected k -group of type



Theorem. We assume that $\text{scd}_2(k) \leq 2$ and that $\text{scd}_3(k) \leq 2$.

- (1) We have $H^1(k, G) = 1$ if A is of index ≤ 4 .
- (2) Let $[z] \in H^1(k, G)$. Then $[z] = 1$ if and only if there exists a quaternion algebra Q and embeddings $SL_1(Q) \rightarrow G$ and $SL_1(Q) \rightarrow {}_z G$ which are geometrically "coroot embeddings".
- (3) If A is of index 1 (resp. 2, 4) then G is split (resp. has k -rang 4, 1) and admits a parabolic k -subgroup of type \emptyset (resp. $\{2, 5, 7\}$, $\{2, 3, 4, 5, 6, 7\}$).

Concluding remarks

- Remarks : If A is of index 8, then G is anisotropic. Also there exists an embedding $SL_1(Q) \rightarrow G$ but we do not know how to show that $SL_1(Q)$ embeds in the twisted k -group ${}_z G$.
- It follows that groups of type E_7 are classified by their Tits algebra in index 1, 2, 4 for fields satisfying $\text{scd}_2(k) \leq 2$ and $\text{scd}_3(k) \leq 2$.
- We do not know whether all triquaternion algebras occur as Tits algebra of groups of some group of type E_7 . A partial result is that biquaternion algebras occur (Quéguiner-Mathieu).
- What about exceptional groups of type E_6, E_8 ?
- There is no Sivatski's theorem for odd primes, this is a first difficulty in type E_6 for the relevant algebras of index 27 and period 3. Also groups of type E_6 with Tits algebra of index 9 are always anisotropic.

A statement for E_8

For simplicity, we assume that k is of characteristic zero.

- Bogolomov conjectured that $\text{cd}(F^{ab}) = 1$ for any field F (possibly containing an algebraically closed field). It would imply that a F -group of type E_8 is split by an abelian extension of degree $2^a 3^b 5^c$. This is an open question which implies conjecture II for E_8 .
- Under the hypothesis of conjecture II, we know that a k -torsor X under E_8 has a 0-cycle of degree one. So answering Serre's question on 0-cycles of degree one for torsors would conjecture II for E_8 . We end with the following statement.
- **Theorem.** Assume that $\text{cd}_2(k) \leq 2$, $\text{cd}_3(k) \leq 2$ and $\text{cd}_5(k) \leq 2$. Let $[z] \in H^1(k, E_8)$ (where E_8 stands for the Chevally group of type E_8). Then $[z] = 1$ if and only if the group ${}_z E_8(k)$ has torsion.

Danke schön.